

# Spectroscopic Studies of Electrochemically Generated Cation Radical of Chlorophyll a in Organic Solvents

著者	KURAWAKI Junichi, KUSUMOTO Yoshihumi
journal or publication title	鹿児島大学理学部紀要=Reports of the Faculty of Science, Kagoshima University
volume	30
page range	17-25
URL	<a href="http://hdl.handle.net/10232/6220">http://hdl.handle.net/10232/6220</a>

# Spectroscopic Studies of Electrochemically Generated Cation Radical of Chlorophyll a in Organic Solvents

著者	KURAWAKI Junichi, KUSUMOTO Yoshihumi
journal or publication title	鹿児島大学理学部紀要=Reports of the Faculty of Science, Kagoshima University
volume	30
page range	17-25
URL	<a href="http://hdl.handle.net/10232/00010005">http://hdl.handle.net/10232/00010005</a>

# Spectroscopic Studies of Electrochemically Generated Cation Radical of Chlorophyll a in Organic Solvents

Junichi KURAWAKI and Yoshihumi KUSUMOTO

(Received July 8, 1997)

**Keywords :** chlorophyll cation radical, electrolysis, spectroscopy, Raman spectra

## Abstract

Spectroscopic studies of electrochemically generated cation radical of chlorophyll a (Chl) were carried out in various organic solvents. Absorption and resonance Raman (RR) spectra of Chl cation radical ( $\text{Chl}^+$ ) exhibited remarkable spectral changes with oxidation. The major prominent changes of Raman bands observed upon oxidation of Chl are a large upshift of the  $\text{C}_9$  keto-carbonyl stretching on ring V ( $\text{C}_9=\text{O}$ ), downshifts of the coordination-state marker bands, and intensity and frequency changes of Raman bands in the  $\text{C}_a\text{C}_m$  and  $\text{C}_a\text{N}$  stretching modes. The large upshift of the  $\text{C}_9=\text{O}$  stretching mode indicates that there is a decrease in the conjugation between  $\text{C}_9=\text{O}$  and the macrocycle following one-electron oxidation. We could find some oxidation-state marker Raman bands in all organic solvents investigated in this study.

## Introduction

The conformational changes of chlorophylls in photosystem (PS) I and photosystem (PS) II serve to control electron transfer by changing the interaction of the chromophore with the surrounding protein and altering the relative distances and orientations between donor and acceptor species (1, 2). As a result, a chlorophyll a (Chl) / Chl cation radical ( $\text{Chl}^+$ ) radical pair is formed in the charge separation process in both PS I and PS II (3). Accordingly, it is important to investigate the structural and electronic properties of  $\text{Chl}^+$  which influence the process of charge separation. Immense effort has been so far expended toward characterizing the spectroscopic properties of  $\text{Chl}^+$  in addition to neutral Chl through absorption, magnetic resonance and vibrational spectroscopies (4-6). In spite of this effort, little information is available on the consequence of the precise structures for  $\text{Chl}^+$ . Resonance

Raman (RR) spectroscopy is, however, a powerful tool for studying the molecular structures of the primary donor and cation and anion radicals produced by electron transfer. Only a few RR spectra of cation and anion radicals of Chl have been reported so far (6-8).

In this paper, therefore, we report spectroscopic results of electrochemically generated  $\text{Chl}^+$  in various organic solvents by means of absorption, fluorescence and Raman spectroscopies. A purpose of our research is to elucidate the vibrational structures of  $\text{Chl}^+$  and its models operating as the primary donor in PS I and PS II.

## Experimental

Commercial Chl (Wako, > 95 %) was used as received and its purity was checked by measuring its absorption spectrum in ethanol. Prior to use, Chl was dried by repetitive codistillation from dichloromethane ( $\text{CH}_2\text{Cl}_2$ ). All organic solvents

used in this work were spectroscopic grade or HPLC one, and were dried by vacuum distillation, and then stored over activated 3 Å molecular sieves. As a supporting electrolyte, tetrabutylammonium perchlorate (TBAP) was used. TBAP was purified by recrystallization from ethyl acetate and dried under vacuum at 100 °C.

Absorption and fluorescence spectra of neutral Chl and  $\text{Chl}^+$  were recorded on a Hitachi 228 spectrophotometer and a Hitachi 850 spectrofluorometer, respectively. Fluorescence lifetimes of  $\text{Chl}^+$  were measured with a Horiba NAES-1100 time-resolved spectrophotometer equipped with a single-photon counting system. ESR spectra of  $\text{Chl}^+$  were performed in an ESR spectrometer (JEOL JM-ME-3X). Raman spectra were recorded on a Jasco R-800 laser-Raman spectrophotometer under the conditions pre-resonating the Soret absorption band by using the 441.6 nm of a He-Cd laser (Kimmon Electric Co., Ltd., CD4601R). Experimental conditions were the same for all Raman experiments and are as follows: resolution 5  $\text{cm}^{-1}$ ; laser power <10 mW at the sample point. All measurements were conducted at 25 °C.

Generation of  $\text{Chl}^+$  and its spectroscopic meas-

urements were performed in a home-made spectro-electrochemical cell (optical path length = 1 and 5 mm) containing a platinum mesh working electrode. Controlled-potential electrogeneration of  $\text{Chl}^+$  was performed at the platinum electrode with a Hokuto Denko HA-301 potentiostat. The concentration of Chl in the sample solutions was adjusted between  $1 \times 10^{-4}$  and  $5 \times 10^{-4}$  M ( $1 \text{ M} = 1 \text{ moldm}^{-3}$ ). The concentration of TBAP was kept at 0.1 M. In order to test the performance of this spectroelectrochemical cell used in this work, we measured representative absorption spectra of zinc tetraphenylporphyrin (ZnTPP) in  $\text{CH}_2\text{Cl}_2$  during electrolysis at 0.8 V and ESR spectrum of the species produced at a final stage of electrolysis. The spectral behavior shown in Fig. 1 and *g*-value (2.002) obtained from ESR spectrum were very close to those reported by Fajer et al. (3); this result indicates that ZnTPP cation radical is produced via one-electron oxidation and this spectroelectrochemical cell can operate successfully.

## Results and Discussion

### Absorption spectroscopy of Chl and $\text{Chl}^+$

Absorption spectra of Chl are dominated by intense  $\pi - \pi^*$  transitions which occur in the visible




Fig. 1. Absorption spectral change of ZnTPP during electrolysis in dichloromethane ( $\text{CH}_2\text{Cl}_2$ ). 1; absorption spectrum of neutral ZnTPP, 4; absorption spectrum of ZnTPP cation radical. Insert is molecular structure of ZnTPP.

region. Fig.2 shows representative absorption spectra at various stages of electrolysis for three organic solvents. For example, the absorption spectrum of neutral Chl in  $\text{CH}_2\text{Cl}_2$  solution shows maxima at 432 nm in the Soret region and at 666 nm in the red region (Fig. 2 (A)). The absorption peaks at 432 and 666 nm correspond to the  $B_x$  and  $Q_y$  transitions, respectively. Upon electrolysis at 0.85 V, the absorbance of these two bands gradually decreased and the Soret absorption maximum shifted

from 432 to 415 nm. In addition, a new strong absorption bands at 415 nm and broad bands around 510, 700 and 815 nm appeared. As shown in Fig.2 (A), the fact that well defined isosbestic points were observed at 426, 446, 609, 640 and 683 nm indicates the smooth course of this reaction. The ESR spectrum of the species produced at a final stage of electrolysis in  $\text{CH}_2\text{Cl}_2$  solution (Fig.2 (A), insert) showed only a single line at  $g=2.004$ ; this species can be assigned to a  $\pi$ -cation radical of Chl ( $\text{Chl}^+$ ).




Fig. 2. Absorption spectral changes of Chl with oxidation in (A)  $\text{CH}_2\text{Cl}_2$ , (B) acetone and (C) methanol solutions. The insert figures in (A) and (C) show ESR spectra of  $\text{Chl}^+$  at a final stage of electrolysis at 0.85 V.

Similar results were also obtained for the other organic solvents such as acetone (Fig. 2 (B)), methanol (Fig. 2 (C)), tetrahydrofuran (THF) and acetonitrile. Therefore, it can be concluded that  $\pi$ -cation radical of Chl ( $\text{Chl}^+$ ) is efficiently produced via one-electron oxidation of Chl.

### Fluorescence spectroscopy of Chl and $\text{Chl}^+$

Typical fluorescence spectra of neutral Chl and  $\text{Chl}^+$  excited at 415 nm are shown in Fig. 3. The fluorescence peak of neutral Chl excited at 415 nm is located at 674 nm for  $\text{CH}_2\text{Cl}_2$  (Fig. 3 (A)) and at 677 nm for methanol (Fig. 3 (B)). The fluorescence peak of  $\text{Chl}^+$  formed by oxidation of Chl for  $\text{CH}_2\text{Cl}_2$  system shifted to 676 nm, while that for methanol system exhibited a blue-shift by *ca.* 6 nm.

A typical fluorescence decay curve of  $\text{Chl}^+$  for the emission at 680 nm upon excitation at 415 nm is shown in Fig. 4 for acetone system. The decay curve shows a single-exponential decay with a lifetime of 8.5 ns. The lifetime of neutral Chl in acetone solution has been reported to be 6.5 ns (9). The lifetime of  $\text{Chl}^+$  is longer than that of Chl, suggesting the electronic stability of the excited  $\text{Chl}^+$  in acetone solution. In this study, we first reported the life-

times of the excited  $\text{Chl}^+$  in organic solvents.

### Raman spectroscopy of Chl and $\text{Chl}^+$

RR spectra of neutral Chl in various organic solvents have been measured to clarify the states of Chl, *e. g.*, axial-coordinate states, self-aggregation and hydrogen bonding between carbonyl oxygen atom and solvent molecule (10-14). The Raman bands in the region of 1750-1000  $\text{cm}^{-1}$  for neutral Chl have been assigned to as follows (10, 15): bands in the higher frequency region (1750-1650  $\text{cm}^{-1}$ ) can be assigned to the keto-carbonyl stretching ( $\text{C}_9=\text{O}$ ), those in the 1630-1520  $\text{cm}^{-1}$  region can be related to both central 16-membered-ring vibrations and  $\text{C}_a\text{C}_m$  or  $\text{C}_b\text{C}_b$  stretching, and those in the region of 1500-1000  $\text{cm}^{-1}$  can be assigned to  $\text{C}_a\text{C}_m$ ,  $\text{C}_m\text{C}_{10}$  and  $\text{C}_a\text{N}$  stretching modes.

It is expected that the RR spectral changes that occur with oxidation of Chl may show a significant effect on the magnitude of the observed band shifts because of the lower symmetry of Chl. Thus, we measured the RR spectra of  $\text{Chl}^+$  formed by oxidation of Chl in organic solvents. For all solvents investigated here, oxidation of Chl resulted in a number of RR spectral changes and almost similar




Fig. 3. Fluorescence spectra of Chl and  $\text{Chl}^+$  in (A)  $\text{CH}_2\text{Cl}_2$  and (B) methanol solvent systems. Excitation wavelength: 415 nm.




Fig. 4. A fluorescence decay curve of  $\text{Chl}^+$  in acetone solution. Excitation wavelength: 415 nm. Monitored wavelength: 680 nm.

spectral behaviors were obtained except for THF system. In this section, therefore, we present typical RR spectra obtained from  $\text{CH}_2\text{Cl}_2$  and THF solvent systems, as shown in Figs. 5 and 6.

For  $\text{CH}_2\text{Cl}_2$  system in Fig. 5, a Raman band at  $1695\text{ cm}^{-1}$  (neutral Chl) assignable to the  $\text{C}_9$  keto-carbonyl stretching ( $\text{C}_9=\text{O}$ ) mode is strongly affected with oxidation and shows a frequency upshift by  $15\text{ cm}^{-1}$ . Peaks at  $1612$  and  $1189\text{ cm}^{-1}$  observed for Chl are downshifted to  $1596$  and  $1184\text{ cm}^{-1}$ , respectively. In addition, loss of intensity is observed for bands at  $1532$ ,  $1337$ ,  $1293$ ,  $1189$  and  $1051\text{ cm}^{-1}$ .

As can be seen from Fig. 6, the RR spectra of  $\text{Chl}^+$  obtained from THF solution were more strongly affected than those from  $\text{CH}_2\text{Cl}_2$ . A Raman band at  $1695\text{ cm}^{-1}$  observed for Chl is upshifted to  $1710\text{ cm}^{-1}$  with oxidation. It can be seen that there are apparent frequency downshifts of  $1596 \rightarrow 1579$ ,  $1180 \rightarrow 1175$  and  $1344 \rightarrow 1332\text{ cm}^{-1}$  with oxidation, while the frequency upshifts of  $1546 \rightarrow 1551$  and  $1289 \rightarrow 1293\text{ cm}^{-1}$  are also observed. Moreover, Fig.

6 shows that a significant decrease in intensity is observed for bands at  $1520$ ,  $1430$ ,  $1344$ ,  $1289$ ,  $1260$ ,  $1180$  and  $1045\text{ cm}^{-1}$  and new bands appear at  $1486$ ,  $1455$ ,  $1230$ ,  $1065$  and  $1032\text{ cm}^{-1}$ . For the purpose of the identification for the prominent RR bands obtained in this work, the frequencies and assignments for  $\text{CH}_2\text{Cl}_2$  and THF solvent systems are summarized in Table 1.

RR bands at  $1695$  ( $\text{CH}_2\text{Cl}_2$ ) and  $1690\text{ cm}^{-1}$  (THF) observed for neutral Chl were the most strongly affected and showed upshifts in frequency by  $15$ - $25\text{ cm}^{-1}$  with oxidation. These bands are assigned to the  $\text{C}_9=\text{O}$  stretching mode (10, 15). Heald and Cotton (7) have measured RR spectra of pyrochlorophyll a cation radical in  $\text{CH}_2\text{Cl}_2$  solution and reported that oxidation of this pigment to form the cation radical results in a decrease in intensity of  $1681\text{ cm}^{-1}$  band and the appearance of a new band at  $1708\text{ cm}^{-1}$ . This behavior is very similar to that observed for  $\text{Chl}^+$  in this work. Thus, this fact suggests that the origin of the Raman bands at  $1710$  ( $\text{CH}_2\text{Cl}_2$ ) and  $1715\text{ cm}^{-1}$  (THF) observed for  $\text{Chl}^+$  is




Fig. 5. Resonance Raman spectra of Chl (top) and Chl<sup>+</sup>· (bottom) in CH<sub>2</sub>Cl<sub>2</sub> solvent system.

not the ester group but the C<sub>9</sub> keto group. The large band shift observed for the C<sub>9</sub>=O stretching mode may reflect a localization of the major electron density charges at the isocyclic ring. It can be concluded that Chl<sup>+</sup> must be in the keto form and not the enol form.

The Raman bands at 1612, 1558 and 1532 cm<sup>-1</sup> (CH<sub>2</sub>Cl<sub>2</sub>) and 1596, 1546 and 1520 cm<sup>-1</sup> (THF) for Chl are known to be sensitive to the coordination-state of Chl and are assigned to C<sub>a</sub>C<sub>m</sub> or C<sub>9</sub>C<sub>9</sub> mode (10, 12, 13). A frequency downshift from 1612 to 1596 cm<sup>-1</sup> and the loss of intensity at 1532 cm<sup>-1</sup> which may accompany the frequency upshift by 6 cm<sup>-1</sup> indicate that oxidation of Chl results in reduced electron density on the methine bridge of macrocycle.

In the region of 1500 - 1200 cm<sup>-1</sup>, the 1337 - and 1293-cm<sup>-1</sup> bands for Chl in CH<sub>2</sub>Cl<sub>2</sub> solution, which are assigned to the C<sub>a</sub>N stretching mode, exhibit a decrease in intensity. A new Raman band at 1360 cm<sup>-1</sup> appears in this solvent; this band can be tentatively assigned to the C<sub>a</sub>C<sub>b</sub> mode. On the contrary, in the case of THF system, the different spectral behaviors could be observed. Loss of intensity is observed at 1430 (C<sub>a</sub>C<sub>m</sub>) and 1326 cm<sup>-1</sup> (C<sub>a</sub>N) and the frequency of the 1344-cm<sup>-1</sup> band (C<sub>a</sub>N) is downshifted to 1332 cm<sup>-1</sup> with Chl oxidation. These results suggest that oxidation of Chl induces the change of substantial spin density at the C<sub>a</sub> position of methine carbons of Chl<sup>+</sup>, resulting in the changes of the C<sub>a</sub>C<sub>m</sub> and C<sub>a</sub>C<sub>b</sub> mode bands as well as bands with C<sub>a</sub>N mode.




Fig. 6. Resonance Raman spectra of Chl (top) and Chl<sup>+</sup> (bottom) in THF solvent system.

In the 1200 - 1000  $\text{cm}^{-1}$  region, 1189- ( $\text{CH}_2\text{Cl}_2$ ) and 1180- $\text{cm}^{-1}$  (THF) bands observed for Chl are, respectively, downshifted to 1184 and 1175  $\text{cm}^{-1}$  and also exhibit a decrease in intensity with oxidation. Although the Raman spectrum of neutral Chl in this region usually contains predominantly  $\text{C}_a\text{N}$  modes according to normal mode assignments (10, 15), Boldt et al. (15) have assigned one band in this region to the  $\text{C}_m\text{C}_{10}$  stretching and reported that this band at 1190  $\text{cm}^{-1}$  shifts to 1185  $\text{cm}^{-1}$ . The 1184- and 1175-  $\text{cm}^{-1}$  bands observed in this work, therefore, reflect a change in the electron density within the isocyclic ring with oxidation. In addition, the 1051- ( $\text{CH}_2\text{Cl}_2$ ) and 1045-  $\text{cm}^{-1}$  (THF) bands ( $\text{C}_s\text{N}$  mode) of Chl disappear, while in the case of THF

system new Raman bands at 1065 and 1032  $\text{cm}^{-1}$  appear upon oxidation; the former band is tentatively assigned to the  $\text{C}_a\text{C}_m$  stretching and the latter one is assigned to the  $\text{C}_m\text{C}_a\text{C}_b$  vibration. This difference in two solvent systems may be ascribed to the difference of the coordination-state of Chl; a THF solvent molecule can bind to the central Mg atom of Chl as the fifth-ligand but in the case of  $\text{CH}_2\text{Cl}_2$  solvent system the coordination number of Chl is four. Thus, the plane-deformation of Chl for THF system may be more strongly induced with oxidation.

Judging from the RR results described above, it can be concluded that oxidation of Chl results in the formation of Chl<sup>+</sup> and induces significant changes

Table 1. Frequencies ( $\text{cm}^{-1}$ ) of the prominent Raman bands of Chl and  $\text{Chl}^+$  in  $\text{CH}_2\text{Cl}_2$  and THF solvent systems

$\text{CH}_2\text{Cl}_2$		THF		Assignment
Chl	$\text{Chl}^+$	Chl	$\text{Chl}^+$	
1695	1710	1690	1715	$\text{C}_9=\text{O}$
	1616	1615	1617	$\text{C}_a\text{C}_m$
1612	1596	1596	1579	$\text{C}_a\text{C}_m$
1558	1560	1546	1551	$\text{C}_b\text{C}_b$
1532	1538	1520	1521	$\text{C}_b\text{C}_b$
			1486	$\text{C}_a\text{C}_m$
			1455	$\text{CH}_3$
1433	1432	1430		
1396	1398	1386	1385	$\text{C}_a\text{N}, \text{C}_a\text{C}_b$
1384	1381	1375	1374	$\text{C}_a\text{C}_b$
	1360			$\text{C}_a\text{C}_b$
1349	1348	1344	1332	$\text{C}_a\text{N}$
1337	1338	1326		$\text{C}_a\text{N}$
1293	1293	1289	1293	$\text{C}_a\text{N}$
1265	1267	1260		$\text{C}_a\text{N}$
1189	1184	1180	1175	$\text{C}_m\text{C}_{10}$
1155	1159	1155	1154	$\text{C}_a\text{N}, \text{C}_a\text{C}_m\text{C}_a$
			1065	$\text{C}_a\text{C}_m$
1051		1045		$\text{C}_a\text{N}$
			1032	$\text{C}_m\text{C}_a\text{C}_b$

in the electron density at the  $\text{C}_9=\text{O}$  bond and in the electron spin density of the macrocyclic  $\pi$ -system. In this study we could observe some  $\text{Chl}^+$ -state marker Raman bands in all organic solvents investigated and conclude that Chl may undergo significant conformational changes with loss of an electron from the  $\text{C}_9$  keto-carbonyl group. Finally, the RR results suggest that the  $\text{C}_9=\text{O}$  group of Chl plays an important role in electron transfer in PS I and PS II through its interaction with the proteins.

## References

1. M. Y. Okamura, G. Feher and N. Nelson, "Energy Conversion by Plants and Bacteria", Ed. by Govindjee, Academic Press, New York (1982), Vol. 1, P.197.
2. H. J. D. Blanken and A. Hoff, *J. Biochim. Biophys. Acta*, **724**, 52 (1983).
3. J. Fajer, D. C. Borg, A. Forman, D. Dolphin and R. H. Felton, *J. Am. Chem. Soc.*, **92**, 3451 (1970).
4. M. S. Davis, A. Forman and J. Fajer, *Proc. Natl. Acad. Sci. U. S. A.*, **76**, 4170 (1979).

5. J. D. Petke, G. M. Maggiora, L. L. Shipman and R. E. Christoffersen, *Photochem. Photobiol.*, **31**, 243 (1980).
6. R. L. Heald, P. M. Callahan and T. M. Cotton, *J. Phys. Chem.*, **92**, 4820 (1988).
7. R. L. Heald and T. M. Cotton, *J. Phys. Chem.*, **94**, 3968 (1990).
8. J. -H. Perng and D. F. Bocian, *J. Phys. Chem.*, **96**, 10234 (1992).
9. A. Agostiano, K. A. Butcher, M. S. Showell, A. J. Gotch and F. K. Fong, *Chem. Phys. Lett.*, **137**, 37, (1987).
10. M. Lutz, "Advances in Infrared and Raman Spectroscopy", Ed. by R. J. H. Clark and R. E. Hester, Wiley, New York (1984), Vol. 11, P.211; *J. Raman Spectry*, **2**, 497 (1974); *Biochim. Biophys. Acta*, **460**, 408 (1977).
11. T. M. Cotton and R. P. Van Duyne, *J. Am. Chem. Soc.*, **103**, 6020 (1981).
12. M. Fujiwara and M. Tasumi, *J. Phys. Chem.*, **90**, 250 (1986).
13. M. Tasumi and M. Fujiwara, "Spectroscopy of Inorganic-based Materials", Ed. by R. J. H. Clark and R. E. Hester, Wiley, New York, (1987), Chap. 6.
14. J. Kurawaki and Y. Kusumoto, *Chem. Phys. Lett.*, **158**, 495 (1989).
15. N. J. Boldt, R. J. Donohoe, R. R. Birge and D. F. Bocian, *J. Am. Chem. Soc.*, **109**, 2284 (1987).