

Changings of Polycyclic Aromatic Hydrocarbons in Aquatic Organisms and Sediment at Guimaras after Oil Spill

著者	Uno Seiichi, Kokushi Emiko, Pahila Ida G., Koyama Jiro
journal or publication title	鹿児島大学水産学部紀要=Memoirs of Faculty of Fisheries Kagoshima University
volume	特別号(2010)
page range	63-66
URL	http://hdl.handle.net/10232/9378

Changings of Polycyclic Aromatic Hydrocarbons in Aquatic Organisms and Sediment at Guimaras after Oil Spill

Seiichi Uno¹, Emiko Kokushi², Ida G. Pahila³, and Jiro Koyama¹

Key words: Oil spill, polycyclic aromatic hydrocarbons, shellfish

Abstract

Following the oil spill accident of the Solar I tanker in 2006 off the coast of Guimaras Island in the Philippines, polycyclic aromatic hydrocarbons (PAHs) in shellfish was investigated at Luzaran in Guimaras Islands, which was heavily polluted with spilled oil, from September 2006 (1 month after the accident) to June 2008. Although concentrations of total PAHs extremely decreased during 3 months after oil spill, those slowly changed in the following period. The relative high molecular weight PAHs such as benzo(b)fluoranthene, benzo(k)fluoranthene, benzo(a)pyrene, dibenz(a,h)-anthracene, indeno(1,2,3-cd)pyrene, and benzo(g,h,i)perylene also tended to be residue in shellfish.

On August 11, 2006, the Solar I tanker was hit by hard rain-storm and sank off the coast of Guimaras Island in the Philippines. About 350,000 tons of heavy fuel oil was spilled from the tanker polluting about 200 km of shoreline around Guimaras. Over 450 ha of huge protected mangrove areas with complex topography at Guimaras was also contaminated¹⁾.

In general, spilled oil degrades with several factors such as evaporation, diffusion, photooxidation, and biodegradation¹⁻⁵⁾. In tropical areas, because of strong sunshine, high temperature, and frequent storms, spilled oil is expected to degrade faster than in the temperate and boreal areas although it is still poorly studied. Since polycyclic aromatic hydrocarbons (PAHs) are persistent in the environment⁵⁻⁷⁾, their toxicities to aquatic organisms have been examined⁸⁻¹¹⁾. The acute toxicity of PAHs to fish generally induces narcosis, which is a reversible anesthetic effect¹²⁾. In addition, acute exposures with early life stages PAHs resulted in developmental abnormalities¹³⁾. When an oil spill occurs in a water system, the concentrations of PAHs are significantly high in water and could have significant threats and several negative impacts on the surrounding

aquatic organisms.

Uno et al. reported the initial impacts of the Solar I, and especially the accumulations of PAHs were examined in some species of fish, shellfish, and crabs immediately after the spill and 1 month later¹⁾. In the present study, the changing of PAH concentrations in shellfish, was caught at the southern tip of Guimaras Island, were investigated for about 1 year immediately after oil spill.

Materials and Methods

Collection of organisms

The five investigations were conducted at Luzaran the southern tip of Guimaras Island, from 2006 to 2008 (September and November 2006, March and September 2007, and June 2008). Luzaran has the huge mangrove forests, and has many species of aquatic organisms. Fig. 1 shows the location of Luzaran on the map. Four species of shellfish, *Modiolus* sp., *Pinctada* sp., *Clypeomorus* sp., and oyster (its taxon has not been identified), were collected in each investigation. After the collections, all samples were stored at -20°C until anal-

¹ 鹿児島大学水産学部海洋資源環境教育研究センター (Education and Research Center for Marine Resources and Environment, Faculty of Fisheries, Kagoshima University, 50-20 Shimoarata 4-Chome, Kagoshima 890-0056, Japan)

² 鹿児島大学大学院連合農学研究科 (The United Graduate School of Agricultural Science, Kagoshima University, 50-20 Shimoarata 4-Chome, Kagoshima 890-0056, Japan)

³ Department of Chemistry, College of Arts and Sciences, University of the Philippines Visayas, Miagao, 5023 Iloilo, Philippines

* Corresponding author, E-mail: uno@fish.kagoshima-u.ac.jp

ysis.

Analysis of PAHs in shellfish

The analysis of PAHs was conducted according to the previous method procedures¹⁾. The target PAHs were as following: naphthalene, acenaphthylene, acenaphthene, fluorene, dibenzothiophene, phenanthrene, anthracene, fluoranthene, benz(a)anthracene, chrysene, benzo(b)fluoranthene, benzo(k)fluoranthene, benzo(a)pyrene, perylene, dibenz(a,h)anthracene, indeno(1,2,3-cd)pyrene, and benzo(g,h,i)perylene. The index total PAHs were calculated as the sum of the quantity of all these compounds.

Results and Discussion

Concentrations of mean total PAHs extremely decreased from September to November 2006, but those slowly changed in the following period (Fig. 2). Especially, shellfish collected at September 2006 could accumulate PAHs directly from water, because it had been just one month after oil spill, and much oil components were suspended in sea water yet. On the other hand, the slow change of PAHs from November 2006 to June 2008 suggested that the pollutions of PAHs had continued around Luzaran since oil spill. In general, when oil spill happens on the sea, the oil components burrow into sediments and are residue with little degradation in sediments for long period¹⁴⁾. The PAHs in shellfish at Luzaran from November 2006 to June 2008 could be derived the oil components, which




Fig. 1. Locations of the sampling site, Luzaran, where shellfish was collected.

were eluted gradually from sediment.

Figure 3 shows the example of the changes for individual PAH concentrations in shellfish (*Clypeomorus* sp.) at individual investigations. Concentration of chrysene was the highest among PAHs in *Clypeomorus* sp. for about six months after spill. However, that was rapidly lowered after September 2007. In addition, concentrations of relatively high molecular weight PAHs such as benzo(b)fluoranthene, benzo(k)fluoranthene, benzo(a)pyrene, dibenz(a,h)anthracene, indeno(1,2,3-cd)pyrene, and benzo(g,h,i)perylene, changed little in *Clypeomorus* sp. Those PAHs could be residue in sediments with little degradation, and/or be more slowly metabolized by shellfish than low molecular weight PAHs. Joint FAO/WHO Expert Committee on Food Additives (JECFA) indicated that those high molecular weight PAHs are mutagenic or carcinogenic compounds¹⁵⁾. The continuous investigation should be conducted at Guimaras Island until the residues of high molecular weight PAHs with mutagenicity and carcinogenicity will be low in shellfish.

Acknowledgements

We thank the people of Guimaras for their great cooperation, and the staff of the University of the Philippines for their help with this investigation. This study was supported by the Asian CORE Program of the Japan Society for the Promotion of Science (JSPS).

References

- 1) Seiichi U., J. Koyama, E. Kokushi, H. Monteclaro, S. Santander, J.O. Cheikyula, S. Miki, N. Añasco, I.G. Pahila, H.S. Taberna Jr., and T. Matsuoka. Monitoring of PAHs and alkylated PAHs in




Fig. 2. Changings of total PAHs in 4 species of shellfish from September 2006 to June 2008. A bar indicates mean concentration of PAHs in shellfish.




Fig. 3. Concentrations of individual PAHs in *Clypeomorus* sp. at September and November 2006, March and September 2007, and June 2008 (ng/g-wet weight). The abbreviations are as follows: Nap, naphthalene; Acy, acenaphthylene; Ace, acenaphthene; Fle, fluorine; Db, dibenzothio-phenylene; Phe, phenanthrene; Ant, anthracene; Fla, fluoranthene; Pyr, pyrene; BaA, benzo(a)anthracene; Chr, chrysene; BbF, benzo(b)fluoran-thene; BkF, benzo(k)fluoranthene; BaP, benzo(a)pyrene; Per, perylene; BP, benzo(g,h,i)perylene DA, dibenz(a,h)anthracene; and IP, indeno(1,2,3-cd)pyrene.

- aquatic organisms after 1 month from the *Solar I* oil spill off the coast of Guimaras Island, Philippines. *Environ. Monit. Assess.* in print.
- 2) Hayes, M.O., and J. Michel (1999). Factors determining the long-term persistence of Exxon Valdez oil in gravel beaches. *Mar. Pollut. Bull.*, **38**: 92-101.
 - 3) Michel, J. and M.O. Hayes (1999). Weathering patterns of oil residues eight years after the Exxon Valdez oil spill. *Mar. Pollut. Bull.*, **38**: 855-863
 - 4) Laffon, B., T. Rábade, E. Pásaro, and J. Méndez (2006). Monitoring of the impact of Prestige oil spill on *Mytilus galloprovincialis* from Galician coast. *Environ. Int.*, **32**: 342-348.
 - 5) Owens, E.H., E. Taylor, and B. Humphrey (2008). The persistence and character of stranded oil on coarsesediment beaches. *Mar. Pollut. Bull.*, **56**: 14-26.
 - 6) Taylor, E. and D. Reimer (2008). Oil persistence on beaches in Prince William Sound—A review of SCAT surveys conducted from 1989 to 2002. *Mar. Pollut. Bull.*, **56**: 458-474.
 - 7) Wang, Z., K. Li, P. Lambert, and C. Yang (2007). Identification, characterization and quantitation of pyrogenic polycyclic aromatic hydrocarbons and other organic compounds in tire fire products. *J. Chromatogr. A*, **1139**: 14-26.
 - 8) Wetzel, D.L., and E.S. van Vleet (2003). Persistence of petroleum hydrocarbon contamination in sediments of the canals in Venice, Italy: 1995 and 1998. *Mar. Pollut. Bull.*, **46**: 1015-1023.
 - 9) Barron, M. G. (1990). Bioconcentration. *Environ. Sci. Tec.*, **24**: 1612-1618.
 - 10) Incardona, J.P., T.K. Collier, and N.L. Scholz (2004). Defects in cardiac function precede morphological abnormalities in fish embryos exposed to polycyclic aromatic hydrocarbons. *Toxicol. Appl. Pharmacol.*, **196**: 191-205.
 - 11) Reynaud, S. and P. Deschaux (2006). The effects of polycyclic aromatic hydrocarbons on the immune system of fish: A review. *Aquat. Toxicol.*, **77**: 229-238.
 - 12) Francioni, E., L.R. de Wagener, A.L. Scofield, M.H. Depledge, B. Cavalier, C.B. Sette, L. Carvalhosa, C. Lozinsky, and R. Mariath (2007). Polycyclic aromatic hydrocarbon in inter-tidal mussel *Perna perna*: Space-time observations, source investigation and genotoxicity. *Mar. Pollut. Bull.* **54**: 329-338.
 - 13) Barron, M.G., M.G. Carls, R. Heintz, and S.D. Rice (2004). Evaluation of fish early life-stage toxicity models of chronic embryonic exposures to complex polycyclic aromatic hydrocarbon mixtures. *Toxicol. Sci.*, **78**: 60-67.
 - 14) Reddy, C.M., T.I. Eglinton, A. Hounshell, H.K. White, L. Xu, R.B. Gaines, and G.S. Frysiner (2002). The West Falmouth oil spill after thirty years: the persistence of petroleum hydrocarbons in marsh sediments. *Environ. Sci. Tec.*, **36**. 4754-4760.
 - 15) Wenzl, T., R. Simon, E. Anklam, and J. Kleiner (2006). Analytical methods for polycyclic aromatic hydrocarbons (PAHs) in food and the environment needed for new food legislation in the European Union. *Anal. Chem.* **25**. 716-725.