

# Inhibitory effect of cyclophilin A from the hard tick *Haemaphysalis longicornis* on the growth of *Babesia bovis* and *Babesia bigemina*

著者	MAEDA Hiroki, BOLDBAATAR Damdinsuren, KUSAKISAKO Kodai, GALAY Remil Linggatong, AUNG Kyaw Min, UMEMIYA-SHIRAFUJI Rika, MOCHIZUKI Masami, FUJISAKI Kozo, TANAKA Tetsuya
journal or publication title	Parasitology research
volume	112
number	6
page range	2207-2213
year	2013
ファイル(説明)	Supplementary Figures table figures
URL	<a href="http://hdl.handle.net/10232/21419">http://hdl.handle.net/10232/21419</a>

doi: info:doi/10.1007/s00436-013-3390-7

1 Inhibitory effect of cyclophilin A from the hard tick *Haemaphysalis longicornis* on the  
2 growth of *Babesia bovis* and *Babesia bigemina*

3 Hiroki Maeda,<sup>1</sup> Damdinsuren Boldbaatar,<sup>1</sup> Kodai Kusakisako,<sup>1</sup> Remil Linggatong

4 Galay,<sup>1,2</sup> Kyaw Min Aung,<sup>1</sup> Rika Umemiya-Shirafuji,<sup>3</sup> Masami Mochizuki,<sup>1,2</sup> Kozo

5 Fujisaki,<sup>4</sup> and Tetsuya Tanaka<sup>1,2\*</sup>

6 <sup>1</sup>*Laboratory of Emerging Infectious Diseases, Joint Faculty of Veterinary Medicine,*

7 *Kagoshima University, Korimoto, Kagoshima 890-0065, Japan*

8 <sup>2</sup>*Department of Pathological and Preventive Veterinary Science, The United Graduate*

9 *School of Veterinary Science, Yamaguchi University, Yoshida, Yamaguchi 753-8515,*

10 *Japan*

11 <sup>3</sup>*National Research Center for Protozoan Diseases, Obihiro University of Agriculture*

12 *and Veterinary Medicine, Inada-cho, Obihiro, Hokkaido 080-8555, Japan*

13 <sup>4</sup>*National Agricultural and Food Research Organization, Tsukuba, Ibaraki 305-0856,*

14 *Japan*

15

16 **Running title:** Babesiacid activity of tick Cyclophilin A

17

18

19 \*To whom editorial correspondence should be addressed:

20 Dr. Tetsuya Tanaka

21 Laboratory of Emerging Infectious Diseases, Joint Faculty of Veterinary Medicine,

22 Kagoshima University, 1-21-24 Korimoto, Kagoshima 890-0065, Japan

23 Tel./Fax: +81-99-285-3539

24 E-mail: [tetsuya@ms.kagoshima-u.ac.jp](mailto:tetsuya@ms.kagoshima-u.ac.jp)

25

26

27

28

29

30

31

32

33

34

35

36

37 **Abstract**

38

39 *Haemaphysalis longicornis* is known as one of the most important ticks  
40 transmitting *Babesia* parasites in East Asian countries, including *Babesia ovata* and  
41 *Babesia gibsoni*, as well as *Theileria* parasites. *H. longicornis* is not the natural vector of  
42 *Babesia bovis* and *Babesia bigemina*. Vector ticks and transmitted parasites are thought  
43 to have established unique host-parasite interaction for their survival, meaning that  
44 vector ticks may have defensive molecules for the growth control of parasites in their  
45 bodies. However, the precise adaptation mechanism of tick-*Babesia* parasites is still  
46 unknown. Recently, cyclophilin A (CyPA) was reported to be important for the  
47 development of *Babesia* parasites in ticks. To reveal a part of their adaptation  
48 mechanism, the current study was conducted. An injection of *B. bovis*-infected RBCs  
49 into adult female *H. longicornis* ticks was found to upregulate the expression profiles of  
50 the gene and protein of CyPA in *H. longicornis* (HlCyPA). In addition, recombinant  
51 HlCyPA (rHlCyPA) purified from *Escherichia coli* exhibited significant inhibitory  
52 growth effects on *B. bovis* and *B. bigemina* cultivated *in vitro*, without any hemolytic  
53 effect on bovine RBCs at all concentrations used. In conclusion, our results suggest that  
54 HlCyPA might play an important role in the growth regulation of *Babesia* parasites in *H.*

55 *longicornis* ticks, during natural acquisition from an infected host. Furthermore,

56 rHlCyPA may be a potential alternative chemotherapeutic agent against babesiosis.

57

58 Key words: Cyclophilin, Tick, recombinant, *Babesia*, RBC, parasite

59

## 60 **Introduction**

61

62 Babesiosis is an important protozoan disease caused by *Babesia* parasites.

63 *Babesia* species are tick-transmitted protozoans that comprise some of the most

64 ubiquitous and widespread parasites of red blood cells (RBCs), affecting a wide range of

65 wild and economically important domestic animals and also humans (Homer et al. 2000;

66 Schnittger et al. 2012; Vannier et al. 2008). From the economic and public health

67 perspectives, sustained and continued research on babesiosis is needed for the

68 development of effective therapeutic medication.

69 *Haemaphysalis longicornis* is one of a well-known vector tick of *Babesia* spp.

70 *Babesia ovata* and *Babesia gibsoni*, as well as *Theileria* parasites are transmitted by *H.*

71 *longicornis*. On the other hand, *Babesia bovis* and *Babesia bigemina* are transmitted by

72 *Rhipicephalus (Boophilus) microplus* (Schnittger et al. 2012). The precise adaptation

73 mechanism of tick-*Babesia* parasites is still unknown.

74           Immunophilin is the generic name of isomerases such as the cyclophilins  
75 (CyPs) and FK-binding protein (FKBPs) families. Immunophilin binds specific  
76 immunosuppressive drugs; e.g., CyPs and FKBP bind the cyclic peptide cyclosporine A  
77 and the macrolactones FK506 (tacrolimus) and rapamycin (sirolimus), respectively.  
78 CyPs have been found in many eukaryotes. They possess peptidyl-prolyl *cis-trans*  
79 isomerase (PPIase) activity. PPIase can catalyze the *cis-trans* isomerization of the  
80 peptide bonds preceding proline residues, which involves in a wide range of cellular  
81 processes, such as cell division, transcriptional regulation, protein trafficking, and RNA  
82 splicing. In addition to enzymatic activities, many immunophilins act as molecular  
83 chaperones. Therefore, most members of the CyP family have been shown to function as  
84 mediators of intra- and inter-cellular communication (Barik et al. 2006; Bell et al. 2006;  
85 Galat 1993, 2004; Krücken et al. 2009).

86           Cyclophilins and their related molecules were identified and characterized  
87 previously in many living organisms; however, in ticks, only a few cyclophilin gene  
88 sequences were identified, and their functions remain unknown. Recently, we reported  
89 that cyclophilin A (CyPA) from the ixodid tick *H. longicornis*, *H. longicornis* CyPA  
90 (HlCyPA), has a conserved PPIase domain and is expressed in multiple organs as well

91 as throughout all developmental stages (Boldbaatar et al. 2008). Recombinant HICyPA  
92 (rHICyPA) was found to exhibit PPIase activity. After knockdown of the *HICyPA* gene  
93 by RNA interference (RNAi), engorged female ticks had significantly lower body  
94 weight and failed to lay eggs. Furthermore, some RNAi-treated ticks died after  
95 engorgement. In addition, there was one report on the putative immunophilin gene in *R.*  
96 (*B.*) *microplus* ticks that showed high homology with the *HICyPA* gene, wherein gene  
97 silencing significantly increased the infection rate of *Babesia bovis* in the larval progeny  
98 (Bastos et al. 2009). These reports strongly suggest that tick immunophilin genes and  
99 their products play important roles in tick physiology and as defensive immunological  
100 mechanisms against parasites. The current study was conducted to evaluate the response  
101 of HICyPA to *Babesia* infection and its inhibitory growth effects on *Babesia* parasites  
102 cultivated *in vitro*. Two bovine *Babesia* parasites, *B. bovis*, *B. bigemina* and non-vector  
103 tick, *H. longicornis* were also used in this study to clarify vector-parasite adaptation  
104 mechanism. This is the first report on the inhibitory effect of cyclophilin from ticks on a  
105 tick-borne pathogen.

106

107

108 **Materials and methods**

109

110 Ticks and Animals

111

112           The parthenogenetic Okayama strain of *H. longicornis* has been maintained by  
113 blood feeding on Japanese white rabbits (Kyudo, Kumamoto, Japan) (Fujisaki 1978) in  
114 the Laboratory of Emerging Infectious Diseases, Joint Faculty of Veterinary Medicine,  
115 Kagoshima University.

116           Rabbits were kept in accordance with the guidelines approved by the Animal  
117 Care and Use Committee of Kagoshima University (Approval number A08010). They  
118 were maintained under regulated conditions throughout the experiments.

119

120 Culture of *Babesia* parasites

121

122           Both *Babesia bovis* (the Texan strain) and *Babesia bigemina* (the Argentine  
123 strain) were used in this study (Bork et al. 2004). They were maintained on purified  
124 bovine RBCs using different culture media for each species (Galay et al. 2012).

125

126 Injection of *B. bovis*-infected RBCs to ticks



127

128           Unfed adult ticks were injected with 0.5  $\mu$ l of *B. bovis*-infected RBCs or  
129 uninfected normal RBCs (Control) through the fourth coxae into the hemocoel, as  
130 previously described (Aung et al. 2012). The degree of parasitemia of the *B.*  
131 *bovis*-infected RBCs was 5%. After injection, nine ticks were collected every 24 h; one  
132 was used for genomic DNA extraction for the detection of *B. bovis*, and three were used  
133 for total RNA extraction and complementary DNA (cDNA) synthesis. The remaining  
134 five were used for protein extraction. The level of expression of the *HlCyPA* gene was  
135 investigated by real-time PCR, and protein expression of HlCyPA was determined by  
136 Western blot analysis. At the first step of real-time PCR, *actin*, *tubulin*, *P0*, and *L23*  
137 genes were selected for tick reference and evaluated for standardization.

138

139 RNA extraction and cDNA synthesis

140

141           To extract total RNA, ticks were homogenized using Automill (Tokken, Tiba,  
142 Japan), to which the TRI<sup>®</sup> reagent (Sigma, MO, USA) was added. The extracted RNA  
143 was purified with the Turbo DNA-free<sup>™</sup> Kit (Applied Biosystems, Tokyo, Japan).  
144 cDNA synthesis was performed with ReverTra Ace- $\alpha$ -<sup>®</sup> (TOYOBO, Osaka, Japan)

145 following the manufacturer's protocol using 1 µg of total RNA.

146

147 DNA extraction

148

149 Genomic DNA was also extracted from collected ticks. Homogenized ticks  
150 were suspended in an extraction buffer [100 mM Tris-HCl (pH 8.0), 0.5% SDS, 100  
151 mM NaCl, 10 mM EDTA], and, after adding proteinase K (10 mg/ml) (KANTO  
152 CHEMICAL, Tokyo, Japan), samples were incubated overnight at 55°C. After removal  
153 of proteins using Phenol:Chloroform:IsoamylAlcohol (Sigma), ethanol precipitation was  
154 performed to collect DNA. DNA samples were purified with an RNaseA solution (4  
155 mg/ml) (Promega, WI, USA).

156

157 Expression analysis of the *HICyPA* gene and detection of the *B. bovis* gene

158

159 The expression analysis of the *HICyPA* gene was performed by real-time PCR  
160 using THUNDERBIRD™ SYBR® qPCR Mix (TOYOBO) with a 7300 real-time PCR  
161 system (Applied Biosystems). Gene-specific primers were designed to target the  
162 *HICyPA* gene (Boldbaatar et al. 2008) and the control genes, as shown in Table 1.

163 Standard curves were made from eight-fold serial dilutions of cDNA of adult ticks fed  
164 for 3 days. The PCR cycle profile was as follows: 95 °C for 10 min, 40 cycles of a  
165 denaturation step at 95 °C for 15 sec, and an annealing/extension step at 60 °C for 60 sec.  
166 The data was analyzed with 7300 system SDS software (Applied Biosystems).

167           Detection of the *B. bovis SSrRNA* gene was performed using PCR as described  
168 by Adham et al. (2009) with a slight modification of the thermo cycle profile at 94 °C for  
169 5 min, 40 cycles of a denaturation step at 94 °C for 1 min, an annealing/extension step at  
170 72 °C for 2 min, and final extension at 72 °C for 7 min.

171

172 Protein extraction and Western blot analysis

173

174           Homogenized ticks were suspended in phosphate-buffered saline (PBS),  
175 ultrasonicated three times (2 min each; Vibra Cell™; Sonics and Materials, CT, USA)  
176 on ice, and finally centrifuged at 500 × g. The supernatant was resolved in 15%  
177 SDS-PAGE (Laemmli 1970) under reducing conditions. After SDS-PAGE, the proteins  
178 were transferred onto a polyvinylidene difluoride membrane (Immobilon®-P; Millipore,  
179 MA, USA). The membrane was blocked overnight with 5% skim milk in PBS and then  
180 incubated with a 1:500 dilution of anti-rHICyPA mouse sera (Boldbaatar et al. 2008) at

181 37°C for 1 h. Tubulin was used as the control protein (Umemiya-Shirafuji et al. 2012).  
182 After washing five times in PBS containing 0.05% Tween20, the membrane was  
183 incubated with a 1:50,000 dilution of horseradish peroxidase (HRP)-conjugated sheep  
184 anti-mouse IgG (GE Healthcare, Buckinghamshire, UK) at 37°C for 1 h. After washing  
185 five times in PBS containing 0.05% Tween20, bands were detected using the  
186 Amersham™ ECL™ Prime Western Blotting Detection Reagent (GE Healthcare) and  
187 viewed using FluorChem®FC2 software (Alpha Innotech, CA, USA).

188

189 Expression and purification of recombinant rHICyPA

190

191 Recombinant plasmids (Boldbaatar et al. 2008) were used to transform into  
192 *Escherichia coli* (BL21), and histidine-tagged rHICyPA expression was induced by 1  
193 mM Isopropyl-β-D(-)-thiogalactopyranoside (IPTG) (Wako, Osaka, Japan) at 37°C for 6  
194 h. The expressed recombinant protein was purified using a His trap FF column (GE  
195 Healthcare) containing 1 ml of chelating sepharose with nickel ions using the Bio Logic  
196 Duo Flow Base System (BIO-RAD, Tokyo, Japan). The purified recombinant protein  
197 was dialyzed against PBS. The concentration of rHICyPA was determined using the  
198 Micro BCA™ protein assay kit (Thermo Fisher Scientific, MA, USA) and rHICyPA

199 was stored at -30°C until use. The PPIase activity was also confirmed as described by  
200 Boldbaatar et al. (2008).

201

202 Hemolysis assay

203

204 The hemolytic activity of rHlCyPA was determined according to the method  
205 described by Stark et al. (2002). Briefly, bovine RBCs were washed with PBS. Then,  
206 from 0.01 to 3.3  $\mu$ M concentrations, rHlCyPA was mixed with bovine RBCs in a  
207 96-well plate (Nunc, Roskilde, Denmark). The plate was incubated at 37 °C for 1 h and  
208 centrifuged at 1000  $\times$  g for 5 min. The supernatant was collected, and the degree of  
209 hemolysis was assessed by measuring the absorbance at 550 nm in a microplate reader  
210 Model 680 (BIO-RAD). PBS and Triton-X were used as agents for preparing the 0 and  
211 100% hemolyses.

212

213 Effect of recombinant HlCyPA on *Babesia* parasites *in vitro*

214

215 The culture media of *Babesia* parasites were changed daily, and rHlCyPA was  
216 added each day at different concentrations of 3.3, 33, 330 nM, and 3.3  $\mu$ M. An equal

217 volume of PBS was used for the control group. Blood smears with Giemsa staining were  
218 made daily to calculate the parasitemia and observe morphology of *Babesia* parasites.  
219 Three replicated wells were tested on the each group. Parasitemia was calculated as the  
220 percentage of infected RBCs to 1,000 RBCs counted.

221

222 Statistical analysis

223

224 All experiments were conducted with two or three separate trials. Data were  
225 statistically analyzed using the Student's *t*-test; results are presented as the mean  $\pm$  SE,  
226 and  $P < 0.05$  was considered statistically significant.

227

228 **Results**

229

230 Expression profiles of the *HlCyPA* gene and protein in *H. longicornis* females injected  
231 with *B. bovis*-infected RBCs

232

233 In the group of adult female *H. longicornis* ticks injected with *B. bovis*-infected  
234 RBCs, the *HlCyPA* gene expression increased faster than the control group (*H.*

235 *longicornis* female ticks injected with normal bovine RBCs), and a significant difference  
236 ( $*P < 0.05$ , Control group vs. *B. bovis*-infected RBC-injected group) was observed 1-3  
237 days after the injection (Fig. 1a). In addition, the expression levels of HICyPA protein  
238 showed a similar pattern compared to gene expression (Fig. 1b). The protein expression  
239 levels were quantified by using the densitometry analysis. During the first two days after  
240 the injection, protein expression of *B. bovis*-infected RBC-injected group tended to  
241 increase faster than in the control group. In addition, *B. bovis* DNA was detected by  
242 PCR to confirm the success of the injection (Supplementary Fig. 1).

243

244 Hemolytic activity of recombinant HICyPA against bovine RBCs

245

246 No hemolysis was observed in bovine RBCs incubated with any concentration  
247 of rHICyPA from 0.01 to 3.3  $\mu\text{M}$ . The hemolytic activity of rHICyPA was compared  
248 with Triton-X and expressed as % hemolysis. The percentage of hemolysis was lower  
249 than 5% and almost negligible. at all concentrations of rHICyPA determined  
250 (Supplementary Fig. 2).

251 Effect of recombinant HICyPA on the growth of *B. bovis* and *B. bigemina in vitro*

252

253           There were no significant differences on the growth of *B. bovis* (Fig. 2a) and *B.*  
254 *bigemina* (Fig. 2b) in the presence of rHlCyPA, at concentrations from 330 nM and  
255 lower. However, the growth of both species was completely inhibited in the culture with  
256 3.3  $\mu$ M rHlCyPA at 3 and 4 days (Fig. 2). In addition, in the presence of 3.3  $\mu$ M  
257 rHlCyPA, *Babesia* parasites were sparsely observed under light microscopy, and most of  
258 them had an abnormal ring-form-like morphology (Fig. 3).

259

## 260 **Discussion**

261

262           *Babesia* is one of best-known parasites transmitted by ticks and has been  
263 considered to be seriously injurious to tick biology (Florin-Christensen and Schnittger  
264 2009). Therefore, it was speculated that ticks may have developed defensive molecular  
265 mechanisms to reduce and/or attenuate the harmful and injurious effects of *Babesia*  
266 parasites. On the other hand, *Babesia* parasites are believed to be capable of avoiding  
267 the defensive mechanisms of ticks (Florin-Christensen and Schnittger 2009; Sonenshine  
268 and Hynes 2008). The existing and sustainable host-parasite relationship between ticks  
269 and *Babesia* parasites is assumed to be maintained on the basis of superb molecular  
270 mechanisms for conflict of interest or potential conflict of interest (Chauvin et al. 2009;



271 Florin-Christensen and Schnittger 2009).

272           The cattle tick, *R. (B.) microplus*, is a known natural vector of *B. bovis* (Bock et  
273 al. 2004; Schnittger et al. 2012), and their immunophilin gene hinders *B. bovis* infection,  
274 which suggests that the gene plays an important role in the control of the transmission of  
275 protozoa (Bastos et al. 2009). *H. longicornis* is also an important tick vector of *Babesia*  
276 spp. (Schnittger et al. 2012); however, it is not a natural vector for *B. bovis* and *B.*  
277 *bigemina* (Bock et al. 2004). These reports suggest that *H. longicornis* might not have  
278 established a control strategy for these *Babesia* species or may have developed some  
279 defense mechanisms for them. Interestingly, in *H. longicornis*, an immunophilin gene  
280 has been identified and characterized, *HICyPA* possessing 90% identity with the  
281 immunophilin gene of *R. (B.) microplus*. Silencing of *HICyPA* through RNAi has led to  
282 a significant reduction in the body weight of engorged ticks and their failure to lay eggs  
283 (Boldbaatar et al. 2008). This result indicates that HICyPA represents a major  
284 cyclophilin protein in *H. longicornis* involved in blood ingestion, tick viability, and  
285 oocyte development. Therefore, HICyPA might also be an important protein involved in  
286 a tick's innate immunity.

287           In this context, this study was conducted to investigate the possible role of  
288 HICyPA against two *Babesia* parasites. To understand the interaction of HICyPA and

289 *Babesia* parasites, *B. bovis*-infected RBCs were injected into *H. longicornis*. As shown  
290 in Fig. 1, the injection of *B. bovis*-infected RBCs may have caused the upregulation of  
291 *HICyPA* gene and its product. These results suggest that HICyPA might be related in the  
292 tick immune response against *Babesia* parasites. In a previous study, we showed that the  
293 *HICyPA* gene was expressed in many organs, and the expression level was the highest in  
294 the midgut and salivary glands (Boldbaatar et al. 2008). Both of them are important  
295 organs involved in the multiplication and transmission of *Babesia* parasites in vector  
296 ticks (Chauvin et al. 2009; Florin-Christensen and Schnittger 2009). These results  
297 suggest that HICyPA may be related to the tick's immune response to *Babesia* parasites.  
298 In *H. longicornis*, a cysteine protease, longipain, is known to be highly expressed in the  
299 midgut as well as HICyPA and act as a defense molecule against invading *Babesia*  
300 parasites (Tsuji et al. 2008). The defensin-like peptide, longicin, was also found to  
301 possess activities against different pathogens, e.g., antimicrobial activity, fungicidal  
302 activity, and parasiticidal activity, including babesiacidal activity (Tsuji et al. 2007).  
303 Additional studies demonstrated that the synthetic partial peptide, P4 of longicin,  
304 showed similar activities, including parasiticidal action, against *Toxoplasma gondii*  
305 (Rahman et al. 2010; Tanaka et al. 2012). These results suggest that HICyPA may act  
306 synergistically with longipain and/or longicin to eliminate parasites, bacteria, and

307 viruses.

308           To further evaluate the effect of HlCyPA on *Babesia* parasites, rHlCyPA was  
309 prepared (Boldbaatar et al. 2008). In this study, a dose-dependent inhibitory effect of  
310 rHlCyPA on the growth of *B. bovis* and *B. bigemina* was observed (Fig. 2), and  
311 rHlCyPA affected their morphology (Fig. 3). Even though lower concentrations of  
312 rHlCyPA seemed to inhibit *B. bigemina* more efficiently than *B. bovis*, this may be due  
313 to the more rapid increase of *B. bovis* than of *B. bigemina*. Coagulation disorders,  
314 cytoadherence, and the hypotensive state seen in acute *B. bovis* infections are not  
315 features of *B. bigemina* infections (Bock et al. 2004). These differences in pathogenicity  
316 probably reflect a distinction in the metabolic or infection mechanisms between *B. bovis*  
317 and *B. bigemina*. Therefore, these differences may affect the susceptibility of *B. bovis*  
318 and *B. bigemina* to rHlCyPA. On the other hand, no toxic effects of rHlCyPA against  
319 host RBCs were observed (Supplementary Fig. 2). This result revealed that the  
320 babesiacidal effect of rHlCyPA is not due to the hemolysis of host RBCs. These results  
321 support the idea that HlCyPA plays an important role in controlling the multiplication of  
322 *Babesia* parasites as a defensive molecule in vector ticks.

323           Many agents possessing babesiacidal activity have been reported. In this study,  
324 rHlCyPA showed an inhibitory growth effect on *Babesia* parasites *in vitro* (Fig. 2) at a

325 lower or similar concentration to triclosan (Bork et al. 2003), heparin (Bork et al. 2004),  
326 nerolidol (AbouLaila et al. 2010), artesunate (Goo et al. 2010; Nagai et al. 2003), or  
327 fusidic acid (Salama et al. 2012). This result suggests that rHCyPA could be a potential  
328 anti-babesial agent as powerful as these drugs.

329           Meanwhile, recent reports suggested that cyclophilin may also play an  
330 important role in the host-response to viruses (Luban et al. 2007; Nagy et al. 2011; Zhou  
331 et al. 2012). Particularly, CyPA in chicken and human cells restricts influenza A virus  
332 replication through interaction and degradation with the viral M1 protein. Moreover, this  
333 inhibitory effect of CyPA on the influenza virus infection process is not dependent on its  
334 isomerase activity (Liu et al. 2009; Liu et al. 2012; Xu et al. 2010). These results suggest  
335 that the CyPA is also involved in some viral diseases and several effects of CyPA do not  
336 require PPIase activity. There are many tick-borne viruses ("tioviruses") have been  
337 detected (Hub á lek and Rudolf 2012). Tick CyPA might be involved tick-virus  
338 interaction.

339           In this study, the biological action of HCyPA was not fully elucidated;  
340 however, it is predicted that HCyPA may also have other functions in ticks and in the  
341 transmission of tick-borne diseases. Thus, further studies will be required to determine  
342 its potential as an alternative chemotherapeutic agent. Moreover, the effect of HCyPA

343 *in vivo* should be evaluated. Further understanding on the underlying mechanism of the  
344 babesiacidal effect of HICyPA, as well as its other functions in the hard tick, may  
345 contribute to control of both babesiosis and its vector.

346

### 347 **Acknowledgments**

348

349 We are grateful to Dr. C. Kubota, Laboratory of Veterinary Theriogenology,  
350 Department of Clinical Veterinary Science, Joint Faculty of Veterinary Medicine,  
351 Kagoshima University, and Dr. H. Yamaguchi, Iriki Livestock Farm, Joint Faculty of  
352 Veterinary Medicine, Kagoshima University, for the supply of bovine blood. This work  
353 was supported by the Bio-oriented Technology Research Advancement Institution  
354 (BRAIN) and Grants-in-Aid for Scientific Research (A) and (C) from the Japan Society  
355 for the Promotion of Science (JSPS).

356

357

358

### 359 **References**

360

361 AbouLaila M, Sivakumar T, Yokoyama N, Igarashi I (2010) Inhibitory effect of terpene  
362 nerolidol on the growth of *Babesia* parasites. *Parasitol Int* 59:278-282

363 Adham FK, Abd-el-Samie EM, Gabre RM, el-Hussein H (2009) Detection of tick blood  
364 parasites in Egypt using PCR assay I-*Babesia bovis* and *Babesia bigemina*. *Parasitol*  
365 *Res* 105:721-730

366 Aung KM, Boldbaatar D, Umemiya-Shirafuji R, Liao M, Tsuji N, Xuan X, Suzuki H,  
367 Kume A, Galay RL, Tanaka T, Fujisaki K (2012) HISRB, a Class B scavenger  
368 receptor, is key to the granulocyte-mediated microbial phagocytosis in ticks. *PLoS*  
369 *ONE* 7:e33504

370 Barik S (2006) Immunophilins: for the love of proteins. *Cell Mol Life Sci* 63:2889-2900

371 Bastos RG, Ueti MW, Guerrero FD, Knowles DP, Scoles GA (2009) Silencing of a  
372 putative immunophilin gene in the cattle tick *Rhipicephalus (Boophilus) microplus*  
373 increases the infection rate of *Babesia bovis* in larval progeny. *Parasit Vectors* 2:57

374 Bell A, Monaghan P, Page AP (2006) Peptidyl-prolyl *cis-trans* isomerases  
375 (immunophilins) and their roles in parasite biochemistry, host-parasite interaction  
376 and antiparasitic drug action. *Int J Parasitol* 36:261-276

377 Bock R, Jackson L, De Vos A, Jorgensen W (2004) Babesiosis of cattle. *Parasitology*  
378 129:S247-S269

379 Boldbaatar D, Kilonzo RM, Battur B, Umemiya R, Liao M, Tanaka T, Xuan X, Fujisaki  
380 K (2008) Identification of two forms of cyclophilin from the hard tick  
381 *Haemaphysalis longicornis*. Process Biochemistry 43:615-625

382 Bork S, Yokoyama N, Matsuo T, Claveria FG, Fujisaki K, Igarashi I (2003) Growth  
383 inhibitory effect of triclosan on equine and bovine *Babesia* parasites. Am J Trop  
384 Med Hyg 68:334-340

385 Bork S, Yokoyama N, Ikehara Y, Kumar S, Sugimoto C, Igarashi I (2004)  
386 Growth-inhibitory effect of heparin on *Babesia* parasites. Antimicrob Agents  
387 Chemother 48:236-241

388 Chauvin A, Moreau E, Bonnet S, Plantard O, Malandrin L (2009) *Babesia* and its hosts:  
389 adaptation to long-lasting interactions as a way to achieve efficient transmission.  
390 Vet Res 40:37

391 Florin-Christensen M, Schnittger L (2009) Piroplasmids and ticks: a long-lasting  
392 intimate relationship. Front Biosci 14:3064-3073

393 Fujisaki K (1978) Development of acquired resistance precipitating antibody in rabbits  
394 experimentally infested with females of *Haemaphysalis longicornis* (Ixodoidea:  
395 Ixodidae). Natl Inst Anim Health Q (Tokyo) 18:27-38

396 Galat A (1993) Peptidylproline *cis-trans* isomerases: immunophilins. Eur J Biochem

397 216:689-707

398 Galat A (2004) Function-dependent clustering of orthologues and paralogues of  
399 cyclophilins. *Proteins* 56:808-820

400 Galay RL, Maeda H, Aung KM, Umemiya-Shirafuji R, Xuan X, Igarashi I, Tsuji N,  
401 Tanaka T, Fujisaki K (2012) Anti-babesial activity of a potent peptide fragment  
402 derived from longicin of *Haemaphysalis longicornis*. *Trop Anim Health Prod*  
403 44:343-348

404 Goo YK, Terkawi MA, Jia H, Aboge GO, Ooka H, Nelson B, Kim S, Sunaga F,  
405 Namikawa K, Igarashi I, Nishikawa Y, Xuan X (2010) Artesunate, a potential drug  
406 for treatment of *Babesia* infection. *Parasitol Int* 59:481-486

407 Homer MJ, Aguilar-Delfin I, Telford SR 3rd, Krause PJ, Persing DH (2000) Babesiosis.  
408 *Clin Microbiol Rev* 13:451-469

409 Hubálek Z, Rudolf I (2012) Tick-borne viruses in Europe. *Parasitol Res* 1:9-36.

410 Krücken J, Greif G, von Samson-Himmelstjerna G (2009) In silico analysis of the  
411 cyclophilin repertoire of apicomplexan parasites. *Parasit Vectors* 2:27

412 Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of  
413 bacteriophage T4. *Nature* 227:680-685

414 Liu X, Sun L, Yu M, Wang Z, Xu C, Xue Q, Zhang K, Ye X, Kitamura Y, Liu W



415 (2009) Cyclophilin A interacts with influenza A virus M1 protein and impairs the  
416 early stage of the viral replication. *Cell Microbiol* 11:730-741

417 Liu X, Zhao Z, Xu C, Sun L, Chen J, Zhang L, Liu W (2012) Cyclophilin A restricts  
418 influenza A virus replication through degradation of the M1 protein. *PLoS ONE*  
419 7:e31063

420 Luban J (2007) Cyclophilin A, TRIM5, and resistance to human immunodeficiency  
421 virus type 1 infection. *J Virol* 81:1054-1061

422 Nagai A, Yokoyama N, Matsuo T, Bork S, Hirata H, Xuan X, Zhu Y, Claveria FG,  
423 Fujisaki K, Igarashi I (2003) Growth-inhibitory effects of artesunate,  
424 pyrimethamine, and pamaquine against *Babesia equi* and *Babesia caballi* in *in vitro*  
425 cultures. *Antimicrob Agents Chemother* 47:800-803

426 Nagy PD, Wang RY, Pogany J, Hafren A, Makinen K (2011) Emerging picture of host  
427 chaperone and cyclophilin roles in RNA virus replication. *Virology* 411:374-382

428 Rahman M, Tsuji N, Boldbaatar D, Battur B, Liao M, Umemiya-Shirafuji R, You M,  
429 Tanaka T, Fujisaki K (2010) Structural characterization and cytolytic activity of a  
430 potent antimicrobial motif in longicin, a defensin-like peptide in the tick  
431 *Haemaphysalis longicornis*. *J Vet Med Sci* 72:149-156

432 Salama AA, Aboulaila M, Moussa AA, Nayel MA, El-Sify A, Terkawi MA, Hassan HY,

433 Yokoyama N, Igarashi I (2012) Evaluation of *in vitro* and *in vivo* inhibitory effects  
434 of fusidic acid on *Babesia* and *Theileria* parasites. Vet Parasitol, In press

435 Schnittger L, Rodriguez AE, Florin-Christensen M, Morrison DA (2012) *Babesia*: A  
436 world emerging. Infect Genet Evol 12:1788-1809

437 Sonenshine DE, Hynes WL (2008) Molecular characterization and related aspects of the  
438 innate immune response in ticks. Front Biosci 13:7046-63.

439 Stark M, Liu LP, Deber CM (2002) Cationic hydrophobic peptides with antimicrobial  
440 activity. Antimicrob Agents Chemother 46:3585-3590

441 Tanaka T, Maeda H, Matsuo T, Boldbaatar D, Umemiya-Shirafuji R, Kume A, Suzuki H,  
442 Xuan X, Tsuji N, Fujisaki (2012) Parasitocidal activity of *Haemaphysalis*  
443 *longicornis* longicin P4 peptide against *Toxoplasma gondii*. Peptides 34:242-250

444 Tsuji N, Battsetseg B, Boldbaatar D, Miyoshi T, Xuan X, Oliver JH Jr, Fujisaki K  
445 (2007) Babesial vector tick defensin against *Babesia* sp. parasites. Infect Immun  
446 75:3633-3640

447 Tsuji N, Miyoshi T, Battsetseg B, Matsuo T, Xuan X, Fujisaki K (2008) A cysteine  
448 protease is critical for *Babesia* spp. transmission in *Haemaphysalis longicornis*  
449 ticks. PLoS Pathog 4:e1000062

450 Umemiya-Shirafuji R, Tanaka T, Boldbaatar D, Tanaka T, Fujisaki K (2012) Akt is an

451 essential player in regulating cell/organ growth at the adult stage in the hard tick  
452 *Haemaphysalis longicornis*. Insect Biochem Mol Biol 42:164-173

453 Vannier E, Gewurz BE, Krause PJ (2008) Human Babesiosis. Infect Dis Clin North Am  
454 22:469-488

455 Xu C, Meng S, Liu X, Sun L, Liu W (2010) Chicken cyclophilin A is an inhibitory  
456 factor to influenza virus replication. Virol J, 7:372

457 Zhou D, Mei Q, Li J, He H (2012) Cyclophilin A and viral infections. Biochem Biophys  
458 Res Commun 424:647-650

459

#### 460 **Figure legends**

461

462 **Fig. 1** (a) Gene expression of *HlCyPA* 1-7 days after injection of *B. bovis*-infected RBCs.  
463 *B. bovis*, *B. bovis*-infected RBC-injected group; RBC, RBC-injected group for control.  
464 \* $P < 0.05$ , significantly different, RBC vs. *B. bovis*. (b) Protein expression of HlCyPA  
465 1-7 days after injection of *B. bovis*-infected RBCs. *B. bovis*, *B. bovis*-infected  
466 RBC-injected group; RBC, RBC-injected group for control; Numbers indicate days after  
467 injection. The line graph shows the relative expression of CyPA to tubulin determined  
468 using densitometry.

469

470 **Fig. 2** Effect of recombinant HlCyPA on the growth of *B. bovis* and *B. bigemina*. In  
471 *vitro* culture of *B. bovis* (a) and *B. bigemina* (b) with different concentrations of  
472 rHlCyPA. Parasitemia was monitored for 4 days. \* $P < 0.05$ , significantly different,  
473 Control vs. rHlCyPA-treated group.

474

475 **Fig. 3** Light micrograph of Giemsa-stained blood smear showing parasite morphology  
476 from Control and 3.3  $\mu\text{M}$  rHlCyPA-treated groups. a, *B. bovis*; b, *B. bigemina*. Arrows  
477 indicate ring-form-like parasites. Bar: 5  $\mu\text{m}$

478

479 **Supplementary Fig. 1** Detection of *B. bovis* DNA from *H. longicornis* injected with *B.*  
480 *bovis*-infected RBCs. *B. bovis*, *B. bovis*-infected RBC-injected group; RBC,  
481 RBC-injected group for control. Numbers indicate days after injection.

482

483 **Supplementary Fig. 2** Hemolysis assay of recombinant HlCyPA. Each percentage  
484 represents the ratio vs. Triton-X as 100% hemolysis. PBS was also used for 0%  
485 hemolysis.