

Ghrelin and glucagon-like peptide-2 increase immediately following massive small bowel resection

著者	MUTO Mitsuru, KAJI Tatsuru, MUKAI Motoi, NAKAME Kazuhiko, YOSHIOKA Takako, TANIMOTO Akihide, MATSUFUJI Hiroshi
journal or publication title	Peptides
volume	43
page range	160-166
ファイル(説明)	Table 3 Table 2 Table 1 Figures Main text Title page
URL	http://hdl.handle.net/10232/18958

doi: info:doi/10.1016/j.peptides.2013.03.006

title

Ghrelin and glucagon-like peptide-2 increase immediately following massive small bowel resection

list of authors and affiliations

Mitsuru Muto ^a, Tatsuru Kaji ^{a*}, Motoi Mukai ^a, Kazuhiko Nakame ^a, Takako Yoshioka ^b, Akihide Tanimoto ^b, Hiroshi Matsufuji ^a

^a Department of Pediatric Surgery, Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima, 8908520 Japan

^b Department of Molecular and Cellular Pathology, Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima, 8908520 Japan

* Corresponding author at: Department of Pediatric Surgery, Kagoshima University Graduate School of Medical and Dental Sciences, 8-35-1 Sakuragaoka, Kagoshima, 8908520 Japan.

Tel: +81 99 275 5444; fax: +81 99 275 2628.

E-mail address: tatu@m2.kufm.kagoshima-u.ac.jp

Abbreviations: GLP-2, glucagon-like peptide-2; 80% SBR, 80% small bowel resection.

abstract and keywords

Children with short bowel syndrome face life-threatening complications. Therefore, there is an urgent need for a new therapy to induce effective adaptation of the remnant intestine. Adaptation occurs only during feeding. We focused on preprandial acyl ghrelin and des-acyl ghrelin, and postprandial glucagon-like peptide-2 (GLP-2), which are known to have active orexigenic and trophic actions. This study aims to clarify the secretion trends of these hormones after massive small bowel resection and to obtain basic data for developing a new treatment. Sixty-three growing male rats were used: 3 were designated as controls receiving no operation and 60 were randomized into the 80% small bowel resection (80% SBR) group and the transection and re-anastomosis group. Changes in body weight, food intake, and remnant intestine morphology were also assessed for 15 days after the operation. Acyl ghrelin and des-acyl ghrelin levels increased immediately, equivalently in both operation groups ($P = 0.09$ and 0.70). Interestingly, in 80% SBR animals, des-acyl ghrelin peaked on day 1 and acyl ghrelin peaked on day 4 ($P = 0.0007$ and $P = 0.049$ vs controls). GLP-2 secretion was obvious in 80% SBR animals ($P = 2.25 \times 10^{-6}$), which increased immediately and peaked on day 4 ($P = 0.009$ vs controls). Body weight and food intake in 80% SBR animals recovered to preoperative levels on day 4. Morphological adaptations were evident after day 4. Our results may suggest a management strategy to reinforce these physiological hormone secretion patterns in developing a new therapy for short bowel syndrome.

Key Words:

Acyl ghrelin, Des-acyl ghrelin, GLP-2, adaptation, massive small bowel resection, short bowel syndrome