

# The fine slips of cutaneous muscle extending from the ventrolateral trunk to the brachium in the rat

著者	MINE Kazuharu, SHIMOTAKAHARA Rie, LEE Hyeyong, OGATA Shigemitsu, TAMATSU Yuichi
journal or publication title	Bulletin of the School of Health Sciences, Faculty of Medicine, Kagoshima University
volume	29
number	1
page range	97-101
year	2019-03-31
URL	<a href="http://hdl.handle.net/10232/00030654">http://hdl.handle.net/10232/00030654</a>

## The fine slips of cutaneous muscle extending from the ventrolateral trunk to the brachium in the rat

Kazuharu Mine<sup>1)</sup>, Rie Shimotakahara<sup>1)</sup>, Hyeyong Lee<sup>2)</sup>, Shigemitsu Ogata<sup>2)</sup>, Yuichi Tamatsu<sup>1)</sup>

1) Department of Gross Anatomy and Forensic Dentistry, Graduate School of Medical and Dental Sciences, Kagoshima University, Sakuragaoka 8-35-1, Kagoshima, 890-8544 Japan  
2) Department of Fundamental and Clinical Nursing, School of Health Science, Faculty of Medicine, Kagoshima University, Sakuragaoka 8-35-1, Kagoshima 890-8544, Japan

### Abstract

A fine slip separated from the external surface of the cutaneous trunci muscle was often found around the axillary region of the rat. However, there has been scarce report concerning such muscular bundle. The aim of this study was to examine it anatomically. The slip extended from the ventrolateral trunk to the fascia of brachium. It covered the thoracic mammary glands in female and the adipose mass in male. The nerve supply from the caudal pectoral nerve was expected, although the true innervation was not clearly identified. Moreover, it was suggested that the murine commonly possesses this type of muscle. The anatomical knowledge about this muscular slip will probably diminish the inconveniences during the skin experiment using rats.

**Key words:** skin, cutaneous trunci muscle, rat, anatomy, innervation

### Introduction

Rat is one of the popular animals used for the skin experiment including the creation of artificial skin wound<sup>1-2)</sup> and validation of skin flap design<sup>3-5)</sup>. Similar to other mammals, the rat has the cutaneous trunci muscle (CTM) that the human never possesses<sup>6-7)</sup>. To appreciate the morphological difference between the laboratory animals and the human in performing the skin experiment, we have studied the subcutaneous structures macro-anatomically<sup>8-9)</sup>.

When we exposed the external surface of the CTM at the axillary region of the rat in the previous studies, a certain muscular slip was often found out there. We searched description concerning such muscular bundle in anatomical textbooks of the rodents<sup>6-7,10)</sup> and literature of comparative myology<sup>11-12)</sup>. However, there was no descriptive text explaining about it.

The axillary region is frequently incised during the skin

experiments intended to the flap design and muscle reflex<sup>13-15)</sup>. At that time, the slip seems likely to be injured unconsciously. Therefore, we planned to examine the cutaneous muscle further focusing on the region from the flank to the forelimb. The aim of this study was to record the muscular slip separated from the CTM of the rat in detail and to confirm its anatomical property.

### Materials and Methods

The animals used in this study were two male and two female Wistar rats of 8 weeks of age. They were sacrificed for the control experiment of cerebral arteries by Dr. S. Okuyama<sup>16)</sup> in 2004. After the experiment, the dead bodies were handed us for exhaustive utilization of animal resources. Immersed in 10% formalin solution for enough fixations, the dermis was excised and the CTM was exposed. Subsequently

---

Address correspondence to Kazuharu Mine  
8-35-1 Sakuragaoka, Kagoshima 890-8544, Japan  
Tel: 099-275-6112  
E-mail: mine@dent.kagoshima-u.ac.jp

to identification of the muscular slips, all specimens were observed macroscopically and photographed. The anatomical nomenclature of muscles in this study followed the *Nomina Anatomica Veterinaria Japonica*<sup>17)</sup> and Ura<sup>18)</sup>.

Preliminarily we inquired anew about the use of rat specimens for a member of the Animal Experimentation Committee of Kagoshima University, and obtained the answer that it is not necessary to submit the experimental plan to use of such recent dead body.

## Result

The thin muscular slips split from the external surface of the CTM in the ventrolateral side of the trunk and extended cranially (Figs.1, 2). Its origin strode over the boundary area between the humeroabdominalis (HA) and humerodorsalis (HD) muscles<sup>18)</sup>. The muscle bundles were sparse and thin, and entire contour was longish trapezoidal. The insertion was on the brachial fascia or within the subcutaneous tissue of the arm to end diffusely. The averages of 8 sides were 17 mm in width at the origin, 35 mm in maximum length, and 0.6 mm in thick at the center of origin. This slip covered the second and third thoracic mammary glands in female and the whitish adipose mass in male (Fig.1). Both organs passed through the CTM at a hiatus between the HA and HD. To confirm the supplying nerve to the slip, the caudal pectoral nerve arising from the brachial plexus and the cutaneous branches of the thoracic nerve were pursued intensively. However, we could not identify the innervation with clarity (Fig.3).

## Discussion

The muscular slip examined in this study has scarcely described in the past. It was separated from the external surface of the CTM. Ura<sup>18)</sup> referred such muscle bundle as to the superficial derivative layer of the CTM. We made literature retrieval again based on his viewpoint, and picked up four descriptions of similar muscles. First one is supramammary muscle of the carnivorous animals<sup>19-20)</sup>, which is homologous to the preputial muscle in male, covering the mammary gland. However, its origin is closely related to the external genital organs. The second is the muscle found in the hystricomorphic rodents, originating at the ventral surface of the CTM and extending to the shoulder<sup>21)</sup>. However, its insertion attaches to the proximal part of the humerus. The third is the muscle found in the axillary fold of a species of marsupials, *Chironectes minimus*. Although Ura<sup>18)</sup> interpreted it an anlage of the derivative layer, he was negative about the relationship between the development of the CTM and the mammary

gland. Lastly there is a short description on the muscular slip of the lemming. Meinertz<sup>22)</sup> reported the slip in a male Greenland lemming splitting off from the lateral side of the CTM and extending to the arm and shoulder. It covered the pad of fat and interlaced partly with the platysma. His description is brief, but it is the most similar to the muscular slips found in this study among the newly retrieved cases. In addition, we have also detected the slips of similar configuration in two laboratory mice (unpublished data). Thus, there is a possibility that the murine commonly possesses the muscle bundle of this type.

In this study, we could not identify the innervation of this slip. Nevertheless we would speculate based on positional relation that it is supplied by a branch of the caudal pectoral nerve running downward just inside the boundary area between the HA and HD (Fig.3). If it is true, the innervating nerve will be coursed recurrently directing cranialward along the muscular bundle. It should be continued to search for the supplying nerve to clear whether the slip is a derivative layer of the CTM or not.

This slip is very delicate and is almost covered over by the forelimb in normal body posture. Even allowing the rat in a supine position during a skin experiment, the slip may provoke some confusion to an unskilled practitioner such as misunderstanding of subcutaneous layers or bleeding from the mammary gland. Moreover, there is a possibility that the slip is injured to cause unnecessary contraction or to delay wound healing<sup>13-15)</sup>. Therefore, the anatomical knowledge about this muscular slip will probably diminish these inconveniences.

## Conclusions

In this study, it is suggested that the fine slips of cutaneous muscle extending from the ventrolateral trunk to the brachium exist not only in the rat but in the common murine animals. For elucidation of phylogenetic origin and its function, further investigation is necessary. Even so, an attention to the muscular slip will improve the procedure of the skin experiment associated with the axillary region of the rat.

## Acknowledgement

We wish to thank Dr. Shinichi Okuyama (Evolution and Disease Laboratory in Sendai, Miyagi Prefecture) for providing animals.

## References

- 1) Sugama J, Sanada H, Nakatani T, et al. Pressure-induced ischemic wound healing with bacterial inoculation in the




Fig. 1 Photographs of the fine muscular slip in the right side extending from the ventrolateral trunk to the brachium. a (external aspect) and b (the slip is picked up): female specimen showing the slip (arrows) and second and third mammary glands (black arrowheads). c: male specimen showing the slip (arrows) and adipose mass (white arrowheads).




Fig. 2 Schematic drawing of the cutaneous trunci muscle system with skeleton of right forelimb and xiphoid process. The mammary gland (asterisk) is partly hidden by the muscular slip (arrow). HA, humeroabdominalis; HD, humerodorsalis; PA, pectoralis abdominalis.




Fig. 3 Schematic drawing of the typical branching pattern in the caudal brachial plexus and a series of pectoralis muscles. Left half viewed from the dorsal side. Black star and circle indicate the origin of the muscular slip outside the HA and HD. C7-T1, ventral branch of the spinal nerves; Cam, medial antebrachial cutaneous nerve; M, median nerve; U, ulnar nerve; CdPN, caudal pectoral nerve; CrPN, cranial pectoral nerve; HA, humeroabdominalis; HD, humero-dorsalis; PA, pectoralis abdominalis; Ppf, pectoralis profundus; Psf, pectoralis superficialis.

- rat. *Wounds* 2005; 17: 157–168
- 2) Stekelenburg A, Oomens CW, Strijkers GJ. Compression-induced deep tissue injury examined with magnetic resonance imaging and histology. *J Appl Physiol* 2006; 100:1946–1954
  - 3) Taylor GI, Gianoutsos MP, Morris SF. The neurovascular territories of the skin and muscles: anatomic study and clinical implications. *Plast Reconstr Surg* 1994; 94: 1–36
  - 4) Taylor GI. The blood supply of the skin. In: Thorne CH, Bartlett SP, Beasley RW, et al. ed. *Grabb and Smith's Plastic Surgery*. 6th ed. Philadelphia: Lippincott Williams and Wilkins; 2007: 33–41
  - 5) McGuire PG, Howdieshell TT. Importance of the engraftment in flap revascularization: confirmation by laser speckle perfusion imaging. *J Surg Res* 2010; 164: e201–e212
  - 6) Greene EC. *Anatomy of the Rat*. New York: Hafner Publishing; 1959
  - 7) Hebel R, Stromberg MR. *Anatomy of the Laboratory Rat*. Baltimore: Williams and Wilkins; 1976
  - 8) Mine K, Shimotakahara R, Lee HY, et al. Dorsal origin of the cutaneous trunci muscle in the rat, with special reference to the tail attachment. *Bull Sch Health Sci Kagoshima Univ* 2016; 26: 9–12
  - 9) Mine K, Shimotakahara R, Lee HY, et al. Source artery of the dorso-cranial part of subcutaneous structures in the rat trunk. *Bull Sch Health Sci Kagoshima Univ* 2018; 28: 77–81
  - 10) Cooper G, Schiller AL. *Anatomy of the Guinea Pig*. Cambridge: Harvard University Press; 1975
  - 11) Kent G, Carr R. *Comparative Anatomy of Vertebrate*. 9th ed. New York: McGraw-Hill Science; 2000
  - 12) Diogo R. *Muscles of Vertebrates: Comparative Anatomy, Evolution, Homologies and Development*. Boca Raton, FL: CRC Press; 2010
  - 13) Kayano S. Why not perforator flap training models in rats? *J Plast Reconstr Aesth Surg* 2010; 63: e134–e135
  - 14) Bali U, Gungor M, Yoleri L. Lateral thoracic artery perforator-based flap: a new experimental model. *J Surg Res* 2016; 200: 738–742

- 15) Petruska JC, Barjer DF, Garraway SM, et al. Organizing of sensory input to the nociceptive-specific cutaneous trunk muscle reflex in rat, an effective experimental system for examining nociceptive and plasticity. *J Comp Neur* 2014; 522:1048–1071
- 16) Okuyama S, Okuyama JN, Okuyama JK, et al. The arterial circle of Willis of the mouse helps to decipher secrets of cerebral vascular accidents in the human. *Med Hypoth* 2004; 63: 997–1009
- 17) Japanese Association of Veterinary Anatomists. *Nomina Anatomica Veterinaria Japonica*, 3rd ed. Tokyo: Gakuso-sha; 2000
- 18) Ura R. Über die allgemeine Differenzierung der oberflächlichen Brustmuskeln mit besonderer Berücksichtigung der Hauttrumpfmuskeln der Säugetiere. *Mitt Med Ges Tokyo* 1937; 51: 216–288, 339–390
- 19) Langworthy OR. A morphological study of the panniculus carnosus and its genetical relationship to the pectoral musculature in rodents. *Am J Anat* 1925; 35: 283–302
- 20) Preuss von F, Budras KD. Die Mm. supramammarius und praepuitalis der Katze. *Anat Anz* 1968; 122: 315–323
- 21) Wood CA, Howland EB. The skin musculature of hystericognath and other selected rodents. *Anat Hist Embryol* 1977; 6: 240–264
- 22) Meinertz T. The skin musculature of the Greenland lemming, *Dicrostonyx groenlandicus* (Traill.). *Meddelelser om Groenland* 1941; 131(3):1–77