

## Enzymatic Synthesis and Characterization of 2-Deoxyamyloses


著者	NAKAMURA Shota, YAMAMOTO Kazuya, KADOKAWA Jun-ichi
journal or publication title	The Research Reports of the Faculty of Engineering, Kagoshima University
volume	62
page range	11-11
year	2020
URL	<a href="http://hdl.handle.net/10232/00031534">http://hdl.handle.net/10232/00031534</a>

## Enzymatic Synthesis and Characterization of 2-Deoxyamyloses

Shota Nakamura<sup>1</sup>, Kazuya Yamamoto<sup>1</sup>, and Jun-ichi Kadokawa<sup>1</sup>

### Abstract

Phosphorylase catalyzes enzymatic polymerization of  $\alpha$ -D-glucose 1-phosphate (Glc-1-P) as a monomer using a maltooligosaccharide primer to produce  $\alpha(1\rightarrow4)$ -glucan (amylose) [1]. Because of weak specificity for the recognition of substrates, phosphorylase recognizes several analogue substrates of Glc-1-P to give non-natural polysaccharides [2]. Recently, we found that different from potato phosphorylase, thermostable phosphorylase (from *Aquifex aeolicus* VF 5) has ability to recognize D-glucal as a monomer for polymerization. In this study, the synthesis of 2-deoxyamylose was investigated by thermostable phosphorylase-catalyzed enzymatic polymerization of D-glucal via the in-situ production of  $\alpha$ -2-deoxy-D-glucose 1-phosphate (2dGlc-1-P) (Scheme 1). The enzymatic copolymerization of D-glucal with Glc-1-P was also carried out. The produced heteropolysaccharide formed a flexible film.



**Scheme 1.** Thermostable phosphorylase-catalyzed enzymatic polymerization of D-Glucal via in-situ production of 2dGlc-1-P.

### References

- [1] G. Ziegast, B. Pfanemüller, *Carbohydr. Res.*, **160**, 185 (1987).
- [2] J. Kadokawa, *Curr. Org. Chem.*, **21**, 1192 (2017).

<sup>1</sup> Department of Chemistry, Biotechnology, and Chemical Engineering, Graduate School of Science and Engineering, Kagoshima University, Kagoshima 890-0065, Japan