

Clinical Significance of 18F-fluorodeoxyglucose Positron Emission Tomography in Superficial Esophageal Squamous Cell Carcinoma

著者	KITA Yoshiaki, OKUMURA Hiroshi, UCHIKADO Yasuto, SASAKI Ken, OMOTO Itaru, MATSUMOTO Masataka, SETOYAMA Tetsuro, TANOUE Kiyonori, MORI Shinichiro, OWAKI Tetsuhiro, ISHIGAMI Sumiya, UENO Shinichi, KAJIYA Yoriko, NATSUGOE Shoji
journal or publication title	Annals of Surgical Oncology
volume	20
number	5
page range	1646-1652
year	2013
URL	http://hdl.handle.net/10232/21219

doi: info:doi/10.1245/s10434-012-2796-2

Table 3**Univariate and multivariate analysis of prognostic factor**

Factors	N	5-years survival rate (%)	Univariate	Multivariate	
			<i>p</i>	Hazard ratio (95% confidence interval)	<i>p</i>
FDG uptake					
Negative	45	87.3	0.041	1.50 (0.68 - 3.82)	0.329
Positive	35	48.3			
Histology					
Well/moderate	72	76.2	0.0001	3.48 (1.57 - 7.70)	0.002
Poor	8	0			
Depth of tumor invasion					
M1-SM1	40	90.2	0.039	0.97 (0.32 - 3.13)	0.96
SM2-SM3	40	51.4			
Lymph node metastasis					
Negative	64	76.9	0.046	1.57 (0.77 - 3.41)	0.218
Positive	16	37.9			
Lymphatic invasion					
Negative	61	73.1	0.379		
Positive	19	50.6			
Vascular invasion					
Negative	56	74.4	0.14		
Positive	24	54.2			
Glut-1 expression in cell membrane					
Negative	56	59.9	0.678		
Positive	24	74.6			