

2006 ANNUAL CONTRIBUTIONS

Author Index

- 1) Abe,T., A. Kurihara, S. Kawaguchi, R. Terada, and M. Masuda (2006). Preliminary report on the molecular phylogeny of *Laurencia complex*.*Coast.Sci.*, **30**: 209-213.
which contains waste stillage from a sweet potato shochu distillery. *Bull. Kagoshima Sci. Scholarship Found.*, **16**: 41-45 (in Japanese).
- 2) Alcantara, L. and T. Noro (2006). Growth of the abalone *Haliotis diversicolor* (Reeve) fed with macroalgae in floating net cage and plastic tank. *Aquacult. Res.*, **37**:708-717.
- 3) Anraku, K., G.Kawamur, M. Nakahara, N. Shigesato, and M.Vazquez Archdale (2006). Fish behaviour control methods in marine ranching in Japan-I. Development of conditioning method on the basis of hearing ability and auditory behaviour, in "Proceedings of the UMS-INOC International Conference on Coastal Oceanography & Sustainable Marine Aquaculture (ICCOSMA) - Confluence & Synergy, May 02-04,2006, Kota Kinabalu, Sabah, Malaysia", (ed.by Chouikhi, A., S. Shahbudin, M.A. Hoque, and M.S. Sitti Raehanah), Inter Islamic Science & Technology Network on Oceanography, Turkey, pp.365-371.
- 4) Anraku,K., T. Makino, F. Okawa, K. Watanabe, S. Masu, H. Ozono, H. Takeshita, G. Kawamura. and M.Vazquez Archdale (2006). Fish behaviour control methods in marine ranching in Japan-III. Development of Fish Guidance Device, in "Proceedings of the UMS-INOC International Conference on Coastal Oceanography & Sustainable Marine Aquaculture (ICCOSMA) - Confluence & Synergy, May 02-04, 2006, Kota Kinabalu, Sabah, Malaysia", (ed. by Chouikhi, A., S. Shahbudin, M.A. Hoque, and M.S. Sitti Raehanah), Inter Islamic Science & Technology Network on Oceanography, Turkey, pp.381-390.
- 5) Bei, J.X., H. Suetake, K. Araki, K. Kikuchi, Y. Yoshiura, H.R. Lin, and Y. Suzuki (2006). Two interleukin (IL)-15 homologues in fish from two distinct origins. *Mol. Immunol.*, **43**(7): 860-869.
- 6) Ebata, K., T. Yoshikawa, T. Sakata, S. Okano, S. Satou, K. Yoshinaga, M. Marugi, K. Tani, S. Uno, and J. Koyama (2006). Characteristic of bacteria on the surface of mortar
- 7) Hosotani, K. (2006). Behavior of the warm ocean water at the south coast of Kyushu. *J. Visual.*, **9**(4): 427-436.
- 8) Hosotani, K. and H. Kikukawa (2006). Behavior of the warm water filament touched with the south coast of Kyushu. *J. Remote Sens. Soc. Jpn.*, **26**(3): 224-233.
- 9) Hosotani, K., H. Kikukawa, and R. Nishi (2006). Analysis of the Dominant Coherent Structures around Kuroshio Using Satellite SST Images, in "Proceedings of the UMS-INOC International Conference on Coastal Oceanography & Sustainable Marine Aquaculture (ICCOSMA) - Confluence & Synergy, May 02-04, 2006, Kota Kinabalu, Sabah, Malaysia", (ed. by Chouikhi, A., S. Shahbudin, M. A. Hoque, and M. S. Sitti Raehanah), Inter Islamic Science & Technology Network on Oceanography, Turkey, pp. 465-474.
- 10) Hosotani, K., T. Suzuki, and Y. Ochiai (2006). Reconstruction of the complex flow measured by PIV -The vertical cross-flow over tube banks-. *J. Visual. Soc. Jpn.*, **26** (12): 114-122 (in Japanese).
- 11) Hossain, M.Y., Z.F. Ahmed, P.M. Leunda, A.K.M.R. Islam, S. Jasmine, J. Oscoz, R. Miranda, and J. Ohtomi (2006). Length-weight and length-length relationships of some small indigenous fish species from the Mathabhangha River, southwestern Bangladesh. *J. Appl. Ichthyol.*, **22**(4): 301-303.
- 12) Hossain, M.Y., Z.F. Ahmed, P.M. Leunda, S. Jasmine, J. Oscoz, R. Miranda, and J. Ohtomi (2006). Condition, length-weight and length-length relationships of the Asian striped catfish *Mystus vittatus* (Bloch, 1794) (Siluriformes: Bagridae) in the Mathabhangha River, southwestern Bangladesh, *J. Appl. Ichthyol.*, **22**(4): 304-307.
- 13) Inoue, Y., T. Murayama, and A. Okino (2006). Monster

- jellyfish excluder trawl net frames. ICES 2006 Boston, Fishing Tech. in the 21th Century Sympo. Proceedings, pp. 110.
- 14) Iqbal, K.M., Y. Masuda, H. Suzuki, and A. Shinomiya (2006). Age and growth of the Japanese silver-biddy, *Gerres equulus*, in western Kyushu, *Jpn. Fish. Res.*, **77**: 45-52.
- 15) Kakinoki, K., S. Imawaki, H. Uchida, H. Nakamura, S. Umatani, A. Nishina, H. Ichikawa, and M. Wimbush (2006). Sea-surface dynamic height of the Kuroshio south of Japan estimated from inverted echo sounders and its error estimation. *Rep. Res. Inst. Appl. Mech., Kyushu University*, No. 130: 11-20 (in Japanese).
- 16) Kadokami, S. and Y. Kitadai (2006). Advantages of environmentally sound poly-eco-aquaculture in fish farms, in "Proceedings of the FFTC-ASPAC/RCA International Workshop on Innovative and Eco-friendly Technologies for the Production of Safe Aquaculture Foods, Dec 4-8, 2006, Denpasar, Bali, Indonesia", (ed. by Hu, S.H. and I.W. Lee), pp. 181-192.
- 17) Kawaguchi, S., S. Shimada, T. Abe, and R. Terada (2006). Morphological and molecular phylogenetic studies of a red alga, *Halymenia durvillei*, (Halymeniaceae, Halymeniales) from Indo-Pacific. *Coast. Sci.*, **30**: 201-208.
- 18) Kawamura G., K. Anraku, T. Nishi, M. Nakahara, and Y. Machida (2006). Fish behaviour control methods in marine ranching in Japan-II. Improved technique for conditioning to sound, in "Proceedings of the UMS-INOC International Conference on Coastal Oceanography & Sustainable Marine Aquaculture (ICCOSMA) – Confluence & Synergy, May 02 – 04, 2006, Kota Kinabalu, Sabah Malaysia", (ed by Chouikhi, A., S. Shahbudin, M.A. Hoque, and M.S. Sitti Raehanah), Inter Islamic Science & Technology Network on Oceanography, Turkey, pp. 372-380.
- 19) Kawamura G., K. Anraku, T. Nishi, H. Yoshida, N. Shigesato, and M. Nakahara (2006). Fish behaviour control methods in marine ranching in Japan-IV. Effect of flicking light on negatively phototactic red sea bream *Pagrus major*, in "Proceedings of the UMS- INOC International Conference on Coastal Oceanography & Sustainable Marine Aquaculture (ICCOSMA) – Confluence & Synergy, May 02-04, 2006, Kota Kinabalu, Sabah, Malaysia", (ed. by Chouikhi, A., S. Shahbudin, M.A. Hoque, and M.S. Sitti Raehanah), Inter Islamic Science & Technology Network on Oceanography, Turkey, pp. 391-397.
- 20) Kobari, T., K. Shimizu, S. Nishikawa, and A. Habano (2006). Seasonal variation in abundance of Oithonidae copepods in the Kagoshima Bay. *Mem. Fac. Fish. Kagoshima Univ.*, **55**: 61-68 (in Japanese).
- 21) Kobari, T. (2006). Plankton in the Kinko Bay. Kinko Bay Expedition. Society of Research of the Kinko Bay, 5 (in Japanese).
- 22) Kobari, T. (2006). Workshop on "Oceanic Ecodynamics Comparison in the Subarctic Pacific (OECOS)": Growth of large interzonal migrating copepods. *PICES Sci. Rept.*, PICES Secretariat Office, **32**: 68-70.
- 23) Koyama, J., S. Imai, K. Fujii, S. Kawami, C.K. Yap, and A. Ismail (2006). Pollution by estrogens in river and estuarine waters around Kuala Lumpur, Malaysia, and their effects on the estuarine Java-medaka, *Oryzias javanicus*. *Jpn. J. Environ. Taxicol.*, **9**: 141-148.
- 24) Michael, F.R., S. Koshio, S. Teshima, M. Ishikawa, and O. Uyan (2006). Effect of choline and methionine as methyl group donors on juvenile kuruma shrimp, *Marsupenaeus japonicus* Beta. *Aquacult.*, **258**: 521-528.
- 25) Mizuno, T. and H. Kikukawa (2006). Three dimensional FEM analysis of the tide in Kagoshima Bay. *Mem. Fac. Fish. Kagoshima Univ.*, **55**: 13-26.
- 26) Nakamura, H., T. Yamashiro, A. Nishina, and H. Ichikawa (2006). Time-frequency variability of Kuroshio meanders in Tokara Strait. *Geophys. Res. Lett.*, **33**, L21605, doi:10.1029/2006GL027516.
- 27) Nakashima, T., Y. Inoue, and T. Shiroyama (2006). Discards in sail-trawl fishery and their survivals in Yatsushiro sound, Japan. *ICES 2006 Boston, Fishing Tech. in the 21th*

- Century Sympo. Proceedings*, pp.116.
- 28) Nguyen, D.T.P., H. Maeda, Y. Taoka, M. Nakashita, M. Hidaka, I. Yoshikawa, and T. Sakata (2006). Chemical and microbiological characterization of the recent surface sediments in Kagoshima Bay and its significance to organic matter accumulation. *J. Sedimentolog. Soc. Japan*, **63**: 3-15.
- 29) Nguyen, D.T.P., T. Yoshikawa, M. Hidaka, H. Maeda, and T. Sakata (2006). Isolation and characterization of sulfate-reducing bacteria from sediments of Kagoshima Bay. *Mem. Fac. Fish. Kagoshima Univ.*, **55**: 69-78.
- 30) Nishi, R. (2006). Visualized rip currents. *Q. J. The Suiro (Hydrography)*, **34**(1): 21-26 (in Japanese).
- 31) Nishi, R. (2006). Safe utilization of coastal area. *Q. J. The Suiro (Hydrography)*, **35**(1): 27-33 (in Japanese).
- 32) Nishi, R. (2006). Field observation on a rip current. *Q. J. The Suiro (Hydrography)*, **35**(2): 21-27 (in Japanese).
- 32) Nishi, R. (2006). Numerical simulation of a rip current. *Q. J. The Suiro (Hydrography)*, **35**(3): 18-24 (in Japanese).
- 34) Nishi, R., N.C. Craus, and A. Kawamori (2006). Fundamental Study on Inlet Geomorphology – An Approach for a Global A-P Relationship, in “Proceedings of Civil Engineering in the Ocean” JSCE, Vol. 22, p. 927-932 (in Japanese).
- 35) Nishi, R., S. Futatsumachi, H. Ito, T. Miyake, A. Nagayama, and A. Ohtani (2006). Fundamental study on safe utilization in a coral reef region. *Ann. J. Coast. Eng.*, *JSCE*, **53**: 111-115 (in Japanese).
- 36) Nishi, T. and G. Kawamura (2006). Magnetosensitivity in the darkbanded rockfish *Sebastes inermis*. *Nippon Suisan Gakkaishi*, **72**(1): 27-33 (in Japanese).
- 37) Nishi, T. (2006). Studies on the magnetosensitivity and magnetocompass orientation in the Japanese eel *Anguilla japonica*. *Mem. Fac. Fish. Kagoshima Univ.*, **55**(2): 107-159 (in Japanese).
- 38) Ohtomi, J. and R. Sakata (2006). Reproductive biology of the grenadier *Coelorinchus jordani* (Gradiformes, Macrouridae) in Kagoshima-Bay, southern Japan. *Nippon Suisan Gakkaishi*, **72**(2): 210-216 (in Japanese).
- 39) Ohtomi, J. (2006). Population structure, condition factor and relative growth of the grenadier *Coelorinchus jordani* (Gradiformes, Macrouridae) in Kagoshima-Bay, southern Japan. *Nippon Suisan Gakkaishi*, **72**(2): 217-222 (in Japanese).
- 40) Ohtomi, J. (2006). Marine organisms in Kagoshima Bay, in “Kinkowan Tankentai” (ed. by Kinkowan Kenkyukai). Kagoshima, pp. 10-11 (in Japanese).
- 41) Ohtomi, J. and K. Ebata (2006). Fishery in Kagoshima Bay, in “Kinkowan Tankentai” (ed. by Kinkowan Kenkyukai). Kagoshima, pp. 14-15 (in Japanese).
- 42) Ohtomi, J., T. Furuta, and H. Kuwazoe (2006). Change in spatial desitribution with development fro larva Japanese mantis *Oratosquilla oratoria* (Stomatopoda, Suillidae) in Tokyo Bay, Japan. *Nippon Suisan Gakkaishi*, **72**(3): 382-389 (in Japanese).
- 43) Ren, T., S. Koshio, S. Teshima, M. Ishikawa, S. Yokoyama, F.R. Michael, and O. Uyan (2006). The efficiency of L-ascorbic acid in micro bound diet for larval red sea bream (*Pagrus major*). *Aquacult. Sci.*, **54**: 561-566.
- 44) Shimabukuro, H., S. Arai, T. Terawaki, and T. Noro (2006). On the record and distribution of *Sargassum carpophyllum* (Phaeophyceae, Fucales) from Japan. *Jpn. J. Phycol. (Sorui)*, **54**: 85-88 (in Japanese).
- 45) Shiozaki, Y., H. Maeda, Y. Yaoka, S. Koshio, T. Yoshikawa, and T. Sakata (2006). The screening of new probiotics-candidate bacteria in aquaculture. *Bokin Bobai*, **34**: 251-261 (in Japanese).
- 46) Shiozaki, Y., Y. Taoka, T. Yoshikawa, and T. Sakata (2006). Characterization of Bacillus sp. PU-T8 isolated from tiger puffer fish *Takifugu rubripes* as probiotics. *Mem. Fac. Fish. Kagoshima Univ.*, **55**: 79-86.

- 47) Shiromoto, T., B. Chokesanguan, and Y. Inoue (2006). Trash fish and discards of push net catch in Thailand. *Mem. Fac. Fish. Kagoshima Univ.*, **55**: 43-50 (in Japanese).
- 48) Suetake, H., N.R. Saha, K. Araki, K. Akatsu, and Y. Suzuki (2006). Lymphocyte surface marker genes in fugu. *Com. Biochem. Physiol. Part D*, **1**(1): 102-108.
- 49) Takada, T., Y. Taoka, A. Yamamoto, and K. Naiki (2006). Effect of total genomic DNA from probiotic bacteria on non-specific immune response of Japanese flounder, *Paralichthys alovaceus*. *Aquacult. Sci.*, **54**(4): 521-524 (in Japanese).
- 50) Tanaka, R., R. Nishi, T. Yuki, and S. Futatsumachi (2006). Photographic survey in a sub-aquatic region applying commercially available digital camera, in "Proceedings of Civil Engineering in the Ocean", JSCE, Vol. 22, pp. 911-915 (in Japanese).
- 51) Taoka, Y., H. Maeda, J. -Y. Jo, S. -M. Kim, S. -I. Park, T. Yoshikawa, and T. Sakata (2006). Use of live and dead probiotic cells in tilapia *Oreochromis niloticus*. *Fish. Sci.*, **72**: 755-766.
- 52) Taoka, Y., H. Maeda, J. -Y. Jo, M. -J. Jeon, S.C. Bai, W. -J. Lee, K. Yuge, and S. Koshio (2006). Growth, stress tolerance and non-specific immune response of Japanese founder *Paralichthys olivaceus* to probiotics in a closed re-circulating system. *Fish. Sci.*, **72**: 310-321.
- 53) Terada, R. and C.M. Smith (2006). Increase of alien seaweed in Hawaiian Islands. *Jpn. J. Phycol.*, **54**: 101-104 (in Japanese).
- 54) Teshima, S., S. Koshio, M. Ishikawa, M.S. Alam, and L.H.H. Hernandez (2006). Protein requirements of the freshwater prawn *Macrobrachium rosenbergii* evaluated by the factorial method. *J. World Aquacult. Soc.*, **37**: 145-153.
- 55) Uyan, S., G. Kawamura, and M. Vazques Archdale (2006). Morphology of the sense organs in larval anchovy *Engraulis japonicus* caught by midwater trawl. *Aquacult. Sci.*, **54**: 275-282.
- 56) Uyan, S., G. Kawamura, and M. Vazques Archdale (2006). Morphology of the sense organs of anchovy *Engraulis japonicus*. *Fish. Sci.*, **72**: 540-545.
- 57) Uyan, O., S. Koshio, S. Teshima, M. Ishikawa, M. Thu, M.S. Alam, and F.R. Michael (2006). Growth and phosphorus loading by partially replacing fishmeal with tuna muscle by-product powder in the diet of juvenile Japanese flounder, *Paralichthys olivaceus*. *Aquacult.*, **257**: 437-445.
- 58) Vazques Archdale, M., C.P. Añasco, and H. Shinji (2006). Comparative fishing trials for invasive crabs *Charybdis japonica* and *Portunus pelagicus* using collapsible pots. *Fish. Res.*, **82**: 50-55.
- 59) Vazques Archdale, M., L. Kariyazino, and C.P. Añasco (2006). The effect of two pot types on entrance rate and entrance behavior of the invasive Japanese swimming crab *Charybdis japonica*. *Fish. Res.*, **77**: 271-274.
- 60) Yamamoto, A. (2006). Bacterial coldwater disease, in "New Atlas of Fish Disease" (ed. by Hatai, K. and K. Ogawa). Midori Shobou, Tokyo, p. 25 (in Japanese).
- 61) Yamamoto, A. (2006). Water molds infection in salmonid eggs, in "New Atlas of Fish Disease" (ed. by Hatai, K. and K. Ogawa). Midori Shobou, Tokyo, p. 29 (in Japanese).
- 62) Yigit, M., M. Erdem, S. Koshio, S. Ergun, A. Turker, and B. Karaalli (2006). Substituting fish meal with poultry by-product meal in diets for black sea turbot *Psetta maeotica*. *Aquacult. Nutr.*, **12**: 1-8.
- 63) Yokoyama, S., S. Koshio, N. Takakura, K. Shima, M. Ishikawa, F.J. Gallardo-Cigarroa, M.R. Catacutan, and S. Teshima (2006). Effect of dietary bovine lactoferrin on growth response, tolerance to air exposure and low salinity stress conditions in arrange spotted grouper *Epinephelus coioides*. *Aquacult.*, **255**: 507-513.
- 64) Winarso, G., K. Hosotani, and H. Kikukawa (2006). Chlorophyll a distribution deduced from MODIS ocean

- color data and its characteristics around Hyuganada. *Mem. Fac. Fish. Kagoshima Univ.*, **55**: 13-26.
- 65) Winarso, G., H. Kikukawa, and T. Nagamatsu (2006). Determination of the oceanographic flow fields using ocean color Modis data, in “Proceedings of the UMS-INOC International Conference on Coastal Oceanography & Sustainable Marine Aquaculture (ICCOSMA) – Confluence & Synergy, May 02-04, Kota Kinabalu, Sabah, Malaysia”, (ed. by Chouikhi, A., S. Shahbudin, M.A. Hoque, and M.S. Sitti Raehanah), Inter Islamic Science & Technology Network on Oceanography, Turkey, pp. 495-501.