

Comparative Study of Leucine Aminopeptidases from Marine Labyrinthulid and Thraustochytrid Strains

Md. Iftekharul Wahid¹, Sadaba R. Bitoon², Tetsuya Fukunaga¹, Takeshi Yoshikawa¹ and Taizo Sakata^{*}

Key words: labyrinthulid; thraustochytrid; leucine aminopeptidase; inhibitor

Abstract

Leucine aminopeptidases (LAPs) from marine labyrinthulid strain 00-Bat-05 and thraustochytrid strain HR-3 cells were partially purified and characterized by enzymological properties. The optimum temperature of LAPs from both strains was 37°C. The thermostability of 00-Bat-05 LAP was indicated by having 80% of maximum activity after heat treatment at 60°C for 10 min, while LAP activity of strain HR-3 was completely inactivated at 60°C. LAP activities from both strains were high at near pH 8.0. Both LAPs were inactivated by 1,10-phenanthroline, *p*-chloromercuribenzoic acid (PCMB), bestatin and sodium dodecyl sulphate (SDS). Enzyme activity of LAP from strain 00-Bat-05 was stimulated by Co²⁺ and inhibited by Zn²⁺, while that from HR-3 was inhibited by Co²⁺ and Zn²⁺. LAP of 00-Bat-05 had a high specificity for L-leucine-*p*-nitroanilide but HR-3 enzyme showed relatively broad specificity for *p*-nitroanilide derivatives of L-amino acids.

Labyrinthulids (Family Labyrinthulaceae) and thraustochytrids (Family Thraustochytriaceae) are the main two groups of the order Labyrinthulales and play the peculiar roles in marine food web systems.^{1,2)} Labyrinthulids are slime mold-like protists and distributed widely in estuarine and near-shore marine habitats throughout the world. In mangrove brackish water environments, *Labyrinthula* spp. live on mangrove leaves, marine vascular plants and benthic algae, and also parasitizes fungal hyphae, which are decomposing plant matter. *Labyrinthula* spp. play an important role in mangrove ecosystem by fixing plant nutrients in a form of hyphae or cells of the organisms and converting them to a nutritious food for plant consuming animals.³⁾ The ectoplasmic networks of *Labyrinthula* spp. are capable of decomposing many different microbes including bacteria, yeast, diatoms, and fungal hyphae.⁴⁾ Thraustochytrids are a group of eukaryotic marine microorganisms, which can play dual roles in nature as bacterial feeders when in an amoeboid form, and organic material degraders in their thallus form.⁵⁾ In mangrove environments, these marine protists are found to be associated with decaying mangrove leaves and decompose mangrove leaves by their degradative enzymes and also to be capable of breaking down several complex

organic substrates.^{6,7)}

In addition, these organisms also deserve more attention in aquaculture because they accumulate high concentrations of polyunsaturated fatty acids such as docosahexaenoic acid (DHA) and eicosapentanoic acid (EPA) which are essential nutrients for many fish and crustaceans.⁸⁻¹¹⁾ Thraustochytrids are used practically as a feed for zooplankton such as rotifers and *Artemia* because of improving the nutritional value of the food animals.¹²⁾

In natural ecosystems, the recycling of nutrients, especially protein degradation is important to maintain all living things and aminopeptidases are found widely distributed among both prokaryotic and eukaryotic organisms, which catalyze the release of amino acid residues from the N-terminal end of proteins.¹³⁾ Leucine aminopeptidase (LAP, EC 3.4.11.1) activity in marine environments indicated an ecological role by bacteria in aquatic biogeochemical cycles.¹⁴⁾ LAP was partially purified from pig intestinal mucosa at first and found to hydrolyze a wide variety of peptides. The LAP activities from swine kidney and *Aspergillus sojae* was determined to be activated by divalent cations like Zn²⁺ and Co²⁺.^{15,16)}

The objective of this study was to characterize and compare the leucine aminopeptidase (LAP) from marine

¹Laboratory of Microbiology, Faculty of Fisheries, Kagoshima University, Kagoshima 890-0056, Japan.

²University of the Philippines in the Visayas, Miyagao, Iloilo, Philippines

*Corresponding author, E-mail: sakata@fish.kagoshima-u.ac.jp

labyrinthulid and thraustochytrid isolates in order to assess nutritional process and degradation activity of these organisms in marine environments.

Materials and Methods

Labyrinthulid and thraustochytrid strains

Labyrinthulid strain 00-Bat-05 was isolated from mangrove leaves at the Batan Bay mangrove area, the Philippines, using a double layer medium containing live diatom cells.¹⁷⁾ Thraustochytrid strain HR-3 was provided by Dr. M. Hayashi, Faculty of Agriculture, Miyazaki University, Japan and which was isolated from coastal area of Hiroshima Prefecture, Japan. Chromosomal DNA of the strains was extracted and 18S rDNAs were amplified by PCR and sequenced by the standard method. A phylogenetic tree inferred from 18S rDNA nucleotide sequences was constructed by the neighbor-joining method.¹⁸⁾

Media and growth conditions

For the enzyme production, labyrinthulid strain 00-Bat-05 was grown at 25°C for 168 h in 1 l of the ES (Provasoli's enrichment seawater) liquid medium in a 2 l Erlenmeyer flask with containing bacterial extract-egg yolk agar medium (NSBEY agar) according to Wahid *et al.*¹⁹⁾ Thraustochytrid strain HR-3 was cultured at 25°C for 96 h in 1 l of the medium containing 30 g glucose, 3 g yeast extract, 5 g polypeptone and 750 ml of artificial seawater (ASW, Herbst's formula composed of NaCl 30.0 g, KCl 7.0 g, MgCl₂·6H₂O 10.8 g, MgSO₄·7H₂O 5.4 g and CaCl₂·2H₂O 1.0 g per l, pH 7.6) in a flask on a reciprocal shaker (Taitec, NR-3, Japan) at 120 rpm.

LAP enzyme preparation from the test organisms

1. Labyrinthulid strain 00-Bat-05

After incubation at the same conditions as above, the cells of labyrinthulid strain 00-Bat-05 collected from 8 l of liquid culture were washed with 50 mM Tris-HCl buffer (pH 7.6) and then suspended in 40 ml of 50 mM Tris-HCl buffer. The suspended cells were disrupted by sonication (4280S, Kaijo Denki, Japan) for 5 min, the cell debris was removed by centrifugation (12,000 × g for 20 min) and the supernatant was filtrated by 0.45 μm

membrane filter (Advantec, Japan) and flow-through solution was used as a cell-free extract. Solid ammonium sulfate was added to the cell-free extract solution to give 50% saturation. After 2 h, the precipitate was removed and the enzyme protein fraction was precipitated from the supernatant with ammonium sulfate (90% saturation) and collected by centrifugation. The precipitate was dissolved with 50 mM Tris buffer and dialyzed overnight against 50 mM Tris buffer. The dialyzed enzyme solution was loaded onto a Toyopearl DEAE-650M column (Tosoh, Japan) previously equilibrated with 50 mM Tris buffer and the proteinous materials were eluted with 50 mM Tris buffer and 500 mM NaCl using a linear gradient. The active fractions were pooled and concentrated by ultrafiltration with a YM-3 membrane (Amicon, USA). The enzyme solution was then applied to a Toyopearl HW-55F column (Tosoh, Japan) equilibrated with 50 mM Tris buffer containing 100 mM NaCl and eluted with the same buffer. The active enzyme fractions were combined, concentrated by ultrafiltration with a YM-3 membrane.

2. Thraustochytrid strain HR-3

After incubation at the same conditions as above, the cells of thraustochytrid strain HR-3 collected from 500 ml of culture broth were washed with 50 mM Tris-HCl buffer (pH 7.6) and then suspended in 50 ml of the same buffer. The suspended cells were disrupted by sonication and the cell debris was removed by centrifugation and the supernatant was filtrated by 0.45 μm membrane filter. Solid ammonium sulfate was added to the crude enzyme solution to give 60% saturation. After 4 h, the precipitate was collected by centrifugation, dissolved with 50 mM Tris buffer and dialyzed overnight against 50 mM Tris buffer. The dialyzed enzyme solution was loaded onto a Toyopearl Super Q-650M column (Tosoh, Japan) previously equilibrated with 50 mM Tris buffer and the proteinous materials were eluted with 50 mM Tris buffer and 500 mM NaCl using a linear gradient. The active fractions of LAP enzyme were pooled and concentrated by freeze drying. The freeze dried enzyme preparation was suspended in 50 mM Tris-HCl buffer and then applied to a Toyopearl HW-55F column (Tosoh, Japan) equilibrated with 50 mM Tris-HCl buffer containing 150 mM NaCl and the proteins were eluted with the same buffer. The

active enzyme fractions were combined and concentrated by freeze drying.

Enzyme assay

Aminopeptidase activity was spectrophotometrically assayed by using L-leucine-*p*-nitroanilide (Leu-*p*-NA) as substrate described by Chien *et al.*¹⁶⁾ The standard assay condition was as follows: 200 μ l reaction mixture consisted of 20 μ l of 10 mM L-leu-*p*-NA solution, 100 μ l of 50 mM Tris-HCl buffer (pH 7.6) and 80 μ l enzyme solution and incubated at 37°C for 2 h. In addition, *p*-nitroanilide (*p*-NA) derivatives binding a series of L-amino acids were used at final concentration of 1 mM to measure the relative activity for LAP enzymes. Absorbance at 405 nm was determined by micro titer plate reader, MPR-A4I (Tosoh, Japan). All data are expressed as average values of duplicate experiments.

Effect of reaction temperature

By using the standard reaction mixture, enzymatic activity was determined at different temperatures between 4 and 80°C for 2 h incubation.

Effect of heat treatment

Enzyme thermostability was determined after preincubation of the enzyme solution in 50 mM Tris-HCl buffer (pH 7.6) at various temperatures (4-80°C) for 10 min and then the solution was cooled on ice for 5 min followed by the standard assay condition to measure residual enzyme activity.

Effect of reaction pH

Enzyme preparations were preincubated in 50 mM various buffer at 30°C for 10 min. The optimum pH for enzyme reaction was examined under the standard assay condition using 50 mM buffer solutions with various pHs: sodium phosphate buffer (pH 6.0-7.0), Tris-HCl (pH 7.0-9.0), carbonate buffer (pH 9.0-11.0) and glycine buffer (pH 11.0-13.0).

Effect of inhibitors

The effect of *p*-chloromercuribenzoic acid (PCMB), phenylmethylsulfonyl fluoride (PMSF), ethylenediaminetetraacetic acid (EDTA), dithiothreitol (DTT), 1,

10-phenanthroline, iodoacetamide (IAA), bestatin and sodium dodecyl sulfate (SDS) on LAP activity was determined after preincubation with the enzyme and the chemicals (final concentration of 1 mM) in 50 mM Tris-HCl buffer (pH 7.6) for 30 min at 30°C. The remaining activity was assayed under the standard condition.

Effect of metal ions

After preincubating the enzyme with various cations, chelating reagents or chemical reagents in 50 mM Tris-HCl buffer (pH 7.6) for 30 min at 30°C. The remaining activity was assayed under the standard assay condition.

Chemicals used

The synthetic chromogenic substrates, L-leucine-*p*-nitroanilide (Leu-*p*-NA), L-alanine-*p*-nitroanilide (Ala-*p*-NA) and L-glutamine-*p*-nitroanilide (Glu-*p*-NA) were obtained from Peptide Institute, INC. (Osaka, Japan) and L-proline-*p*-nitroanilide (Pro-*p*-NA), L-methionine-*p*-nitroanilide (Met-*p*-NA), L-arginine-*p*-nitroanilide (Arg-*p*-NA) were from Sigma Chemical Co. (St. Louis, Mo., USA). The protease inhibitors used in this study were purchased from Sigma and Nakarai Tesuque (Kyoto, Japan).

Results

Phylogenetic analysis of strains 00-Bat-05 and HR-3

Figures 1 A and 1 B show cell morphology of labyrinthulid strain 00-Bat-05 and thraustochytrid strain HR-3, respectively. The individual cells of labyrinthulid 00-Bat-05 (Fig. 1A) were typically spindle form with averaged 2.5-5.0 μ m wide by 8.0-12.0 μ m long. On the other hand, those of thraustochytrid HR-3 (Fig. 1B) were round form with 10.0-15.0 μ m in diameter, which contained numerous granules in the cytoplasm. The 18S rDNA sequence of strain 00-Bat-05 (GenBank accession no. AB290459) showed 93% homology with that of *Labyrinthula* sp. AN-1565 (AB022105), while strain HR-3 had 99% homology with *Thraustochytriidae* sp. NIOS-1 haplotype NIOS1-D00-1 (AY705769). A phylogenetic tree constructed by the neighbor-joining method is shown in Fig. 2.




Fig. 1. Photograph of vegetative cells of labyrinthulid strain 00-Bat-05 (A) and thraustochytrid strain HR-3 (B). Scale bars indicate 10 µm.




Fig. 2. Phylogenetic tree inferred from 18S rDNA of labyrinthulid sp. 00-Bat-05 and thraustochytrid sp. HR-3 by the neighbour-joining method. Bootstrap values greater than 50% are shown above the branches. A scale bar indicates genetic distance. LPG, labyrinthulid phylogenetic group; TPG, thraustochytrid phylogenetic group.

Enzyme preparation

Partially purified enzyme preparations were obtained after ion-exchange (Fig. 3) and gel filtration chromatography. One active fraction of LAP from each test strain was isolated although it still included several protein molecules as judged from polyacrylamide gel electrophoresis (PAGE). In this experiment, the partially purified enzymes were used for characterization of LAP activity.




Fig. 3. Chromatographic profiles of LAP enzymes from test strains. A, Toyopearl DEAE-650M chromatogram of LAP from strain 00-Bat-05; B, Toyopearl Super Q-650M chromatogram of LAP from strain HR-3. ●: LAP activity (OD at 405 nm), —: protein concentration (OD at 280 nm), ---: NaCl concentration (M).

Characterization of LAP enzymes

1. Optimal temperature

LAP activities of both strains were at their maxima at 37°C as shown in Fig. 4. More than 50 % of maximum activity was found between 15 to 45°C in both strains. More than 80% activity of 00-Bat-05 LAP was remaining after heat treatment at 60°C for 10 min, while HR-3 LAP enzyme was almost inactivated with heat treatment at the same temperature (Fig. 5).

2. Optimal pH

As shown in Fig. 6, the optimum pH was 8.0 for 00-Bat-05 LAP and 7.0-8.0 for HR-3 LAP. However, the enzyme activity of 00-Bat-05 strain was found to be considerably low at pH 9.0.




Fig. 4. Effect of reaction temperature on enzyme activity from strains 00-Bat-05 (A) and HR-3 (B). Enzyme activity was measured at various temperatures in 50 mM Tris-HCl buffer (pH 7.6). Activities are expressed relative to the maximum value.

3. Effect of inhibitors

Effect of various inhibitor substances on LAP activity is shown in Table 1. In strain 00-Bat-05, the LAP activity was strongly inhibited by bestatin, which is generally known to be an inhibitor of aminopeptidase (Table 1). Strong inhibition was also observed in the presence of EDTA, SDS, PCMB, DTT, and 1,10-phenanthroline. EDTA was known to be a metal-chelating agent, and PCMB to be a SH inhibitor. On the other hand, LAP activity from HR-3 strain was inhibited by bestatin, SDS, PCMB, and 1,10-phenanthroline.

4. Effect of metal ions

The effect of several metal ions on enzyme activity is shown in Table 2. Among the metal ions tested in LAP of strain 00-Bat-05, enzyme activity was strongly inhibited




Fig. 5. Effect of heat treatment on enzyme stability of strains 00-Bat-05 (A) and HR-3 (B). Enzyme preparations were heated in 50 mM Tris-HCl buffer (pH 7.6) at various temperatures for 10 min

by Zn^{2+} , Hg^{2+} and Ni^{2+} and moderately inhibited by Mn^{2+} , Mg^{2+} , Cu^{2+} , Ba^{2+} and K^+ . However, Co^{2+} strongly activated the enzyme activity from strain 00-Bat-05. On the other hand, the LAP activity of strain HR-3 was strongly inhibited by Zn^{2+} , Hg^{2+} and Ni^{2+} , and slightly inhibited by Co^{2+} . Metal ions such as Na^+ , K^+ , and Mg^{2+} stimulated the LAP activity of strain HR-3.




Fig. 6. Effect of pH on enzyme activity of strains 00-Bat-05 (A) and HR-3 (B). Enzyme preparations were preincubated in 50 mM various pH buffer at 30°C for 10 min. ◆: sodium phosphate buffer, ○: Tris-HCl buffer, ▲: carbonate buffer, ×: glycine-NaOH buffer.

Table 1. Effect of chemical reagents on LAP activity from test strains

Chemical reagents ¹	Relative activity (%)	
	00-Bat-05	HR-3
No addition	100.0	100.0
EDTA ^{*2}	40.2	88.0
PCMB ^{*2}	13.6	10.0
DTT ^{*2}	16.4	78.0
IAA ^{*2}	132.4	91.4
1,10-phenanthroline	12.8	16.0
PMSF ^{*2}	106.8	89.0
SDS ^{*2}	3.4	17.5
Bestatin	5.2	35.7

¹Each chemical reagent was added to a final concentration of 1 mM in the reaction mixture.

²EDTA: ethylenediaminetetraacetic acid, PCMB: *p*-chloromercuribenzoic acid, DTT: dithiothreitol, IAA: iodoacetamide, PMSF: phenylmethylsulfonyl fluoride, SDS: sodium dodecyl sulfate.

Table 2. Effect of metal ions on LAP activity from test strains

Metal ions ¹	Relative activity (%)	
	00-Bat-05	HR-3
No addition	100.0	100.0
NaCl	108.5	187.8
KCl	55.8	155.7
MnCl ₂	76.0	104.3
MgCl ₂	62.8	178.8
ZnCl ₂	31.0	29.7
CuSO ₄	52.7	100.0
CaCl ₂	79.9	99.5
HgCl ₂	30.2	23.1
NiCl ₂	12.9	46.7
CoSO ₄	298.6	75.0
BaCO ₃	55.5	87.3

¹ Each salt was added to a final concentration of 1 mM in the reaction mixture.

5. Substrate specificity

The ability of the enzyme to catalyze the hydrolysis of various *p*-NA derivatives is shown in Table 3. In strain 00-Bat-05, Leu-*p*-NA was most efficiently hydrolyzed by LAP. Met-*p*-NA, Ala-*p*-NA and Arg-*p*-NA were also to be considerably hydrolyzed, while Pro-*p*-NA and Glu-*p*-NA were practically resistant to the action of the enzyme. In strain HR-3, Ala-*p*-NA was most efficiently hydrolyzed by the enzyme and Leu-*p*-NA, Met-*p*-NA and Arg-*p*-NA were good substrate for LAP. On the other hand, Pro-*p*-NA and Glu-*p*-NA were not optimal to the enzyme activity from stain HR-3.

Table 3. Hydrolytic activities of LAP from test strains on various substrates

Substrate ¹	Relative activity (%)	
	00-Bat-05	HR-3
Leu- <i>p</i> -NA	100.0	100.0
Ala- <i>p</i> -NA	21.0	347.0
Pro- <i>p</i> -NA	1.3	34.2
Met- <i>p</i> -NA	51.0	109.4
Arg- <i>p</i> -NA	20.1	125.0
Glu- <i>p</i> -NA	5.2	22.7

¹All substrates were used at a final concentration of 1 mM under the standard assay conditions.

Discussion

Both labyrinthulids and thraustochytrids have a role in decomposing plant matters in coastal environments and their degrading enzymes are involved in this process.¹¹

This study compared the enzymological properties of LAP enzymes from labyrinthulid and thraustochytrid strains isolated from marine environments.

The LAP enzyme of strain 00-Bat-05 had an optimal temperature of 37°C but 80% of its maximal activity remained even after heat treatment at 60°C for 10 min. On the other hand, the optimal temperature for the LAP activity of strain HR-3 was 37°C and the enzyme was inactivated completely with heat treatment at 60°C for 10 min. The optimum pH of LAP from strain 00-Bat-05 was 8.0 and high activity was still detectable at pH 10.0 and 11.0. However, the activity was relatively low under neutral conditions (pH 7.0). The HR-3 LAP had a broad pH range from 6.0 to 11.0 and the activity was still found

about 50% at pH 11.0 and 12.0. These results indicated that both enzymes have optimal activity at alkaline region similar to other aminopeptidases.^{16, 20)}

The LAP activities from strains 00-Bat-05 and HR-3 were inhibited by PCMB, suggesting both LAP enzymes to be SH enzymes. The LAP activity from 00-Bat-05 was inhibited by 1,10-phenanthroline and EDTA and similar results were found in LAP from *Geobacillus thermoleovorans*.²¹⁾ Most of LAP enzymes from bacteria and fungi were reported to be inhibited by metal-chelating agents and to be largely restored by divalent cations such as Zn²⁺ and Co²⁺.^{16, 20)} The LAP activity of strain 00-Bat-05 was inhibited by Zn²⁺ and stimulated by Co²⁺, while the enzyme activity of strain HR-3 was slightly inhibited by Co²⁺ and strongly inhibited by Zn²⁺. The enzyme of strain 00-Bat-05 was inhibited by K⁺, but the enzyme of strain HR-3 was distinctively activated by Na⁺ and K⁺. These results indicate that effect of metal ions on LAP activity varies from organism to organism.

As substrates, Leu-*p*-NA was best for the LAP enzyme of strain 00-Bat-05 LAP, followed by Met-*p*-NA (51 %), Ala-*p*-NA (21 %) and Arg-*p*-NA (20 %) and negligible activity was detected for Glu-*p*-NA and Pro-*p*-NA. On the other hand, the enzyme of strain HR-3 showed high activity for Leu-*p*-NA (100 %), Ala-*p*-NA (347 %), Met-*p*-NA (109 %) and Arg-*p*-NA (125 %) and slight activity for Glu-*p*-NA and Pro-*p*-NA.

This is the first report on the isolation and enzymological properties of LAP enzymes from labyrinthulid and thraustochytrid strains. The information obtained in this study will help to reveal the ecological roles in degradation process of plant materials and microbial cells in marine environments and physiological response of these organisms as live food of aquaculture animals.

Acknowledgments

The authors thank Dr. Masahiro Hayashi, Faculty of Agriculture, Miyazaki University, Japan for providing of thraustochytrid HR-3 strain and useful comments on the properties of this strain. This work was supported by a Core University program from the Japan Society for the Promotion of Science (JSPS).

References

- 1) Porter, D. (1989). Phylum Labyrinthulomycota. In "Handbook of Protoctista" (ed. by Margulis, L., J. O. Corliss, M. Melkonian, and D. J. Chapman) Jones and Bartlett Publishers, Boston. pp. 388-398.
- 2) Cavalier-Smith, T. (1993). Kingdom Protozoa and its 18 phyla. *Microbiol. Rev.*, **57**: 953-994.
- 3) Honda, D., T. Yokochi, T. Nakahara, S. Raghukumar, A. Nakagiri, K. Schaumann and T. Higashihara (1999). Molecular phylogeny of labyrinthulids and thraustochytrids based on the sequencing of 18S ribosomal RNA gene. *J. Eukaryot. Microbiol.*, **46**: 637-646.
- 4) Nakagiri A. (2001). Ecology of *Labyrinthula* and note on modified methods for cultivation and preservation of the isolates. *Aquabiol.*, **23**: 32-38 (in Japanese).
- 5) Raghukumar, S. (1992). Bacteriovory: a novel dual role for thraustochytrids in the sea. *Mar. Biol.*, **113**: 165-169.
- 6) Bremer, G. B. and G. Talbot (1995). Cellulolytic enzyme activity in the marine protist *Schizochytrium aggregatum*. *Bot. Mar.*, **38**: 37-41.
- 7) Raghukumar, S., V. Sathe-Pathak, S. Sharma and C. Raghukumar (1995). Thraustochytrid and fungal component of marine detritus. III. Field studies on decomposition of leaves of the mangrove *Rhizophora apiculata*. *Aquat. Microbiol. Ecol.*, **9**: 117-125.
- 8) Singh, A. and O. P. Ward (1997). Microbial production of docosahexaenoic acid (DHA, C22:6). *Adv. Appl. Microbiol.*, **45**: 271-312.
- 9) Nakahara, T., T. Yokochi, T. Higashihara, S. Tanaka, T. Yaguchi and D. Honda (1996). Production of docosahexaenoic and docosapentaenoic acids by *Schizochytrium* sp. isolated from Yap Islands. *J. Am. Oil Chem. Soc.*, **73**: 1421-1426.
- 10) Sakata, T., T. Fujisawa and T. Yoshikawa (2000). Colony formation and fatty acid composition of marine labyrinthulid isolates grown on agar media. *Fish. Sci.*, **66**: 84-90.
- 11) Miller, M. R., P. D. Nichols and C. G. Carter (2007). Replacement of fish oil with thraustochytrid *Schizochytrium* sp. L oil in Atlantic salmon parr (*Salmo salar* L) diets. *Com. Biochem. Physiol., Part A*, **148**: 382-392.
- 12) Hayashi, M., R. Matsumoto, T. Yoshimatsu, S. Tanaka and S. Shimizu (2002). Isolation of highly DHA-accumulated Labyrinthulales and their utilization for nutritional enrichment of rotifers and *Artemia*. *Nippon Suisan Gakkaishi*, **68**: 674-678 (in Japanese).
- 13) Gonzales, T. and J. Robert-Baudouy (1996). Bacterial amino peptidases: Properties and functions. *FEMS Microbiol. Rev.*, **18**: 319-344.
- 14) Caruso, G. and R. Zaccone (2000). Estimates of leucine aminopeptidase activity in different marine and brackish environments. *J. Appl. Microbiol.*, **89**: 951-959.
- 15) Himmelhoch, S. R. (1969). Leucine aminopeptidase: a zinc metalloenzyme. *Arch. Biochem. Biophys.*, **134**: 597-602.
- 16) Chien, H. R., L. Lin, S. Chao, C. Chen, W. Wang, C. Shaw, Y. Tsai, H. Hu and W. Hsu (2002). Purification, characterization, and genetic analysis of a leucine aminopeptidase from *Aspergillus sojae*. *Biochim. Biophys. Acta*, **1576**: 119-126.
- 17) Sakata, T. and K. Iwamoto (1995). Isolation of marine algicidal microorganisms on diatom double layer agar plates. *Fish. Sci.*, **61**: 173-174.
- 18) Saitou, H. and M. Nei (1963). The neighbor-joining: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.*, **4**: 406-425.
- 19) Wahid, M. I., T. Yoshikawa, and T. Sakata (2007). Bacteriolytic activity and growth of marine isolates of labyrinthulids on dead bacterial cells. *Fish. Sci.*, **73**: 1286-1294.
- 20) Lee, G., S. Chun, Y. Kho, and H. Chun (1998). Purification and properties of an extracellular leucine aminopeptidase from the *Bacillus* sp. N2. *J. Appl. Microbiol.*, **84**: 561-566.
- 21) Deejing, S., K. Yoshimune, S. Lumyong and M. Moriguchi (2005). Purification and characterization of hyperthermotolerant leucine aminopeptidase from *Geobacillus thermoleovorans* 47b. *J. Ind. Microbiol. Biotechnol.*, **32**: 269-276.