

Supplementary Note on the Previous Papers, "Observations of the Distribution of Giant Sea-Salt Particles in the Sea-Surface Boundary Layer" and "On the Variation of the Amount of Giant Sea-Salt Particles near the Ground"

Masaaki CHAEN*

In the author's two papers, "Observations of the distribution of giant sea-salt particles in the sea-surface boundary layer" and "On the variation of the amount of giant sea-salt particles near the ground", published in Memoirs of the Faculty of Fisheries, Kagoshima University, Vol. 20, No. 1, 1971, there was a confusion in the description of the dry mass of sea-salt particles, between a tentative classification and true classes. In the former paper, in the first paragraph of page 58, and in the latter paper in the first paragraph of section 3.1 on page 74, "sea-salt particles larger than 1.9×10^{-11} gm in mass of salt ($\log m = 0.5$, $m : 10^{-12}$ gm unit)" should be read "sea-salt particles larger than 1.9×10^{-11} gm in mass of salt ($\log m = 1.28$, $m : 10^{-12}$ gm unit)". Accordingly, $\log m$ values throughout the papers should be corrected as shown in Table 1.

Table 1. Corrigenda and supplement.

Figures in the papers	Correct figures	Dry mass of salt
$\log m = 0.50$	$\log m = 1.28$	$1.9 \times 10^{-11}(\text{gm})$
0.75	1.45	2.8 "
1.00	1.64	4.4 "
1.25	1.80	6.3 "
1.50	1.97	9.4 "
1.75	2.15	1.4×10^{-10}
2.00	2.34	2.2 "
2.25	2.53	3.4 "
2.50	2.72	5.2 "
2.75	2.91	8.1 "
3.00	3.10	1.3×10^{-9}

* Laboratory of Oceanography, Faculty of Fisheries, Kagoshima University.