



Doctoral Thesis

**AN ANALYSIS OF A US CASE-CONTROL
STUDY ON THE ASSOCIATION BETWEEN
HIGH BIRTH WEIGHT AND THE RISKS
OF CHILDHOOD CNS TUMORS AND
LEUKEMIA**

TRAN THANH LONG

Department of Epidemiology and Preventive Medicine

Kagoshima University Graduate School of Medical and Dental Sciences

Kagoshima, Japan

2013-2017

Doctoral Thesis

**AN ANALYSIS OF A US CASE-CONTROL
STUDY ON THE ASSOCIATION BETWEEN
HIGH BIRTH WEIGHT AND THE RISKS
OF CHILDHOOD CNS TUMORS AND
LEUKEMIA**

TRAN THANH LONG

Department of Epidemiology and Preventive Medicine

Kagoshima University Graduate School of Medical and Dental Sciences

Kagoshima, Japan

2013-2017

DECLARATION

I assure that the thesis entitled “**An analysis of a US case-control study on the association between high birth weight and the risks of childhood CNS tumors and leukemia**” submitted to Kagoshima University for the award of the degree of Doctor of Philosophy in Medical Science, is my original research work. In the thesis, all materials were collected and processed objectively and honestly. It has not been submitted previously to any other university.

TRAN THANH LONG

PUBLICATION LIST

Long TT, Hang TML, Koriyama C, Futoshi U, Akiba S. The association between high birth weight and the risks of childhood CNS tumors and leukemia: an analysis of a US case-control study in an epidemiological database. *BMC Cancer* 2017, 17(1):687.

Hang TML, Koriyama C, Tokudome S, Tran HH, **Long TT**, Athira N, Suminori A and Ngoan TL. Waterpipe tobacco smoking and gastric cancer risk among Vietnamese men. *PLOS ONE* 2016, 11(11): e0165587.

Dang TN, Seposo XT, Duc NH, Thang TB, An DD, Hang TML, **Long TT**, Loan BT, Honda Y. Characterizing the relationship between temperature and mortality in tropical and subtropical cities: a distributed lag non-linear model analysis in Hue, Viet Nam, 2009-2013. *Glob Health Action*. 2016 Jan 13;9:28738. doi: 10.3402/gha.v9.28738. eCollection 2016.

The part of this thesis is originally based on the first paper above published in BMC Cancer.

TABLE OF CONTENT

1	BACKGROUND	1
1.1	Epidemiology of childhood cancer	1
1.2	The association between birth weight and the risk of all childhood cancer	2
1.3	The association between birth weight and the risks of childhood CNS tumors and leukemia.....	3
1.3.1	The association between high birth weight and the risks of childhood CNS tumors and leukemia	3
1.3.2	The association between birth weight itself and the risks of childhood CNS tumors and leukemia	4
1.3.3	The association between gestational age and the risks of childhood CNS tumors and leukemia	5
1.3.4	The association between birth weight corrected for gestational age and the risks of childhood CNS tumors and leukemia	6
2	STUDY PURPOSE	8
3	METHODS	9
3.1	Overview of data from the CEDR database.....	9
3.2	Inclusion/exclusion criteria	11
3.3	Statistical analysis	11
3.4	Ethical clearance	13
4	RESULTS	14
4.1	Characteristics of CNS tumor and leukemia cases and controls	14
4.2	The associations between birth weight and the risks of CNS tumors and leukemia	15
5	DISCUSSION	22
6	CONCLUSION	29
7	FUTURE PROSPECTIVES	30
8	ACKNOWLEDGEMENT	31
9	REFERENCES	32
10	APPENDICES	37

LIST OF ABBREVIATIONS

AGA	Appropriate for gestational age
ALL	Acute lymphoblastic leukemia
AML	Acute myeloid leukemia
APC	Annual percent change
BW	Birth weight
CEDR	Comprehensive Epidemiologic Data Resource
CI	Confidence Interval
CNS	Central nervous system
DOE	Department of Energy
GA	Gestational age
IR	Incidence rate
LGA	Large for gestational age
OR	Odds ratio
SES	Socio-economic status
SGA	Small for gestational age

LIST OF TABLES

Table 1. Characteristics of cases and controls by factors matched (or surrogate factors) in the original study	14
Table 2. The association between birth weight and the risk of CNS tumors.....	17
Table 3. The association between birth weight and the risk of leukemia	18
Table 4. The association between gestational age and the risks of CNS tumors and leukemia	19
Table 5. CNS tumor risk among small-for-gestational-age and large-for-gestational-age children	19
Table 6. Leukemia risk among small-for-gestational-age and large-for-gestational-age children	20

LIST OF APPENDICES

Additional Table 1. The association between birth weight and CNS tumor risk without adjustment for gestational age	37
Additional Table 2. The association between birth weight and the CNS tumor risk among children with gestational age of 37-42 weeks.....	38
Additional Table 3. The association between birth weight and leukemia risk among children with gestational age of 37-42 weeks	39
Additional Table 4. The risk of CNS tumors or leukemia among small-for-gestational-age and large-for-gestational-age children with gestational age of 37-42 weeks	40

ABSTRACT

Background: The incidence of childhood cancer from 2001-2010 has increased since the 1980s in most parts of the world. The most common cancers among children are leukemia and central nervous system (CNS) tumors. According to a recent study conducted in the US, a significant upward trend in the incidence rate of acute lymphocytic leukemia was noticed in children aged 5 to 9 years between 2000 and 2010; however, the incidence rates of CNS tumors remained stable. High birth weight (BW), 4,000g or larger, is an established risk factor for childhood leukemia. However, its association with CNS tumor risk is yet unclear. The present study examined it, analyzing data obtained from a case-control study conducted among three states from the US. The association with childhood leukemia risk was also further examined.

Methods: In this study, a data set provided by the Comprehensive Epidemiologic Data Resource was analyzed with an official permission. The original case-control study was conducted to examine the association between paternal preconception exposure to ionizing radiation and childhood cancer risk. Cases with childhood cancer were mainly ascertained from local hospitals, and controls were selected, matched with birth year (1-year category), county of residence, sex, ethnicity and maternal age (+/-2 years). The controls in the original study consisted of children identified from birth certificates. Since the ID numbers were unavailable, conventional logistic analyses were conducted adjusting for those matching variables except for the county of residence. In addition to those variables, gestational age, age at diagnosis and study sites as covariables were included in the logistic models.

Results: Analyzed subjects were 72 CNS tumor cases, 124 leukemia cases and 822 controls born from 1945-1989. Cases and controls showed similar distributions regarding demographic factors and gestational age at birth, except birth year. There was no CNS tumor cases who were born before 1952. The proportion of children with CNS tumors born in later years, especially after 1970, was higher than that of children with leukemia. The odds ratios (ORs) of CNS tumor risk for children with low BWs (<2,500 g) and high BWs (>4,000 g) were 2.0 (95% confidence interval [CI] = 0.7, 5.9) and 2.5 (95% CI=1.2, 5.2)], respectively. When high-BW children were restricted to those who were large for gestational age (LGA), the OR for high-BW children remained similar (OR=2.7; 95%CI=1.1, 6.2). On the other hand, the ORs of leukemia risk for children with low and high BWs were 0.8 (95%CI=0.2, 3.0) and 1.4 (95%CI=0.7, 2.6), respectively. In the normal range of BW (2,500-4,000 g), higher BW was positively associated with CNS tumor risk (beta=0.0011, p for trend=0.012). However, the association with leukemia risk was not significant (beta= -0.0002, p for trend=0.475).

Conclusion: High-BW and LGA children had an elevated childhood CNS tumor risk. Low-BW was not associated with the CNS tumor risk. In the normal BW range, the BW itself was positively related to CNS tumor risk. No significant association between BW and childhood leukemia risk was observed in this study. Further investigations are required to explore the biologic mechanisms underlying the association between high BW and CNS tumor risk.

1 BACKGROUND

1.1 Epidemiology of childhood cancer

A recent study, reported by Steliarova-Foucher *et al.* [1], revealed that the incidence of childhood cancer from 2001-2010 has increased since the 1980s in all regions except sub-Saharan Africa, and the highest increase was observed in Southeast Asia. The most common cancers among children are leukemia and central nervous system (CNS) tumors. According to a recent study conducted in the US, a significant upward trend in the incidence rate of acute lymphocytic leukemia (ALL) was noticed in children aged 5 to 9 years between 2000 and 2010; however, the incidence rates of CNS tumors remained stable [2]. For children aged 10 to 14 years, however, the incidence rates of both ALL and CNS tumors increased significantly [2].

CNS tumor is the second most frequent solid tumor among children in the US between 2000 and 2010. The cancer incidence rate (IR) varies by site, and the highest IRs were observed in children with ALL and CNS tumors (3.84 and 3.57 per 100,000 population, respectively) [2]. For children aged 0 to 14 years old, boys experienced increases in ALL with annual percent change (APC) of 1.3, whereas malignant CNS tumors increased significantly in girls only (APC=0.8). The IR of ALL increased significantly in non-Hispanic white (APC=0.7) and black (APC=2.3) children, but remained stable in white-Hispanic children.

Malignant CNS tumors increased significantly in white-Hispanics (APC=0.8) and blacks (APC=1.7), and non-malignant CNS tumors tended to increase significantly in non-Hispanic white children (APC=2.0) [2].

The etiology of childhood cancer has been investigated for several decades but it is still largely unknown. Many environmental factors, including N-nitroso compounds, pesticides, tobacco smoke, electromagnetic frequencies, infectious agents, trauma, parental occupation, medications and vitamins, are suspected to be attributed to the development of childhood cancer. A few genetic syndromes and ionizing radiation are established risk factors for both childhood leukemia and CNS tumors [3, 4]. High birth weight (BW), 4,000 g or larger, is also known to be a risk factor for childhood leukemia, especially ALL [5-8]. However, its association with childhood CNS tumor risk is yet unclear [5, 6, 9, 10].

1.2 The association between birth weight and the risk of all childhood cancer

Among several intrinsic characteristics of children or their parents, BW has been studied whether it is associated with childhood cancer risk or not. One of the most consistent findings is that high BW, 4,000 g or larger, increased the risk of leukemia risk, both ALL and acute myeloid leukemia (AML) [7, 8].

High BW is also suspected to be risk factor of CNS tumors, lymphomas and Wilms tumors [6, 9, 11, 12]. An international population-based case-control

studies revealed that total cancer risk increased linearly with each 500 g increase in BW in both the UK (odds ratio [OR]=1.06; 95% confidence interval [CI]=1.04, 1.10) and the US (OR=1.10; 95%CI=1.04, 1.08). For approximately half of all childhood cancers, BW was positively associated with the cancer risk [5]. In a population-based case-control study conducted in four Nordic countries, compared to children with 3,000-3,500 g, those with BWs of 4,500 g or larger had ORs of 1.2 (95%CI=0.9, 1.6), 1.8 (95%CI=1.3, 2.6), 1.4 (95%CI=1.3, 1.5) for lymphomas, Wilms tumors, and overall childhood cancer, respectively [12].

1.3 The association between birth weight and the risks of childhood CNS tumors and leukemia

1.3.1 The association between high birth weight and the risks of childhood CNS tumors and leukemia

In a large case-control study of children younger than 5 years of age, conducted in Texas, the US, the leukemia risk was elevated among those with high BWs (OR)=1.36; 95%CI=1.10, 1.69). However, the CNS tumor risk was not evidently increased among them (OR=1.14; 95%CI=0.83, 1.56) [6]. Similar results were obtained by a German study. High-BW children had ORs of 1.41 (95%CI=1.08, 1.84), 1.56 (95%CI=0.88, 2.79) and 1.34 (95%CI=0.97, 1.85), respectively, for ALL, AML and CNS tumors when compared to normal-BW children [11]. However, it should be noted that gestational age (GA) was not adjusted in those studies. Another large case-control study, conducted in

California, which focused on CNS tumors reported a GA-adjusted OR of 1.12 (95%CI=0.91, 1.38) [9]. In a population-based case-control study conducted in four Nordic countries, the ORs of ALL, AML and CNS tumors were 1.3 (95%CI=1.1, 1.5), 1.5 (95%CI=1.3, 1.8) and 1.3 (95%CI=0.85, 2.0), respectively, when children with BWs of 4,500 g or larger were compared to those with 3,000-3,500 g, adjusting for GA [12]. Taken together, the leukemia risk was increased by 30% to 50% even after the adjustment for GA. In the case of CNS tumor risk, the association appears to be weaker.

1.3.2 The association between birth weight itself and the risks of childhood CNS tumors and leukemia

Regarding the effect of BW itself, several studies have investigated its effect on the risks of childhood leukemia and CNS tumors. Previous studies in Texas [6] and California [13] consistently found that an each 1,000-g increase in BW was associated with leukemia risk: the ORs (95%CI) were 1.28 (1.12, 1.44) and 1.11 (1.01, 1.21), respectively. A pooled analysis using the data obtained from nationwide studies in the US (16,554 cases) and UK (23,772 cases) found similar results: ORs per 500-g increase were 1.10 (95%CI=1.06, 1.13) and 1.07 per 500 g (95%CI=1.04, 1.10), respectively [5]. Higher BW is also suspected to be related to an increased CNS tumor risk. The studies conducted in Texas and California reported ORs per 1,000-g increase of 1.17 (95%CI=0.98, 1.40) [6] and 1.11 (95%CI=0.99, 1.24) [9], respectively. Similar results were obtained by the

pooled study above. ORs per 500-g increase were 1.05 (95%CI=1.01, 1.08) and 1.07 (95%CI=1.04, 1.10) in the US and UK studies, respectively [5]. In the UK study, however, GA was not taken into account.

1.3.3 The association between gestational age and the risks of childhood CNS tumors and leukemia

Longer GAs are also suspected to be a risk factor for CNS tumors. A French study [14] reported that children with longer GAs (41 weeks or longer) were at an increased CNS tumor risk (OR=1.4; 95%CI=0.6, 3.3) when compared to those with the GA of 37-40 weeks, although there was no statistical significance. A Swedish study observed a similar trend in which children with the GA of 43 weeks or longer had a 1.2-fold increase of brain tumor risk (OR=1.2; 95%CI=0.4, 3.8) when compared to those with the GA of 38-42 weeks [15]. Only a slight increase in the CNS tumor risk was observed in the Texas study (OR=1.07; 95%CI=0.78, 1.47) [6]. However, the findings on the association of leukemia risk with GA were inconsistent. The Texas study reported that children with the GA of 41 weeks or longer had a slightly decreased leukemia risk (OR=0.91; 95%CI=0.71, 1.15) when compared to those with the GA of 37-40 weeks [6]. A contrary result was reported in a study conducted in Denmark, Sweden, Norway and Iceland, which pointed to an OR of 1.08 (95%CI=0.90, 1.29) for longer GAs (42 weeks or longer) compared to the GA of 40-41 weeks [16].

1.3.4 The association between birth weight corrected for gestational age and the risks of childhood CNS tumors and leukemia

BW is strongly related to GA [17]. Based on GA, BW can be divided into three categories: small for gestational age (SGA), appropriate for gestational age (AGA) and large for gestational age (LGA). In the Texas study, the LGA was significantly associated with an increased ALL risk (OR=1.66; 95%CI=1.32, 2.10), but not for CNS tumor risk (OR=1.14;95%CI=0.82, 1.58) [6]. A study in California also showed no significant association between LGA and the risk of CNS tumors (OR=1.09; 95%CI=0.89, 1.27) [9]. In the German study, the OR of ALL was 1.45 (95%CI=1.07, 1.97) in LGA children compared to AGA children. However, the OR for CNS tumor was not statistically significant: 1.18 (95%CI=0.80, 1.72) [11]. In the Nordic study, LGA was related neither ALL risk (OR=1.2; 95%CI=0.91, 1.5) nor CNS tumor risk (OR=1.1; 95%CI=0.85, 1.4) [12]. Taken together, those studies suggested that LGA children may be at an elevated ALL risk. The association with the risk of CNS tumors is unlikely.

The studies described above showed no association between CNS tumor risk and SGA. The ORs in the studies of California [9], Texas [6], West Germany [11], and the Nordic countries [12] were 0.96 (95%CI=0.75, 1.23), 0.98 (95%CI=0.70, 1.38), 0.96 (95%CI=0.67, 1.37) and 0.95 (95%CI=0.77, 1.20), respectively. However, the findings on the association between leukemia risk and SGA are inconsistent. The ORs for all types of leukemia and ALL were 0.88

(95%CI=0.68, 1.13) and 0.78 (95%CI=0.57, 1.05), respectively, in the Texas study [6]. The ORs for ALL and AML were 1.00 (95%CI=0.74, 1.35) and 0.89 (95%CI=0.43, 1.83), respectively, in the German study [11], and 1.2 (95%CI=0.96, 1.50) and 1.8 (95%CI=1.1, 3.1), respectively, in the Nordic study [12].

2 STUDY PURPOSE

We analyzed data from a case-control study which was originally conducted in the US to examine the association between paternal preconception exposure to ionizing radiation and the risk of childhood cancer, and this study found no association between them [18]. Using this dataset, we examined the association between BW and childhood cancer risk.

3 METHODS

3.1 Overview of data from the CEDR database

We used data from a case-control study of childhood cancers and paternal preconception occupational exposure to ionizing radiation in counties surrounding three US Department of Energy (DOE) nuclear facilities. The data, which were obtained by the study conducted by Sever *et al* [18], are available in the Comprehensive Epidemiologic Data Resource (CEDR) database through CEDR website [19] after getting an official permission. The three facilities were the Hanford (Hanford), Idaho National Engineering Laboratory (INEL) and Oak Ridge (K-25, Y-12, and X-10 at Oak Ridge laboratories). The counties selected for the study in each of 3 DOE nuclear facilities were as follows: the Benton and Franklin counties in Hanford; the Bannock, Bingham, Bonneville, Buttee, Jefferson and Madison counties in INEL; and the Anderson, Knox and Roane counties in Oak Ridge. Those counties were selected, as most of the workers of the corresponding DOEs at those sites resided in them [18].

This study included 75 CNS tumor cases, 132 leukemia cases and 26 non-Hodgkin's lymphoma cases, which were diagnosed prior to the age of 15 years, from 1957-1991. According to the original report [18], cases had to be born to residents of one of the study counties and be residents of one of them when their cancer was diagnosed. Cases were ascertained from each of the populations, using multiple sources (local primary care hospitals, regional referral hospitals, cancer

registries and death certificates), as population-based cancer registries were unavailable in those areas during the period of 1957-1991. The controls analyzed in the present study (N=1,047) were matched based on year of birth (1-year category), county of residence, sex, ethnicity and maternal age (+/-2 years). The controls in the original study consisted of children identified from birth certificates. In the case of Hanford, the birth certificate controls were selected from a computer file provided by the Technical and Data Services Section, Center Health Statistics, Washington State Department of Health [18]. Server *et al.* identified all the births that matched each case on the basis of the year of birth, race, sex and maternal age. A file of potential controls was developed; this included all the births matching each case.

For all the cases, information on diagnosis and cause of death was abstracted from hospital records, tumor registries and death certificates in the original study. Sever *et al.* [18] stated in their report that "each source was utilized to provide as complete an ascertainment as possible". Pathological reports were reviewed to obtain the most accurate histopathological data.

Demographic information including sex, ethnicity, year of birth and address at the time of the diagnosis was abstracted from birth certificates or electronic birth files. Information on parental employment was collected from records at the DOE sites. Information on pregnancy (parity, date of the mother's last menstrual period, initiation of prenatal care, viral infections during pregnancy

and X-ray during pregnancy), delivery (breach or other malpresentation and clinical estimation of GA), and newborn characteristics (plurality, BW and congenital malformation) was obtained from medical records [18].

3.2 Inclusion/exclusion criteria

In our study, we excluded children in whom information on BW, GA and year of diagnosis was lacking. Those whose ethnicities were categorized as others or unknown were also excluded. Non-Hodgkin's lymphoma cases were not used because the number of cases was few for statistical analysis. After excluding ineligible subjects, the number of eligible subjects for CNS tumor cases, leukemia cases and controls used in statistical analysis were 72, 124 and 822, respectively.

3.3 Statistical analysis

We analyzed the association between BW and the risks of CNS tumors and leukemia, using a conventional logistic model [20]. All p values were two-sided and calculated, using the likelihood ratio test. The p values for trend were calculated, using continuous variables. Data analyses were performed, using Software Stata 14.0.

In the original study, the cases and controls were matched according to the year of birth (1 year category), county of residence, sex, ethnicity (black or white), and maternal age (+/-2 years). However, information on the county of residence is unavailable in the data, which we downloaded from the CEDR database.

Therefore, we generated a new variable on DOE sites as surrogate variable based on birth places of the study subject. In the CEDR database, the birth places were divided into the following eight categories: Hanford hospitals, Idaho hospitals, Tennessee hospitals, home, birth center, maternity hospitals and unknown. Those who were born at home, or in birth centers, maternity hospitals and unknown were coded as a missing value in the variable on DOE sites (23 and 8 subjects in the original study and present study, respectively).

In the available data set, the ID number to identify the matched control(s) for each case was unavailable; therefore, we could not conduct conditional logistic models. Therefore, we conducted conventional logistic analysis. When the analysis of matched case-control data ignores case-control matching, all the matched factors should be treated as potential confounders in statistical analysis [21]. Therefore, we adjusted for the matching variables (birth year, county of residence, sex, ethnicity and maternal age). In addition, we also included GA, DOE sites and age at diagnosis as independent variables in the logistic model as well. Age at diagnosis for controls was calculated, using the year of diagnosis, which was assigned to the controls by the original study (the year of diagnosis of each case was assigned to the corresponding controls by the original study). The DOE sites were used as a surrogate variable for the county of residence.

Low BW is defined, by the World Health Organization, as a BW smaller than 2500 g. High BW is defined by Centers for Disease Control and Prevention

as a BW larger than 4,000 g [22]. Furthermore, we used BW corrected for GA to categorize the subjects as being LGA, AGA and SGA. In the present study, LGA children were those with BWs greater than the 90th percentile for their GAs. Children whose BW was below the 10th percentile for their GAs were classified as SGA. AGA children were those whose BWs were in the 10-90 percentile for their GAs. Those categories were constructed, using the US national reference for fetal growth [23].

3.4 Ethical clearance

Ethical Guidelines for Medical and Health Research Involving Human Subjects, Japan, is not applicable to this study because the data base used in the present study is anonymous and open-access.

4 RESULTS

4.1 Characteristics of CNS tumor and leukemia cases and controls

Table 1. Characteristics of cases and controls by factors matched (or surrogate factors) in the original study

Variables	Controls (N=822)	Cases	
		CNS tumor (N=72)	Leukemia (N=124)
Year of birth	1946-1989	1952-1989	1949-1989
1946-1959	120 (14.6%)	11 (15.3%)	22 (17.7%)
1960-1969	240 (29.2%)	16 (22.2%)	44 (35.5%)
1970-1979	294 (35.8%)	28 (38.9%)	35 (28.2%)
1980-1989	168 (20.4%)	17 (23.6%)	23 (18.6%)
Age at diagnosis (years)^a			
Mean (SD)	6.1 (4.4)	5.6 (4.3)	5.3 (4.1)
Min-Max	0-15	0-14	0-14
Sex			
Male	493 (60.0%)	45 (62.5%)	71 (57.3%)
Female	329 (40.0%)	27 (37.5%)	53 (42.7%)
Ethnicity			
Black	18 (2.2%)	3 (4.2%)	2 (1.6%)
White	804 (97.8%)	69 (95.8%)	122 (98.4%)
Maternal age (years)			
Mean (SD)	25.5 (5.4)	25.2 (5.2)	25.5 (5.8)
Min-Max	14-44	15-37	16-42
DOE sites^b			
Hanford	271 (32.9%)	19 (26.4%)	28 (22.6%)
INEL	183 (22.3%)	15 (20.8%)	33 (26.6%)
Oak Ridge	363 (44.2%)	37 (51.4%)	61 (49.2%)
Unknown	5 (0.6%)	1 (1.4%)	2 (1.6%)
Gestational age (weeks)^c			
Mean (SD)	39.3 (1.9)	38.5 (2.4)	39.4 (1.8)
Min-Max	28-45	27-43	34-44

SD: Standard Deviation, DOE: Department of Energy

^aAge at diagnosis for controls was calculated, using the year of diagnosis assigned by the original study, which was matched case-control study

^bDOE sites: a surrogate variable for county of residence

^cNot matched in the original study

The characteristics of the CNS tumor and leukemia cases and the controls, according to the factors matched (or surrogate factors) in the original study, are presented in table 1. Cases and controls showed similar distributions regarding those factors. One exception was the year of birth. CNS tumor cases did not have those born before 1952. The proportion of children with CNS tumors born in later years, especially after 1970, was higher compared to that of children with leukemia. In this table, DOE sites are a surrogate factor for the county of residence, which was matched in the original study, but was unavailable in the database. Regarding the DOE sites' distribution, the control group had more subjects in Hanford and less in Oak Ridge. In order to control those potential confounders, we included those variables in the conventional logistic models in the risk analysis.

4.2 The associations between birth weight and the risks of CNS tumors and leukemia

In the following tables, the results of the logistic analysis are summarized. The analysis for leukemia risk was also conducted and their results are included in those tables for comparison. As shown in table 2, CNS tumor risk increased with BW (p value for trend =0.010). When those with BW less than 2,500 g were excluded, the association became stronger (p for trend <0.001). Even among those in the normal-BW range (2,500-4,000 g), the p for trend was significant (p=0.012). The increasing trend was mainly from those larger than 4,000 g. The

OR for this high BW adjusted for GA was 2.5 (95%CI=1.2, 5.2) when compared to normal BW (2,500-4,000 g). The GA-unadjusted OR was 2.0 (95%CI=1.0, 4.1) (additional table 1). In this table, we also made a comparison between low-BW and normal-BW children. The CNS tumor risk was also increased among low-BW children, and the OR was 2.0 (95%CI=0.7, 5.9); however, the increase was not statistically significant (p=0.241).

Among the high-BW children, SGA, AGA and LGA accounted for 2, 33 and 51 children, respectively. When high-BW children were restricted to LGA, the OR for CNS tumors was 2.7 (95%CI=1.1, 6.2; p=0.035) as shown in the middle panel of table 2. When high-BW children were restricted to SGA/AGA (the lower panel of table 2), the OR for CNS tumors became smaller (OR=2.2; 95%CI=0.7, 6.7; p=0.209).

Table 2. The association between birth weight and the risk of CNS tumors

Birth weight	Controls	CNS tumors	OR	95%CI		P value
				Lower	Upper	
Total subjects						
<2,500 g	24	7	1.8	0.5	5.8	0.363
2,500-<3,000 g	137	7	0.6	0.2	1.4	0.214
3,000-<3,500 g	305	21	1	Reference		
3,500-4,000 g	276	25	1.5	0.8	2.8	0.205
>4,000 g	80	12	2.9	1.3	6.6	0.012
<i>P for homogeneity=0.017</i>						
<i>For all: P for trend=0.010 (beta=0.0007)</i>						
<i>For birth weight $\geq 2,500$ g: P for trend <0.001 (beta=0.0011)</i>						
<i>For birth weight 2,500-4,000 g: P for trend =0.012(beta=0.0011)</i>						
<2,500 g	24	7	2.0	0.7	5.9	0.241
2,500-4,000 g	718	53	1	Reference		
>4,000 g	80	12	2.5	1.2	5.2	0.018
<i>P for homogeneity=0.028</i>						
The risk of high-birth-weight and LGA children compared to normal-birth-weight children*						
2,500-4,000 g	718	53	1	Reference		
>4,000 g and LGA	48	8	2.7	1.1	6.2	0.035
The risk of high-birth-weight and SGA/AGA children compared to normal-birth-weight children*						
2,500-4,000 g	718	53	1	Reference		
>4,000 g and SGA/AGA	32	4	2.2	0.7	6.7	0.209

LGA: large for gestational age, SGA: small for gestational age, AGA: appropriate for gestational age ORs and corresponding 95% CIs and p values were adjusted for sex, ethnicity, year of birth, age at diagnosis, gestational age (continuous variable), maternal age and DOE sites.

* Children with low-birth weight were not included in the analyses.

Leukemia risk was not associated with BW (table 3). In the lower panel of table 3, among high-BW children, the risk was increased by 40%, but the increase was not statistically significant.

Table 3. The association between birth weight and the risk of leukemia

Birth weight	Controls	Leukemia cases	OR	95%CI		P value
				Lower	Upper	
Total subjects						
<2,500 g	24	3	0.7	0.2	2.6	0.564
2,500-<3,000 g	137	20	0.7	0.4	1.3	0.300
3,000-<3,500 g	305	54	1	Reference		
3,500-4,000 g	276	33	0.7	0.4	1.1	0.092
>4,000 g	80	14	1.1	0.6	2.2	0.752
<i>P for homogeneity=0.396</i>						
<i>For all: P for trend=0.778 (beta=0.00006)</i>						
<i>For birth weight $\geq 2,500$ g: P for trend=0.833 (beta=0.00005)</i>						
<i>For birth weight 2,500-4,000 g: P for trend=0.475 (beta=-0.00022)</i>						
<2,500 g	24	3	0.8	0.2	3.0	0.765
2,500-4,000 g	718	107	1	Reference		
>4,000 g	80	14	1.4	0.7	2.6	0.343
<i>P for homogeneity=0.611</i>						
The risk of high-birth-weight and LGA children compared to normal-birth-weight children*						
2,500-4,000 g	718	107	1	Reference		
>4,000 g and LGA	48	10	1.7	0.8	3.7	0.166
The risk of high-birth-weight and SGA/AGA children compared to normal-birth-weight children*						
2,500-4,000 g	718	107	1	Reference		
>4,000 g and SGA/AGA	32	4	0.9	0.3	2.7	0.865

LGA: large for gestational age, SGA: small for gestational age, AGA: appropriate for gestational age
 ORs and corresponding 95% CIs and p values were adjusted for sex, ethnicity, year of birth, age at diagnosis, gestational age (continuous variable), maternal age and DOE sites.

* Children with low-birth weight were not included in the analyses.

We examined the association of GA with the risks of CNS tumor and leukemia (table 4). The CNS tumor risk was positively associated with longer GA (42 weeks or longer) after adjustment for BW (p for trend=0.001). On the other hand, the leukemia risk was elevated among children with longer GA.

Table 4. The association between gestational age and the risks of CNS tumors and leukemia

Gestational age	Controls	Cases	OR	95%CI		P value
				Lower	Upper	
For the analysis of CNS tumor risk						
<37 weeks	54	9	5.1	1.7	14.9	0.005
37-39 weeks	331	38	3.3	1.8	6.3	<0.001
40-41 weeks	366	20	1	Reference		
>41 weeks	71	5	1.3	0.5	3.6	0.644
<i>P for trend=0.001</i>						
For the analysis of Leukemia risk						
<37 weeks	54	6	1.2	0.4	3.5	0.682
37-39 weeks	331	51	1.4	0.9	2.2	0.175
40-41 weeks	366	52	1	Reference		
>41 weeks	71	15	1.6	0.8	3.0	0.162
<i>P for trend=0.930</i>						

ORs and corresponding 95% CIs and p values were adjusted for sex, ethnicity, year of birth, age at diagnosis, maternal age, birth weight (5-category variable) and DOE sites.

Table 5. CNS tumor risk among small-for-gestational-age and large-for-gestational-age children

Birth weight	Controls	CNS tumors	OR	95%CI		P value
				Lower	Upper	
Total subjects						
SGA	189	15	0.9	0.4	1.7	0.643
AGA	566	48	1	Reference		
LGA	63	9	1.8	0.8	3.9	0.163
<i>P for homogeneity=0.307</i>						
Birth weight 2,500 g or larger						
SGA	177	12	0.8	0.4	1.6	0.494
AGA	544	44	1	Reference		
LGA	63	9	2.0	0.9	4.5	0.101
<i>P for homogeneity=0.173</i>						
Birth weight 3,000 g or larger						
SGA	97	9	1.2	0.5	3.1	0.672
AGA	497	40	1	Reference		
LGA	63	9	2.0	0.9	4.4	0.113
<i>P for homogeneity=0.279</i>						

SGA: Small for gestational age, AGA: Appropriate for gestational age, LGA: Large for gestational age

ORs and corresponding 95% CIs and p values were adjusted for sex, ethnicity, year of birth, age at diagnosis, maternal age and DOE sites.

We examined the association of LGA and SGA with the risks of CNS tumors and leukemia (tables 5 and 6). LGA children were at higher risks of CNS tumors and leukemia, but neither increase was statistically significant. Even when the subjects were limited to those with BWs 2,500 g or larger, or those with BWs 3,000 g or larger, the results did not change sizably. The risk of CNS tumors or leukemia was not statistically significantly associated with SGA.

Table 6. Leukemia risk among small-for-gestational-age and large-for-gestational-age children

Birth weight	Controls	Leukemia cases	OR	95%CI		P value
				Lower	Upper	
Total subjects						
SGA	189	30	0.9	0.6	1.5	0.696
AGA	566	83	1	Reference		
LGA	63	11	1.4	0.7	2.9	0.342
						<i>P for homogeneity=0.555</i>
Birth weight 2,500 g or larger						
SGA	177	29	0.9	0.5	1.5	0.714
AGA	554	81	1	Reference		
LGA	63	11	1.4	0.7	2.9	0.340
						<i>P for homogeneity=0.561</i>
Birth weight 3,000 g or larger						
SGA	97	18	0.9	0.5	1.8	0.841
AGA	497	72	1	Reference		
LGA	63	11	1.5	0.7	3.1	0.298
						<i>P for homogeneity=0.547</i>

SGA: Small for gestational age, AGA: Appropriate for gestational age, LGA: Large for gestational age

ORs and corresponding 95% CIs and p values were adjusted for sex, ethnicity, year of birth, age at diagnosis, maternal age and DOE sites.

The American Congress of Obstetricians and Gynecologists has redefined "term pregnancy" and replaced it with four new definitions of "term" deliveries:

early term (37 weeks 0 day - 38 weeks 6 days), full term (39 weeks 0 day - 40 weeks 6 days), late term (41 weeks 0 day - 41 weeks 6 days) and postterm (42 weeks 0 day and beyond). We relaxed the definition for normal GA to avoid losing the number of cases, and used children with GA of 37-42 weeks. This decision increased the number of CNS tumor and leukemia cases, and the controls by 5, 11 and 51, respectively. However, the associations of BW or LGA/SGA with the risk of CNS tumors or leukemia did not change appreciably (additional tables 2, 3 and 4).

5 DISCUSSION

The present study showed that higher BW was positively associated with childhood CNS tumor risk with or without adjustment for GA. This observed association was mainly from those larger than 4,000 g. The OR among the high-BW children was 2.5 (95%CI=1.2, 5.2) with adjustment for GA, and 2.0 (95%CI=1.0, 4.1) without adjustment. Those values are higher than those reported by the previously conducted studies [6, 9, 11, 12].

Leukemia risk was increased (OR=1.4; 95%CI=0.7, 2.6; p=0.343) among the high-BW children. A meta-analysis reported a similar OR (OR=1.35; 95%CI=1.24, 1.48) on the basis of 32 studies [7]. The fact that this study was unable to establish a significant association between high BW and leukemia risk could be attributed to the fact that the effect estimate of high BW might be too small, relative to the sample size.

Even in the normal-BW range (2,500-4,000 g), higher BW was still positively associated with childhood CNS tumor risk (p for trend=0.012), but not with leukemia risk (p for trend=0.475). To date, no study has found that BW is related to childhood CNS tumor risk in the normal-BW range.

Several studies examined the association of BW itself with the risk of childhood CNS tumors and leukemia [5, 6, 9, 13]. Based on their findings as already mentioned in section 1.3.2, we conducted a meta-analysis on BW and the

risks of CNS tumor and leukemia. Pooled ORs per 1,000g increase were 1.12 (95%CI=1.08, 1.17) and 1.17 (95%CI=1.13, 1.21) for CNS tumors and leukemia, respectively. In the previous studies, the association between BW and CNS tumor risk was weak and its magnitude was smaller than that for leukemia. On the other hand, the present study showed a larger magnitude of CNS tumor risk by 1,000g increase (OR=2.07, 95%CI=1.19, 3.67) comparing to that of leukemia (OR=1.07, 95%CI=0.69, 1.65).

In the present study, GA was inversely associated with CNS tumor risk (p for trend=0.001). This finding is at variance with those obtained from the other studies, which reported a weak positive association between BW and CNS tumor risk [6, 14, 15]. The association between leukemia risk and GA was not found in our study (p for trend=0.930) as was the case with the other studies [6, 16].

BW and GA are known to be closely related to each other [17]. When the high-BW children were restricted to those who were LGA, the OR was 2.7 (95%CI=1.1, 6.2). When high-BW children were restricted to those without LGA, the OR was 2.2 (95%CI=0.7, 6.7), which is smaller than the OR for high-BW and LGA children. In the present study, SGA was not statistically related to the risk of CNS tumors or leukemia.

Our study found an increased risk of CNS tumors among LGA children, but the increase was not statistically significant. The OR obtained in our study (OR=1.8; 95%CI=0.8, 3.9), which was larger than those reported by the other

studies (in which the ORs were in the range of 1.09-1.18) [6, 9, 11, 12]. In the case of leukemia, our study obtained an OR of 1.4 (95% CI=0.7, 2.9), which is similar to those reported by other studies (in which the ORs were in the range of 1.45-1.66) [6, 11].

In the present study, CNS tumor risk was not associated with SGA (OR=0.9; 95% CI=0.4, 1.7) as was the case with the other studies [6, 9, 11, 12]. The OR for leukemia was 0.9 (95% CI=0.6, 1.5). The association between leukemia risk and SGA on the literature is inconsistent. The OR obtained from the US and German studies were in the range of 0.78 to 1.00 [6, 11], and were 1.2 to 1.8 in Nordic study [12]. Our result is similar to the values reported by the Texan and German studies [6, 11].

CNS tumors have various histological types which may have different etiological backgrounds. The three most common types of childhood CNS tumors include medulloblastomas, astrocytomas and malignant gliomas, which accounted for 50% of those tumors in a US study [24]. A meta-analysis of eight studies reported in 2008 showed that high-BW children had slightly elevated risks of astrocytoma (OR=1.38, 95% CI=1.07, 1.79) and medulloblastoma (OR=1.27, 95% CI=1.02, 1.60) [10]. Among the eight studies, only California study considered the GA as a potential confounder [15, 25-31]. In the present study, we did not have information on the pathological types of the tumors.

Several mechanisms which stimulate prenatal weight gain and act simultaneously as long-term carcinogens might explain the association between high BW and the increased risk of CNS tumors. First, high BW could be an indicator of a greater number of cells, leading to more cell divisions. It is strongly suspected that such a condition could make them more vulnerable to carcinogenic agents and therefore, the cancer risk increases after birth [32]. BW is known to be positively correlated with insulin-like growth factor-1, which is strongly suggested to be involved in brain ontogenesis and carcinogenesis [33, 34]. Second, Heuch *et al.* [27] proposed the involvement of excess prenatal nutrition in medulloblastoma development, and suspected that high BW is an important indicator of excess nutrition in the last gestational trimester. They suspected that ample nutrition may interfere with the migration of granular neuronal cells, which starts at approximately 30 gestational weeks. If the cells migrate incompletely, they may remain immature. As a result, neoplastic potential of the cell may increase.

In the present study, childhood cancer patients were diagnosed from 1957 to 1991. As shown in table 1, the proportion of CNS tumor patients seems to have increased with calendar year, though this upward trend was not observed in the case of childhood leukemia. The improvement in diagnostic technologies could have led to artifactual increases in the rate CNS tumor occurrence [35]. It is to be noted that computed tomography and magnetic resonance imaging scans were

widely used in the 1970s and 1980s, respectively. The number of cases was small in the first 15 years (1946-1959) probably because the cases in original study might be obtained from the limited sources. The number of subjects in the period of 1980-1989 reduced due to the small number of children reached 15 years old.

Our study has several limitations. First, the results should be treated with considerable caution because of the limited number of cases. Regarding the leukemia risk, we failed to find a significant association. The effect estimate of high BW might be too small compared to the sample size. Second, cases were ascertained mainly from hospitals. Although the original study described “cancer registry” as a source of case ascertainment, we assumed that this might have been a hospital-based registry, as population-based cancer registries were unavailable in the 1957-1991 period. Thus, we could deny the possibility that cases without consultation at the hospitals or diagnosed outside of the study areas could be missed. Third, we lacked information on the subtypes of CNS tumors and leukemia. Typically, tumor registries did not cover those years. Death certificates did not provide identification of a hospital where diagnostic information might be located. The data in hospital records were insufficient for those years. Fourth, the study encountered problems in obtaining the birth records of the cases and controls. While Sever *et al.* received high level of cooperation from many hospitals that provided them with access to records, the medical records themselves were often missing and the data were incomplete [18]. Since these

problems were mainly with newborn records, that they did not affect the cases and controls differently. Fifth, the study did not collect sufficient information on the socio-economic status (SES) of the subjects. Unlike in the case of the relationship between SES and low BW, the association between SES and high BW risk is not consistent [36]. Many studies have been conducted to examine the association between SES and leukemia risk. On reviewing studies published until 1982, higher SES was suspected to be related to childhood leukemia risk [37]. A review by Poole *et al.* [38], however, noted that most later studies consistently reported inverse associations of childhood leukemia with SES; it was concluded, therefore, that associations between SES measures and childhood leukemia likely vary with the time and place. A study based on 5,240 leukemia cases from the Canadian cancer registries, that covered at least 95% of all the cases, reported a slightly lower relative risk of leukemia in the poorest group (RR=0.87; 95%CI=0.80, 0.95) [39]. A similar finding was also reported in a large case-control study from the UK (OR=0.99, 95%CI=0.96, 1.01) [40]. Thus, the effect of SES on the association between BW and leukemia risk may be considerably small even if SES is a potential confounding factor. The association between SES and CNS tumor risk was still inclusive [41-44]. Sixth, information on maternal comorbidities was not available in this data set. Although gestational diabetes mellitus is the most important risk factor for high BW and LGA, we could not examine the effect of gestational diabetes mellitus on childhood cancer risk.

Finally, SGA was not a risk factor for childhood cancers in our study. The Barker hypothesis shows that low BW is associated to the risk of developing chronic diseases in later life [45-47]. However, the association of low BW and childhood cancer risk has not been clarified.

6 CONCLUSION

High-BW and LGA children had an elevated childhood CNS tumor risk. In the normal BW range, BW itself was positively related to CNS tumor risk. Low BW was not associated with an increased CNS tumor risk. No significant association between BW and childhood leukemia risk was observed in this study.

7 FUTURE PROSPECTIVES

Further investigations are recommended to explore the biologic mechanisms underlying the association between high birth weight and CNS tumor risk.

8 ACKNOWLEDGEMENT

With a profound sense of gratitude, I acknowledge that this work was completed under the kind support and supervision of Dr. Suminori Akiba, Former Chairman Department of Epidemiology and Preventive Medicine.

I would like to express my sincere thanks and gratitude to co-supervisor Dr. Chihaya Koriyama, Current Chairman Department of Epidemiology and Preventive Medicine, for her keen interest and skilled guidance at every stage of my research.

Many thanks go to my friends in Kagoshima and my friends at the Department of Epidemiology and Preventive Medicine for constant moral encouragement and support when I was in difficult situations.

I am extremely thankful to my wife, Mrs.Lai Thi Minh Hang and all my loving family in Vietnam for their supports, understanding and encouragement.

Especially, I would like to thank the Japanese Ministry of Education, Culture, Sports, Science and Technology (Monbukagakusho) for the financial support.

Finally, on behalf of the authors for the paper, which I used for my thesis, I would like to express our sincere thanks for sharing the data from Comprehensive Epidemiologic Data Resource (CEDR) database by The U.S Department of Energy.

9 REFERENCES

1. Steliarova-Foucher E, Colombet M, Ries LAG, Moreno F, Dolya A, Bray F, Hesselning P, Shin HY, Stiller CA, contributors I-. International incidence of childhood cancer, 2001-10: a population-based registry study. *The Lancet Oncology* (2017);18 (6):719-731. doi:10.1016/S1470-2045(17)30186-9
2. Gittleman HR, Ostrom QT, Rouse CD, Dowling JA, de Blank PM, Kruchko CA, Elder B, Rosenfeld SS, Selman WR, Sloan AE, Barnholtz-Sloan JS. Trends in central nervous system tumor incidence relative to other common cancers in adults, adolescents, and children in the United States, 2000 to 2010. *Cancer* (2015)
3. Baldwin RT, Preston-Martin S. Epidemiology of brain tumors in childhood--a review. *Toxicology and applied pharmacology* (2004);199 (2):118-131. doi:10.1016/j.taap.2003.12.029
4. Spector LG, Pankratz N, Marcotte EL. Genetic and nongenetic risk factors for childhood cancer. *Pediatric clinics of North America* (2015);62 (1):11-25. doi:10.1016/j.pcl.2014.09.013
5. O'Neill KA, Murphy MF, Bunch KJ, Puumala SE, Carozza SE, Chow EJ, Mueller BA, McLaughlin CC, Reynolds P, Vincent TJ, Von Behren J, Spector LG. Infant birthweight and risk of childhood cancer: international population-based case control studies of 40 000 cases. *International journal of epidemiology* (2015);44 (1):153-168. doi:10.1093/ije/dyu265
6. Sprehe MR, Barahmani N, Cao Y, Wang T, Forman MR, Bondy M, Okcu MF. Comparison of birth weight corrected for gestational age and birth weight alone in prediction of development of childhood leukemia and central nervous system tumors. *Pediatric blood & cancer* (2010);54 (2):242-249. doi:10.1002/pbc.22308
7. Caughey RW, Michels KB. Birth weight and childhood leukemia: a meta-analysis and review of the current evidence. *International journal of cancer Journal international du cancer* (2009);124 (11):2658-2670. doi:10.1002/ijc.24225
8. Hjalgrim LL, Westergaard T, Rostgaard K, Schmiegelow K, Melbye M, Hjalgrim H, Engels EA. Birth weight as a risk factor for childhood leukemia: a meta-analysis of 18 epidemiologic studies. *American journal of epidemiology* (2003);158 (8):724-735
9. Oksuzyan S, Crespi CM, Cockburn M, Mezei G, Kheifets L. Birth weight and other perinatal factors and childhood CNS tumors: a case-control

study in California. *Cancer epidemiology* (2013);37 (4):402-409. doi:10.1016/j.canep.2013.03.007

10. Harder T, Plagemann A, Harder A. Birth weight and subsequent risk of childhood primary brain tumors: a meta-analysis. *American journal of epidemiology* (2008);168 (4):366-373. doi:10.1093/aje/kwn144

11. Schuz J, Forman MR. Birthweight by gestational age and childhood cancer. *Cancer causes & control : CCC* (2007);18 (6):655-663. doi:10.1007/s10552-007-9011-y

12. Bjorge T, Sorensen HT, Grotmol T, Engeland A, Stephansson O, Gissler M, Tretli S, Troisi R. Fetal growth and childhood cancer: a population-based study. *Pediatrics* (2013);132 (5):e1265-1275. doi:10.1542/peds.2013-1317

13. Oksuzyan S, Crespi CM, Cockburn M, Mezei G, Kheifets L. Birth weight and other perinatal characteristics and childhood leukemia in California. *Cancer epidemiology* (2012);36 (6):e359-365. doi:10.1016/j.canep.2012.08.002

14. Mallol-Mesnard N, Menegaux F, Lacour B, Hartmann O, Frappaz D, Doz F, Bertozzi AI, Chastagner P, Hemon D, Clavel J. Birth characteristics and childhood malignant central nervous system tumors: the ESCALE study (French Society for Childhood Cancer). *Cancer detection and prevention* (2008);32 (1):79-86. doi:10.1016/j.cdp.2008.02.003

15. Linet MS, Gridley G, Cnattingius S, Nicholson HS, Martinsson U, Glimelius B, Adami HO, Zack M. Maternal and perinatal risk factors for childhood brain tumors (Sweden). *Cancer causes & control : CCC* (1996);7 (4):437-448

16. Hjalgrim LL, Rostgaard K, Hjalgrim H, Westergaard T, Thomassen H, Forestier E, Gustafsson G, Kristinsson J, Melbye M, Schmiegelow K. Birth weight and risk for childhood leukemia in Denmark, Sweden, Norway, and Iceland. *Journal of the National Cancer Institute* (2004);96 (20):1549-1556. doi:10.1093/jnci/djh287

17. Buck Louis GM, Grewal J, Albert PS, Sciscione A, Wing DA, Grobman WA, Newman RB, Wapner R, D'Alton ME, Skupski D, Nageotte MP, Ranzini AC, Owen J, Chien EK, Craigo S, Hediger ML, Kim S, Zhang C, Grantz KL. Racial/ethnic standards for fetal growth: the NICHD Fetal Growth Studies. *American Journal of Obstetrics and Gynecology* (2015);213 (4): 449.e1-449.e41. doi: 10.1016/j.ajog.2015.08.032.

18. Sever LE, Gilbert ES, Tucker K, Greaves JA, Greaves C, Buchanan JA (1997) *Epidemiologic evaluation of Childhood leukemia and paternal exposure to ionizing radiation*. Seattle, USA

19. Oak Ridge Institute for Science and Education (ORISE). Comprehensive Epidemiologic Data Resource (CEDR). U.S. Department of Energy (DOE). <https://apps.ornl.gov/cedr/#.WVxB1YSGNdg>.
20. David WH, Stanley L (1989) Applied Logistic Regression. Wiley Series in Probability and Mathematical Statistics. Wiley-Interscience, United State of America
21. Introduction to Regression Modeling. In: Rothman KJ, Greenland S, Lash TH (eds) Modern Epidemiology. USA: Lippincott Williams & Wilkins; (2008).
22. Deval LP, Timothy PM, JudyAnn B, John A, Ron B, Judy H, Hafsatou D (2009) 2008 Pregnancy Data Report. Centers for Disease Control and Prevention (CDC), Massachusetts, USA
23. Alexander GR, Kogan MD, Himes JH. 1994-1996 U.S. singleton birth weight percentiles for gestational age by race, Hispanic origin, and gender. *Maternal and child health journal* (1999);3 (4):225-231
24. Surawicz TS, McCarthy BJ, Kupelian V, Jukich PJ, Bruner JM, Davis FG. Descriptive epidemiology of primary brain and CNS tumors: results from the Central Brain Tumor Registry of the United States, 1990-1994. *Neuro-oncology* (1999);1 (1):14-25
25. Von Behren J, Reynolds P. Birth characteristics and brain cancers in young children. *International journal of epidemiology* (2003);32 (2):248-256
26. Emerson JC, Malone KE, Daling JR, Starzyk P. Childhood brain tumor risk in relation to birth characteristics. *Journal of clinical epidemiology* (1991);44 (11):1159-1166
27. Heuch JM, Heuch I, Akslen LA, Kvale G. Risk of primary childhood brain tumors related to birth characteristics: a Norwegian prospective study. *International journal of cancer Journal international du cancer* (1998);77 (4):498-503
28. Kuijten RR, Bunin GR, Nass CC, Meadows AT. Gestational and familial risk factors for childhood astrocytoma: results of a case-control study. *Cancer research* (1990);50 (9):2608-2612
29. McCredie M, Little J, Cotton S, Mueller B, Peris-Bonet R, Choi NW, Cordier S, Filippini G, Holly EA, Modan B, Arslan A, Preston-Martin S. SEARCH international case-control study of childhood brain tumours: role of index pregnancy and birth, and mother's reproductive history. *Paediatric and perinatal epidemiology* (1999);13 (3):325-341
30. Schuz J, Kaletsch U, Kaatsch P, Meinert R, Michaelis J. Risk factors for pediatric tumors of the central nervous system: results from a German

population-based case-control study. *Medical and pediatric oncology* (2001);36 (2):274-282. doi:10.1002/1096-911X(20010201)36:2<274::AID-MPO1065>3.0.CO;2-D

31. Mogren I, Malmer B, Tavelin B, Damber L. Reproductive factors have low impact on the risk of different primary brain tumours in offspring. *Neuroepidemiology* (2003);22 (4):249-254. doi:70567

32. Gold E, Gordis L, Tonascia J, Szklo M. Risk factors for brain tumors in children. *American journal of epidemiology* (1979);109 (3):309-319

33. Ross JA, Perentesis JP, Robison LL, Davies SM. Big babies and infant leukemia: a role for insulin-like growth factor-1? *Cancer causes & control : CCC* (1996);7 (5):553-559

34. Del Valle L, Enam S, Lassak A, Wang JY, Croul S, Khalili K, Reiss K. Insulin-like growth factor I receptor activity in human medulloblastomas. *Clinical cancer research : an official journal of the American Association for Cancer Research* (2002);8 (6):1822-1830

35. Jukich PJ, McCarthy BJ, Surawicz TS, Freels S, Davis FG. Trends in incidence of primary brain tumors in the United States, 1985-1994. *Neuro-Oncology* (2001);3 (3): 141-151.

36. Dubois L, Girard M, Tatone-Tokuda F. Determinants of high birth weight by geographic region in Canada. *Chronic Diseases in Canada* (2007);28 (1-2):63-70.

37. Greenberg RS, Shuster JL Jr. Epidemiology of cancer in children. *Epidemiology Review* (1985);7 : 22-48.

38. Poole C, Greenland S, Luetters C, Kelsey JL, Mezei G. Socioeconomic status and childhood leukaemia: a review. *International journal of epidemiology* (2006);35 (2):370-384. doi:10.1093/ije/dyi248

39. Borugian MJ, Spinelli JJ, Mezei G, Wilkins R, Abanto Z, McBride ML. Childhood leukemia and socioeconomic status in Canada. *Epidemiology* (2005);16 (4):526-531.

40. Smith A, Roman E, Simpson J, Ansell P, Fear NT, Eden T. Childhood leukaemia and socioeconomic status: fact or artefact? A report from the United Kingdom childhood cancer study (UKCCS). *International Journal of Epidemiology* (2006);35 (6):1504-1513.

41. McNally RJ, Alston RD, Eden TO, Kelsey AM, Birch JM. Further clues concerning the aetiology of childhood central nervous system tumours. *European journal of cancer* (2004);40 (18):2766-2772. doi:10.1016/j.ejca.2004.08.020

42. Del Risco Kollerud R, Blaasaas KG, Claussen B. Poverty and the risk of leukemia and cancer in the central nervous system in children: A cohort study in a high-income country. *Scandinavian journal of public health* (2015);43 (7):736-743. doi:10.1177/1403494815590499
43. Keegan TJ, Bunch KJ, Vincent TJ, King JC, O'Neill KA, Kendall GM, MacCarthy A, Fear NT, Murphy MF. Case-control study of paternal occupation and social class with risk of childhood central nervous system tumours in Great Britain, 1962-2006. *British journal of cancer* (2013);108 (9):1907-1914. doi:10.1038/bjc.2013.171
44. Ramis R, Tamayo-Uria I, Gomez-Barroso D, Lopez-Abente G, Morales-Piga A, Pardo Romaguera E, Aragonés N, Garcia-Perez J. Risk factors for central nervous system tumors in children: New findings from a case-control study. *PloS one* (2017);12 (2):e0171881. doi:10.1371/journal.pone.0171881
45. Morley R. Fetal origins of adult disease. *Seminars in fetal & neonatal medicine* (2006);11 (2):73-78. doi:10.1016/j.siny.2005.11.001
46. de Boo HA, Harding JE. The developmental origins of adult disease (Barker) hypothesis. *The Australian & New Zealand journal of obstetrics & gynaecology* (2006);46 (1):4-14. doi:10.1111/j.1479-828X.2006.00506.x
47. Miles HL, Hofman PL, Cutfield WS. Fetal origins of adult disease: a paediatric perspective. *Reviews in endocrine & metabolic disorders* (2005);6 (4):261-268. doi:10.1007/s11154-005-6184-0

10 APPENDICES

Additional Table 1. The association between birth weight and CNS tumor risk without adjustment for gestational age

Birth weight	Controls	CNS tumors	OR	95%CI		P value
				Lower	Upper	
<2,500 g	24	7	4.1	1.6	10.1	0.007
2,500-4,000 g	718	53	1	Reference		
>4,000 g	80	12	2.0	1.0	4.1	0.058

P for homogeneity=0.006

ORs and corresponding 95% CIs and p values were adjusted for sex, ethnicity, year of birth, age at diagnosis, maternal age and DOE sites.

Description of data: The GA-unadjusted OR for high BW was 2.5 (95%CI=1.2, 5.2) when compared to normal BW (2,500-4,000 g).

Additional Table 2. The association between birth weight and the CNS tumor risk among children with gestational age of 37-42 weeks

Birth weight	Controls	CNS tumors	OR	95%CI		P value
				Lower	Upper	
Total subjects						
<2,500 g	11	3	3.8	0.9	15.7	0.091
2,500-<3,000 g	112	5	0.6	0.2	1.6	0.251
3,000-<3,500 g	288	19	1	Reference		
3,500-4,000 g	262	24	1.7	0.9	3.3	0.099
>4,000 g	75	11	3.6	1.5	8.7	0.005
				<i>P for homogeneity=0.007</i>		
				<i>P for trend=0.012</i>		
				<i>P for trend=0.001 (birth weight \geq2,500 g)</i>		
				<i>P for trend=0.019 (birth weight 2,500-4,000 g)</i>		
<2,500 g	11	3	3.4	0.9	13.2	0.111
2,500-4,000 g	662	48	1	Reference		
>4,000 g	75	11	2.8	1.3	5.9	0.014
				<i>P for homogeneity=0.016</i>		
The risk of high-birth-weight and LGA children compared to normal-birth-weight children*						
2,500-4,000 g	662	48	1	Reference		
>4,000 g and LGA	44	7	2.7	1.1	6.7	0.048
The risk of high-birth-weight and SGA/AGA children compared to normal-birth-weight children*						
2,500-4,000 g	662	48	1	Reference		
>4,000 g and GA/AGA	31	4	2.9	0.9	9.6	0.098

LGA: large for gestational age, SGA: small for gestational age, AGA: appropriate for gestational age
ORs and corresponding 95% CIs and p values were adjusted for sex, ethnicity, year of birth, age at diagnosis, gestational age (continuous variable), maternal age and DOE sites.

* Children with low-birth weight were not included in the analyses

Description of data: When compared to the results in table 2, the ORs and 95% CIs for high or low BW did not change appreciably.

Additional Table 3. The association between birth weight and leukemia risk among children with gestational age of 37-42 weeks

Birth weight	Controls	Leukemia cases	OR	95%CI		P value
				Lower	Upper	
<2,500 g	11	1	0.5	0.1	3.7	0.413
2,500-<3,000 g	112	18	0.8	0.4	1.4	0.425
3,000-<3,500 g	288	51	1	Reference		
3,500-4,000 g	262	30	0.7	0.4	1.1	0.106
>4,000 g	75	14	1.3	0.7	2.7	0.429
<i>P for homogeneity=0.275</i>						
<i>P for trend=0.715</i>						
<i>P for trend =0.314 (birth weight 2,500-4,000 g)</i>						
<2,500 g	11	1	0.5	0.1	4.4	0.529
2,500-4,000 g	662	99	1	Reference		
>4,000 g	75	14	1.6	0.8	3.1	0.158
<i>P for homogeneity=0.296</i>						

LGA: large for gestational age, SGA: small for gestational age, AGA: appropriate for gestational age

ORs and corresponding 95% CIs and p values were adjusted for sex, ethnicity, year of birth, age at diagnosis, gestational age (continuous variable), maternal age and DOE sites.

Description of data: When compared to the results in table 3, the ORs and 95% CIs for high or low BW did not change appreciably.

Additional Table 4. The risk of CNS tumors or leukemia among small-for-gestational-age and large-for-gestational-age children with gestational age of 37-42 weeks

Birth weight	Controls	Cases	OR	95%CI		P value
				Lower	Upper	
CNS tumors						
Total subjects						
SGA	186	15	0.9	0.5	2.0	0.963
AGA	507	39	1	Reference		
LGA	55	8	2.0	0.9	4.7	0.119
						<i>P for homogeneity=0.286</i>
Birth weight 2,500 g or larger						
SGA	175	12	0.8	0.4	1.7	0.563
AGA	507	39	1	Reference		
LGA	55	8	2.1	0.9	4.8	0.113
						<i>P for homogeneity=0.207</i>
Birth weight 3,000 g or larger						
SGA	97	9	1.1	0.4	2.9	0.797
AGA	473	37	1	Reference		
LGA	55	8	2.0	0.9	4.7	0.131
						<i>P for homogeneity=0.319</i>
LEUKEMIA						
Total subjects						
SGA	186	30	0.9	0.6	1.5	0.752
AGA	507	73	1	Reference		
LGA	55	11	1.8	0.8	3.7	0.148
						<i>P for homogeneity=0.306</i>
Birth weight 2,500 g or larger						
SGA	175	29	0.9	0.6	1.6	0.812
AGA	507	73	1	Reference		
LGA	55	11	1.7	0.8	3.6	0.157
						<i>P for homogeneity=0.334</i>
Birth weight 3,000 g or larger						
SGA	97	18	0.9	0.5	1.9	0.918
AGA	473	66	1	Reference		
LGA	55	11	1.9	0.9	4.0	0.112
						<i>P for homogeneity=0.267</i>

SGA: Small for gestational age, AGA: Appropriate for gestational age, LGA: Large for gestational age

ORs and corresponding 95% CIs and p values were adjusted for sex, ethnicity, year of birth, age at diagnosis, maternal age and DOE sites.

Description of data: When compared to the results in tables 5 and 6, the ORs for LGA/SGA did not change appreciably.