

# Novel pathogenic *XK* mutations in McLeod syndrome and interaction between *XK* protein and chorein

Yuka Urata, MD, Masayuki Nakamura, MD, PhD, Natsuki Sasaki, MD, PhD, Nari Shiokawa, MD, PhD, Yoshiaki Nishida, MD, Kaoru Arai, MD, Hanae Hiwatashi, MS, Izumi Yokoyama, BS, Shinsuke Narumi, MD, Yasuo Terayama, MD, PhD, Takenobu Murakami, MD, PhD, Yoshikazu Ugawa, MD, PhD, Hiroki Sakamoto, MD, Satoshi Kaneko, MD, PhD, Yusuke Nakazawa, MD, Ryo Yamasaki, MD, PhD, Shoko Sadashima, MD, Toshiaki Sakai, MD, Hiroaki Arai, MD, and Akira Sano, MD, PhD

## Correspondence

Dr. Nakamura  
nakamu36@  
m.kufm.kagoshima-u.ac.jp

*Neurol Genet* 2019;5:e328. doi:10.1212/NXG.0000000000000328

## Abstract

### Objective

To identify *XK* pathologic mutations in 6 patients with suspected McLeod syndrome (MLS) and a possible interaction between the chorea-acanthocytosis (ChAc)- and MLS-responsible proteins: chorein and *XK* protein.

### Methods

Erythrocyte membrane proteins from patients with suspected MLS and patients with ChAc, ChAc mutant carriers, and normal controls were analyzed by *XK* and chorein immunoblotting. We performed mutation analysis and *XK* immunoblotting to molecularly diagnose the patients with suspected MLS. Lysates of cultured cells were co-immunoprecipitated with anti-*XK* and anti-chorein antibodies.

### Results

All suspected MLS cases were molecularly diagnosed with MLS, and novel mutations were identified. The average onset age was  $46.8 \pm 8$  years, which was older than that of the patients with ChAc. The immunoblot analysis revealed remarkably reduced chorein immunoreactivity in all patients with MLS. The immunoprecipitation analysis indicated a direct or indirect chorein-*XK* interaction.

### Conclusions

In this study, *XK* pathogenic mutations were identified in all 6 MLS cases, including novel mutations. Chorein immunoreactions were significantly reduced in MLS erythrocyte membranes. In addition, we demonstrated a possible interaction between the chorein and *XK* protein via molecular analysis. The reduction in chorein expression is similar to that between Kell antigens and *XK* protein, although the chorein-*XK* interaction is a possibly noncovalent binding unlike the covalent Kell-*XK* complex. Our results suggest that reduced chorein levels following lack of *XK* protein are possibly associated with molecular pathogenesis in MLS.

From the Department of Psychiatry (Y. Urata, M.N., N. Sasaki, N. Shiokawa, Y. Nishida, K.A., H.H., I.Y., A.S.), Kagoshima University Graduate School of Medical and Dental Sciences; Department of Neurology and Gerontology (S.N., Y.T.), Iwate Medical University, Morioka; Department of Neurology (T.M.), School of Medicine, Fukushima Medical University; Department of Neuro-regeneration, Department of Neurology (Y. Ugawa), School of Medicine, Fukushima Medical University; Department of Neurology (H.S., S.K.), Kansai Medical University, Hirakata; Department of Neurology (Y. Nakazawa, R.Y., S.S.), Neurological Institute, Graduate School of Medical Sciences, Kyushu University, Fukuoka; Department of Neurology (T.S.), Nagano Matsushiro General Hospital; and Department of General Medicine (H.A.), Nagano Matsushiro General Hospital, Japan.

Funding information and disclosures are provided at the end of the article. Full disclosure form information provided by the authors is available with the full text of this article at [Neurology.org/NG](http://Neurology.org/NG).

The Article Processing Charge was paid for by the Authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

## Glossary

**MLS** = McLeod syndrome; **ChAc** = chorea-acanthocytosis; **NA** = neuroacanthocytosis.

---

Neuroacanthocytosis (NA) syndromes are rare neurodegenerative disorders exhibiting neurologic abnormalities and erythrocyte acanthocytosis. The core NA syndromes are characterized by degeneration of the striatum and huntingtonism. They comprise 2 main diseases: chorea-acanthocytosis (ChAc) and McLeod syndrome (MLS). ChAc is caused by loss-of-function mutations in *VPS13A*,<sup>1,2</sup> leading to an absent or markedly reduced level of the encoding protein, chorein.<sup>3,4</sup> MLS is caused by loss-of-function mutations in the *XK*, leading to absent XK protein.<sup>5</sup> Although later onset and cardiomyopathy may occur predominantly in MLS,<sup>6</sup> the 2 diseases share almost their entire symptomology in the CNS and erythrocyte membrane. Although molecular interactions are assumed to exist between these diseases, no studies have as yet established a direct association.

## Methods

### Human samples and mutation analysis

All 6 patients with suspected MLS were Japanese males with clinically suspected NA (table). Six healthy male controls and 6 male patients with ChAc were matched to suspected MLS cases by age. A further 6 heterozygous ChAc mutant carriers were used for the analyses. Lymphoblastoid cell lines from MLS\_1<sup>7</sup> and a healthy control were established by SRL (Tokyo, Japan).

Coding and flanking regions of *XK* (NC\_000023.10) and *VPS13A* (NC\_000009.11) were analyzed by Sanger sequencing on an ABI PRISM 3100 Avant Genetic Analyzer (Thermo Fisher Scientific, Waltham, MA).<sup>3</sup> In the case of MLS\_6, we performed a whole-genome sequence, long-range PCR covering the deletion region, and Sanger sequencing.

### Immunoprecipitation and immunoblot analysis

Co-immunoprecipitation (co-IP) and reverse co-IP assays were performed using Dynabeads Protein G (Thermo Fisher Scientific). K562 and HEK293 cells that stably overexpressed chorein<sup>8</sup> were lysed with Mammalian Protein Extraction Reagent (Thermo Fisher Scientific). K562 cells that were subcultured at  $1 \times 10^6$  cells/mL and incubated for 24 hours were used. The cell lysates (input) were used for the Dynabeads-antibody complex and Dynabeads-IgG complex. The cell lysate was diluted 5 times with  $1 \times$  Tris-buffered saline because delicate surfactant conditions were required to maintain the IP interaction. The cell lysate and each bead were incubated for 2 hours at room temperature.

Protein samples were analyzed by immunoblotting using rabbit anti-chorein (HPA021662; Atlas Antibodies, Bromma,

Sweden) and rabbit anti-XK protein (HPA019036; Atlas Antibodies) primary antibodies, which show no cross-reactivity with spectrin. Donkey anti-rabbit IgG, HRP-linked whole Ab (GE Health care, Little Chalfont, England) and VeriBlot for IP Detection Reagent (HRP) (ab131366; Abcam, Cambridge, UK) were used as secondary antibodies. Proteins were visualized using ECL Prime Western Blotting Detection Reagent (GE Health care), and images were recorded with a digital analyzer (FUSION-SOLO.7S.WL; Vilber Lourmat, Marne-la-Vallée, France).

### Standard protocol approvals, registrations, and patient consents

Genomic DNAs and/or proteins from peripheral blood samples were taken from all participants who provided written informed consent. The research protocol and consent form were approved by the Institutional Review Boards of Kagoshima University.

### Data availability statement

The data sets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Results

### Molecular diagnosis and clinical features of MLS cases

For all suspected MLS cases, XK protein immunoreactivity was lacking in the immunoblot analysis of the erythrocyte membrane (figure 1A). Clinical symptoms and pathologic *XK* mutations are presented in the table. In MLS\_6, comprehensive mutation analysis revealed a mutation, which was a combination of a gross deletion and an insertion (figure 1, B–D).

### Chorein immunoreactivity reductions in all MLS cases


We found a marked reduction in chorein immunoreactivity in all patients with MLS (figure 2A). The mean density level of patients with MLS was significantly lower ( $p = 0.00127$ ,  $d = 2.6$ ) at 0.55, relative to controls (figure 2D). The average reductions in the levels of chorein immunoreactivity in the erythrocyte membranes of MLS patients were equivalent to those found in heterozygous ChAc mutation carriers (figure 2, C and D), although no pathogenic mutations were identified in *VPS13A* in any patients with MLS. On the other hand, the average density levels of the XK immunoreactions did not significantly differ between ChAc and ChAc mutant carriers and healthy controls in either the immunoblot or densitometric analyses (figure 2, B–E). Chorein immunoreactions of

**Table** Profile of patients with MLS in this study

Clinical symptoms											XK pathogenic mutations			
Case	Age at sampling	Age at onset	Initial symptom	Main psychiatric symptom	Acanthocytes	Chorea	EEG abnormality	CK (IU/L)	DTRs	Cardiomyopathy	Atrophy of the corpus striatum on MRI	Exon	DNA change	Protein change
MLS_1	64	47	Muscle weakness	Insomnia	+	+	Intermittent theta wave	5465	Absent	-	+	Exon 3	c.del669_673TGAGinsGGTCCTTTTACC	p.V225Lfs*12
MLS_2	61	53	Involuntary movement	?	?	+	?	?	<sup>a</sup> Hyperreflexia	?	+	Exon 3	c.1013delT	p.F338Sfs*70
MLS_3	69	43	Muscle weakness	Persecutory delusion	+	?	?	920	Absent	+	?	Exon 2	c.451dupC	p.Q151Pfs*47
MLS_4	56	33	Involuntary movement	Depression	+	+	3-4 Hz slow wave	821	Normal	-	+	Exon 2	c.370C>T	p.Q124*
MLS_5	65	50	Involuntary movement	Cognitive decline	+	+	?	2,422	?	?	?	Exon 2	c.397C>T	p.R133*
MLS_6	70	55	Involuntary movement	Obsessiveness	+	+	?	1,052	Absent	-	-	Exon 3	Gross deletion	Unknown

Abbreviations: CK = creatine kinase; DTR = deep tendon reflex. Novel mutations were identified in MLS\_4 and MLS\_6. Age at onset: age when first signs or symptoms appeared (yr). <sup>a</sup>When DTR tests were performed, MLS\_2 was affected with bacterial meningitis.

**Figure 1** Molecular diagnosis of 6 MLS cases



(A) The results of XK immunoblotting revealed a lack of XK immunoreactivity in all patients with MLS. Equal loading was shown by staining with MemCode reversible protein stain (Pierce), shown in the lower panel. (B) In the case of MLS\_6, the *XK* gene mutation was predicted to be a gross deletion including exon 3 based on the results of gDNA amplification. To identify the breakpoints of this mutation, we performed a long-range PCR covering the deletion region. The results of long-range PCR for MLS\_6 showed a gross deletion mutation that was approximately 5500 bp in size. (C) Sanger sequencing results revealed a combination of a gross deletion, from intron 2 (c.509-636) to an *XK-CYBB* intergenic region (c.\*3667 + 670), and an insertion of a complementary sequence of 380 bp in intron 2. (D) The schematic shows the structure of the gross deletion region in MLS\_6.

the lymphoblastoid cell lysates from MLS\_1 and the control were equivalent. In addition, there was no immunoreaction corresponding to the XK protein band in both control and MLS\_1 lymphoblastoid cells (figure 2F).

### Chorein-XK protein interaction in cultured cells


Cell lysates extracted from K562 cells were immunoprecipitated with anti-XK antibody. In the subsequent immunoblot analysis, positive chorein bands were detected in the XK immunoprecipitants (figure 2G). Because the endogenous chorein level was low, XK protein immunoreactivity was not visually observed in chorein immunoprecipitants. Therefore, co-IP and reverse co-IP assays were conducted in a similar manner using the lysate extracted from chorein stably

overexpressing HEK293 cells. Signals positive for chorein and XK protein were detected in the XK and chorein immunoprecipitants, respectively (figure 2H).

### Discussion

In the present study, we analyzed 6 cases with MLS and confirmed the molecular diagnosis, as well as identifying 2 additional novel pathogenic mutations (table). The profile of 6 cases of MLS in this study was similar to those reported previously.<sup>6</sup> In our MLS cases, the average onset age was 46.8 ± 8 years, which is approximately 13 years older than that of patients with ChAc.<sup>3</sup> The disease duration for MLS may be longer than 30 years, which is typically longer than for ChAc.<sup>6</sup>

**Figure 2** Immunoblotting of erythrocyte membranes, lymphoblastoid cells, and co-immunoprecipitants



(A and C) Chorein immunoblotting results revealed reduced chorein immunoreactivity in the erythrocyte membranes of patients with MLS, which are equivalent to the heterozygous mutant carriers of ChAc (ChAc\_hetero). (B and C) XK immunoblotting results revealed normal XK immunoreactivity in the erythrocyte membranes of ChAc and ChAc\_hetero. (A–C) Each imaging was performed by underexposed condition for semiquantitative analysis. (D) The histograms show the chorein relative protein density ratio in patients with MLS and ChAc and ChAc\_hetero. Each band density was normalized by protein density staining with MemCode reversible protein stain. Error bars represent the SD (each n = 6); 2-sample Student t tests were performed accordingly on different sets of data. \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ .  $d$  shows the effect size (Cohen's  $d$ ). (E) The histograms show the XK protein relative density ratio in patients with MLS and ChAc and ChAc\_hetero. Each band density was normalized by protein density staining with memcode reversible protein stain. (F) Chorein immunoblotting results revealed equivalent chorein immunoreactivity in the lymphoblastoid cells of MLS\_1 and the control (MLS\_1\_lympho and Control\_lympho, respectively). The XK immunoblotting of lymphoblastoid cells shown in the lower panel of figure 2F reveals a lack of XK immunoreactivity. The Control\_EM and MLS\_EM lanes show the erythrocyte membrane for control and MLS, respectively. (G) Co-immunoprecipitation (IP) assay using K562 cells was performed with anti-XK antibody. Immunoblot analyses used anti-chorein (upper panel) and anti-XK antibodies (lower panel). (H) Co-IP and reverse co-IP assays using human embryonic kidney 293 (HEK293) cells stably overexpressing Myc-DDK-tagged chorein confirmed an interaction between XK protein and chorein. Immunoblot analyses used anti-XK protein and anti-chorein antibodies.

In the present study, semiquantitative chorein immunoblotting using erythrocyte membranes from all patients with MLS revealed significantly reduced chorein immunoreactions compared with age- and sex-matched healthy controls (figure 2A). Chorein immunoreactivities in heterozygous ChAc mutation carriers are also reduced to the same level as in patients with MLS (figure 2, C and D). These findings were demonstrated in at least triplicate independent experiments. Some ChAc mutation carriers exhibit partial symptoms of NA such as acanthocytosis.<sup>9</sup> Taken together, the later onset and slower progression found for MLS compared with ChAc suggest that the chorein level reductions found in MLS may be directly associated with MLS molecular pathology. The erythrocyte membrane from 1 patient with MLS and lymphoblastoid cells from another patient with MLS showed normal chorein levels in previous study.<sup>4</sup> In that study, chorein immunoblotting of heterozygous ChAc mutant carriers showed normal chorein levels, suggesting that the results

of immunoblotting analysis might be unavailable for semiquantification.

In the present study, XK immunoblotting of lymphoblastoid cell lysate from healthy controls showed no XK protein band, suggesting no expression of XK protein in lymphoblastoid cells. This may account for the normal chorein immunoreaction found by chorein immunoblotting of lymphoblastoid cell lysate from MLS\_1, although further investigation is required.

XK protein covalently interacts with Kell antigens, which are remarkably reduced in erythrocyte membranes of MLS patients.<sup>6</sup> In this study, based on the finding of reduced chorein in the erythrocyte membranes of patients with MLS, we hypothesized that the XK protein directly or indirectly interacts with chorein. In the present study, we performed IP assays, which revealed the possible interaction. In erythrocyte

membranes, the absence of XK led to reduced chorein levels, although the absence of chorein was unrelated to XK levels. Computational analysis revealed a number of the corresponding impaired phosphorylation pathways in MLS and ChAc, suggesting a common molecular background bridging the generation of acanthocytes.<sup>10</sup> In the present study, protein staining on blotting membranes revealed upshift in band 3 from both MLS and ChAc (data not shown), suggesting results of phosphorylation. Taken together, these results suggests that reduced chorein is associated with MLS phosphorylation-related molecular pathology in the erythrocyte membranes. However, the direct mechanisms of reduced chorein in erythrocytes of MLS are unknown. In addition, our study did not include molecular investigations of the CNS. Further studies are needed to elucidate the molecular mechanism of NA.

## Acknowledgment

The authors thank the patients with MLS, patients with ChAc, ChAc mutant carriers, and healthy controls for their participation, and Ms. Kyoko Meguro for her technical assistance.

## Study funding

This study was supported by Grants-in Aid from the Research Committee of CNS Degenerative Diseases, Research on Policy Planning and Evaluation for Rare and Intractable Diseases, Health, Labour and Welfare Sciences Research Grants, and the Ministry of Health, Labour and Welfare, Japan, and in part by the Ministry of Education, Culture, Sports, Science and Technology KAKENHI (Grant No. 17H04250 to A.S. and No. 18K07606 to M.N.).

## Disclosure

Disclosures available: [Neurology.org/NG](http://Neurology.org/NG).

## Publication history

Received by *Neurology: Genetics* November 18, 2018. Accepted in final form March 7, 2019.

## Appendix Authors

Name	Location	Role	Contribution
<b>Yuka Urata, MD</b>	Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima	Author	Performed laboratory work and data analysis and prepared the manuscript
<b>Masayuki Nakamura, MD, PhD</b>	Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima	Corresponding author	Supervised the project, advised on laboratory work and data analysis, and prepared the manuscript
<b>Natsuki Sasaki, MD, PhD</b>	Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima	Author	Advised on laboratory work

## Appendix (continued)

Name	Location	Role	Contribution
<b>Nari Shiokawa, MD, PhD</b>	Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima	Author	Advised on laboratory work
<b>Yoshiaki Nishida, MD</b>	Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima	Author	Advised on data analysis
<b>Kaoru Arai, MD</b>	Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima	Author	Advised on data analysis
<b>Hanae Hiwatashi, MS</b>	Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima	Author	Performed laboratory work
<b>Izumi Yokoyama, BS</b>	Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima	Author	Performed laboratory work
<b>Shinsuke Narumi, MD</b>	Iwate Medical University, Morioka	Author	Collected clinical data and blood samples of study patients
<b>Yasuo Terayama, MD, PhD</b>	Iwate Medical University, Morioka	Author	Collected clinical data and blood samples of study patients
<b>Takenobu Murakami, MD, PhD</b>	Fukushima Medical University, Fukushima	Author	Collected clinical data and blood samples of study patients
<b>Yoshikazu Ugawa, MD, PhD</b>	Fukushima Medical University, Fukushima	Author	Collected clinical data and blood samples of study patients
<b>Hiroki Sakamoto, MD</b>	Kansai Medical University, Hirakata	Author	Collected clinical data and blood samples of study patients
<b>Satoshi Kaneko, MD, PhD</b>	Kansai Medical University, Hirakata	Author	Collected clinical data and blood samples of study patients
<b>Yusuke Nakazawa, MD</b>	Kyushu University, Fukuoka	Author	Collected clinical data and blood samples of study patients
<b>Ryo Yamasaki, MD, PhD</b>	Kyushu University, Fukuoka	Author	Collected clinical data and blood samples of study patients
<b>Shoko Sadashima, MD</b>	Kyushu University, Fukuoka	Author	Collected clinical data and blood samples of study patients

## Appendix *(continued)*

Name	Location	Role	Contribution
<b>Toshiaki Sakai, MD</b>	Nagano Matsushiro General Hospital, Nagano	Author	Collected clinical data and blood samples of study patients
<b>Hiroaki Arai, MD</b>	Nagano Matsushiro General Hospital, Nagano	Author	Collected clinical data and blood samples of study patients
<b>Akira Sano, MD, PhD</b>	Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima	Author	Advised on data analysis, prepared the manuscript, and served as a scientific advisor

## References

1. Ueno S, Maruki Y, Nakamura M, et al. The gene encoding a newly discovered protein, chorein, is mutated in chorea-acanthocytosis. *Nat Genet* 2001;28:121–122.
2. Rampoldi L, Dobson-Stone C, Rubio JP, et al. A conserved sorting-associated protein is mutant in chorea-acanthocytosis. *Nat Genet* 2001;28:119–120.
3. Tomiyasu A, Nakamura M, Ichiba M, et al. Novel pathogenic mutations and copy number variations in the VPS13A Gene in patients with chorea-acanthocytosis. *Am J Med Genet B* 2011;156:620–631.
4. Dobson-Stone C, Velayos-Baeza A, Filippone LA, et al. Chorein detection for the diagnosis of Chorea-acanthocytosis. *Ann Neurol* 2004;56:299–302.
5. Ho M, Chelly J, Carter N, Danek A, Crocker P, Monaco AP. Isolation of the gene for McLeod syndrome that encodes a novel membrane transport protein. *Cell* 1994;77:869–880.
6. Jung HH, Danek A, Walker RH. Neuroacanthocytosis syndromes. *Orphanet J Rare Dis* 2011;6:68.
7. Narumi S, Natori T, Miyazawa H, et al. Case of McLeod syndrome with a novel genetic mutation. *Neurol Clin Neurosci* 2016;4:115–117.
8. Shiokawa N, Nakamura M, Sameshima M, et al. Chorein, the protein responsible for chorea-acanthocytosis, interacts with  $\beta$ -adducin and  $\beta$ -actin. *Biochem Biophys Res Commun* 2013;441:96–101.
9. Ichiba M, Nakamura M, Kusumoto A, et al. Clinical and molecular genetic assessment of a chorea-acanthocytosis pedigree. *J Neurol Sci* 2007;263:124–132.
10. De Franceschi L, Scardoni G, Tomelleri C, et al. Computational identification of phospho-tyrosine sub-networks related to acanthocyte generation in neuroacanthocytosis. *PLoS One* 2012;7:e31015.