

平成24年退職予定者の研究業績

坂田祐介氏（生物生産学科 園芸生産学講座 観賞園芸学研究室）

学術論文

- 坂田祐介・上本俊平：スイートピーの花色に関する研究（第1報）。春咲種のアントシアニン色素生成について。園芸学会雑誌, 45 (2), 181-186 (1976)
- 坂田祐介・永吉実孝・有隅健一：ツバキ属植物の花色素に関する研究. I. 長崎県平戸市周辺におけるヤブツバキ, サザンカおよびハルサザンカのアントシアニン色素について. 鹿大農学術報告, 30, 35-40 (1980)
- 坂田祐介・有隅健一：トサカケイトウのベタレイン系色素生成に関する研究. I. 光の影響について. 鹿大農学術報告, 30, 41-46 (1980)
- Sakata, Y., Nagayoshi, S. and Arisumi, K.: Studies on the flower colours in the *Camellia*. II. On the anthocyanin constitution in the cultivars of *C. japonica*, *C. japonica* subsp. *rusticana* *C. sasanqua*, *C. hiemalis*, *C. vernalis* and *C. wabisuke*. Mem. Fac. Agr. Kagoshima Univ. 17, 79-94 (1981)
- Matsuo, E., Arisumi, K., Ohmachi, K. and Sakata, Y.: Effect of scale planting depth on leaf and shoot emergence in scale-propagated Easter lily. HortScience, 17, 806-807 (1982)
- 坂田祐介・有隅健一：スイートピーの花色に関する研究(第2報)。種々の品種群のアントシアニンおよびフラボノール色素構成について。鹿大農学術報告, 33, 13-19 (1983)
- Arisumi, K., Sakata, Y. and Miyajima, I.: Studies on the flower colours in *Rhododendron*. I. Pigment constitutions of the elepidote and lepidote rhododendrons. Mem. Fac. Agr. Kagoshima Univ., 21, 133-147 (1985)
- Sakata, Y. and Arisumi, K.: Studies on the flower colours in the *Camellia*. On the anthocyanin constitution in *C. reticulata*, *C. saluenensis*, *C. pitardii*, *C. hongkongensis*, *C. rosaeiflora* and *C. maliflora* and in the cultivars derived from interspecific hybridization. Mem. Fac. Agr. Kagoshima Univ., 21, 149-156 (1985)
- Sakata, Y., Arisumi, K. and Miyajima, I.: Cyanidin 3-galactoside, a new anthocyanin from *Camellia japonica* subsp. *rusticana* (Honda) Kitamura and its occurrence in the garden forms of *Camellia* of Japanese origin. J. Japan. Soc. Hort. Sci., 55 (1), 82-88 (1986)
- Sakata, Y., Arisumi, K. and Miyajima, I.: Constitution of anthocyanins in flowers of the wild forms of section Camellia of Japanese and Formosan origin. J. Japan. Soc. Hort. Sci., 56 (2), 208-214 (1987)
- Matsuo, E., Matsuzawa, M., Sakata, Y. and Arisumi, K.: Asexual propagation of variegated *Lilium longiflorum* 'Chotaro'. Scientia Hortic., 39, 349-354 (1989)
- Sakata, Y., Arisumi, K. and Miyajima, I.: Some morphological and pigmental characteristics in *Rhododendron kaempferi* Planch., *R. kiusianum* Makino and *R. eriocarpum* Nakai in southern Kyushu. J. Japan. Soc. Hort. Sci., 60 (3), 669-675 (1991)
- Sakata, Y. and Arisumi, K.: Anthocyanins in *Camellia polyodonta* How, *C. semiserrata* Chi and *C. chekiangoleosa* Hu, and their phylogenetic positions in section Camellia based on the pigmentation. J. Japan. Soc. Hort. Sci., 61 (2), 375-381 (1992)
- Terahara, N., Saito, N., Toki, K., Sakata, Y. and Honda, T.: Cyanidin 3-lathyroside from berries of *Fatsia japonica*. Phytochemistry, 31 (4), 1446-1448 (1992)
- Sakata, Y., Miyajima, I. and Arisumi, K.: Variations in some morphological and pigmental characteristics in *Rhododendron kaempferi* Planch., *R. kiusianum* Makino and their natural hybrids on Kirishima mountain mass. J. Japan. Soc. Hort. Sci., 61 (4), 925-932 (1993)
- 上野敬一郎・野添博昭・坂田祐介・有隅健一：鹿児島県下で見出された*Lycoris*新種の成立について。園芸学会雑誌, 63 (2), 409-417 (1994)
- Sakata, Y., Aoki, N., Tsunematsu, S., Nishikouri, H. and Johjima, T.: Petal coloration and pigmentation of tree peony bred and selected in Daikon Island. J. Japan. Soc. Hort. Sci., 64 (2), 351-357 (1995)
- Miyajima, I., Uemoto, S., Sakata, Y. and Arisumi, K.: Morphology and flower pigments of wild evergreen azaleas (*Rhododendron sataense* Nakai) in southern Kyushu. J. Japan. Soc. Hort. Sci., 66 (2), 385-391 (1997)

- 松添直隆・山口雅篤・川伸修治・渡部由香・東 華枝・坂田祐介：果実への暗黒処理がナスの果色と果皮のアントシアニン組成に及ぼす影響。園芸学会雑誌, 68 (1), 138-145 (1999)
- Hashimoto, F., Tanaka, M., Maeda, H., Shimizu, K. and Sakata, Y.: Characterization of cyanic flower color of *Delphinium* cultivars. J. Japan. Soc. Hort. Sci., 69 (4), 428-434 (2000)
- 宮島郁男・黒瀬克之・松田鹿徳・上本俊平・坂田祐介：九重、阿蘇山系およびその周辺地域におけるミヤマキリシマ (*Rhododendron kiusianum* Makino) の花器と花色素構成の変異。園芸学会雑誌, 70 (1), 108-114 (2001)
- Wang, L.-S., Shiraishi, A., Hashimoto, F., Aoki, N., Shimizu, K. and Sakata, Y.: Analysis of petal anthocyanins to investigate flower coloration of Zhongyuan (Chinese) and Daikon island (Japanese) tree peony cultivars. J. Plant Res., 114 (1113), 33-43 (2001)
- Wang, L.-S., Hashimoto F., Shiraishi, A., Aoki, N., Li, J.-J., Shimizu, K. and Sakata, Y.: Phenetics in tree peony species from China by flower color cluster analysis. J. Plant Res., 114 (1115), 213-221 (2001)
- Uddin, A. F. M. J., Hashimoto, F., Nishimoto, S., Shimizu, K. and Sakata, Y.: Flower growth, coloration and petal pigmentation of four *lislanthus* cultivars. J. Japan. Soc. Hort. Sci., 71 (1) : n40-47 (2002)
- Hashimoto, F., Tanaka, M., Maeda, H., Fukuda, S., Shimizu, K. and Sakata, Y.: Changes in flower coloration and sepal anthocyanins in cyanic *Delphinium* cultivars during flowering. Biosci. Biotechnol. Biochem., 66 (8), 1652-1659 (2002)
- 西本慎一・清水圭一・橋本文雄・坂田祐介：胚珠培養によるツバキ属種間雑種（キンカチャ×ヤブツバキ）の作出。園芸学会雑誌, 72 (3), 236-242 (2003)
- Shimizu, K., Hashimoto, M., Hashimoto, F. and Sakata, Y.: Plant regeneration from suspension cultures in Japanese morning glory (*Ipomoea nil* (L.) Roth.). J. Japan. Soc. Hort. Sci., 72 (5), 409-414 (2003)
- Uddin, A. F. M. J., Hashimoto, F., Shimizu, K. and Sakata, Y.: Seasonal variation in pigmentation and anthocyanidin phenetics in commercial *Eustoma* flowers. Scientia Hortic., 100, 103-115 (2004)
- Wang, L.-S., Hashimoto, F., Shiraishi, A., Aoki, N., Li, J.-J. and Sakata, Y.: Chemical taxonomy of the Xibei tree peony from China by floral pigmentation. J. Plant Res., 117, 47-55 (2004)
- Hashimoto, F., Jamal Uddin, A. F. M., Shimizu, K. and Sakata, Y.: Multiple allelism in flavonoid hydroxylation in *Eustoma grandiflorum* (Raf.) Shinn. flowers. J. Japan. Soc. Hort. Sci., 73(3), 235-240 (2004)
- Shimizu, K., Yogai, T., Hashimoto, F. and Sakata, Y.: Plant regeneration from suspension cell culture of *Delphinium*. J. Japan. Soc. Hort. Sci., 73(5), 435-440 (2004)
- Shimizu, K., Tokumura, T., Hashimoto, F. and Sakata, Y.: In vitro propagation of sterile mutant strains in Japanese morning glory by culture of embryoid derived from immature embryo. J. Japan. Soc. Hort. Sci., 74(4), 311-317 (2005)
- Li, Jian-Bin, Hashimoto, F., Shimizu, K. and Sakata, Y.: Anthocyanins from the red flowers of *Camellia reticulata* Lindl. Biosci. Biotechnol. Biochem., 71(11), 2833-2836 (Nov. 2007)
- Li, Jian-Bin, Hashimoto, F., Shimizu, K. and Sakata, Y.: Anthocyanins from the red flowers of *Camellia saluenensis* Stapf ex Bean. J. Japan. Soc. Hort. Sci., 77(1), 75-79 (Jan. 2008)
- Li, Jian-Bin, Hashimoto, F., Shimizu, K. and Sakata, Y.: Anthocyanins from the red flowers of Tali camellia (Dalicha). Phytochemistry, 69, 3166-3171 (2008)
- Li, Jian-Bin, Hashimoto, F., Shimizu, K. and Sakata, Y.: A new anthocyanin from the red flowers of *Camellia hongkongensis* and characterization of anthocyanins in the Section Camellia species. Jour. Integrative Plant Biol., 51(6), 545-552 (2009)
- Li, Jia-Hua, Nesumi, A., Shimizu, K., Sakata, Y., Liang, Ming-Zhi, He, Qing-Yuan, Zhou, Hong-Jie and Hashimoto, F.: Chemosystematics of tea trees based on tea leaf polyphenols as phenetic markers. Phytochemistry, 71, 1342-1349 (2010).
- Li, Jia-Hua, Zhang, Guag-Hui, Shao, Wan-Fang, Sakata, Y., Nesumi, A. and Hashimoto, F.: Hereditary analysis of tea polyphenols in *Camellia sinensis* F₁ hybrids based on Principal Component and Cluster Analysis. Molecular Plant Breeding, 9, 491-497 (2011)

藤崎幸藏氏（獣医学科 新興感染症学研究室）

学術論文

- Battsetseg, B., Matsuo, T., Xuan, X., Boldbaatar D., Chee, H.S., Umemiya, R., Sakaguchi, T., Hatta, T., Zhou, J., Rodolfo, V.A., Taylor, D., Fujisaki, K.: *Babesia* parasites develop and are transmitted by the non-vector soft tick *Ornithodoros moubata* (Acari: Argasidae). *Parasitology* 134, 1-8 (2007).
- Miyoshi, T., Tsuji, N., Islam, M.K., Huang, X., Motobu, M., Alim, M.A., Fujisaki, K.: Molecular and reverse genetic characterization of serine proteinase-induced hemolysis in the midgut of the ixodid tick *Haemaphysalis longicornis*. *J. Insect Physiol.* 53, 195-203 (2007).
- Umemiya, R., Hatta, T., Liao, M., Tanaka, M., Zhou, J., Inoue, N., Fujisaki, K.: *Haemaphysalis longicornis*: Molecular characterization of a homologue of the macrophage migration inhibitory factor from the partially fed ticks. *Exp. Parasitol.* 115, 135-142 (2007).
- Huang, X., Tsuji, N., Miyoshi, T., Nakamura-Tsuruta, S., Hirabayashi, J., Fujisaki, K.: Molecular characterization and oligosaccharide binding properties of a galectin from the argasid tick *Ornithodoros moubata*. *Glycobiology* 17, 313-323 (2007).
- Huang, X., Tsuji, N., Miyoshi, T., Motobu, M., Islam, M.K., Alim, M.A., Fujisaki, K.: Characterization of glutamine: fructose-6-phosphate aminotransferase from the ixodid tick, *Haemaphysalis longicornis*, and its critical role in host blood feeding. *Int. J. Parasitol.* 37, 383-392 (2007).
- Liao, M., Zhou, J., Hatta, T., Umemiya, R., Miyoshi, T., Tsuji, N., Xuan, X., Fujisaki, K.: Molecular characterization of *Rhipicephalus (Boophilus) microplus* Bm86 homologue from *Haemaphysalis longicornis* ticks. *Vet. Parasitol.* 146, 148-157 (2007).
- Zhou, J., Liao, M., Ueda, M., Gong, H., Xuan, X., Fujisaki, K.: Sequence characterization and expression patterns of two defensin-like antimicrobial peptides from the tick *Haemaphysalis longicornis*. *Peptides* 28, 1304-1310 (2007).
- Boldbaatar, D., Battsetseg, B., Hatta, T., Miyoshi, T., Tsuji, N., Xuan, X., Fujisaki, K.: Valosin-containing protein from the hard tick *Haemaphysalis longicornis*: effects of dsRNA-mediated HLVCP gene silencing. *Biochem. Cell Biol.* 85, 384-394 (2007).
- Motobu, M., Tsuji, N., Miyoshi, T., Huang, X., Islam, M.K., Alim, M.A., Fujisaki, K.: Molecular characterization of a blood-induced serine carboxypeptidase from the ixodid tick *Haemaphysalis longicornis*. *FEBS J.* 274, 3299-3312 (2007).
- Tsuji, N., Battsetseg, B., Boldbaatar, D., Miyoshi, T., Xuan, X., Oliver, J.H. Jr., Fujisaki, K.: Babesial vector tick defensin against *Babesia parasites*. *Infect. Immun.* 75, 3633-3640 (2007).
- Liao, M., Hatta, T., Umemiya, R., Huang, P., Jia, H., Gong, H., Zhou, J., Nishikawa, Y., Xuan, X., Fujisaki, K.: Identification of three disulfide isomerase members from *Haemaphysalis longicornis*. *Insect Biochem. Mol. Biol.* 37, 641-654 (2007).
- Alim, M.A., Tsuji, N., Miyoshi, T., Islam, M.K., Huang, X., Motobu, M., Fujisaki, K.: Characterization of asparaginyl endopeptidase, legumain induced by blood feeding in the ixodid tick *Haemaphysalis longicornis*. *Insect Biochem. Mol. Biol.* 37, 911-922 (2007).
- Umemiya, R., Matsuo, T., Hatta, T., Sakakibara, S., Boldbaatar, D., Fujisaki, K.: Cloning and characterization of an autophagy-related gene, ATG12, from the three-host tick *Haemaphysalis longicornis*. *Insect Biochem. Mol. Biol.* 37, 975-984 (2007).
- Gong, H., Zhou, J., Liao, M., Hatta, T., Harnnoi, T., Umemiya, R., Inoue, N., Xuan, X., Fujisaki, K.: Characterization of a carboxypeptidase inhibitor from the tick *Haemaphysalis longicornis*. *J. Insect Physiol.* 53, 1079-1087 (2007).
- Umemiya, R., Matsuo, T., Hatta, T., Sakakibara, S., Boldbaatar, D., Fujisaki, K.: Autophagy-related genes

- from a tick, *Haemaphysalis longicornis*. Autophagy 4, 79-81 (2008).
- Gong, H., Liao, M., Zhou, J., Hatta, T., Huang, P., Zhang, G., Kanuka, H., Nishikawa, Y., Xuan, X., Fujisaki, K.: Gene silencing of ribosomal protein P0 is lethal to the tick *Haemaphysalis longicornis*. Vet. Parasitol. 151, 268-278 (2008).
 - Liao, M., Boldbaatar, D., Gong, H., Huang, P., Umemiya, R., Harnnoi, T., Zhou, J., Tanaka, T., Suzuki, H., Xuan, X., Fujisaki, K.: Functional analysis of protein disulfide isomerase in blood feeding, viability and oocyte development in *Haemaphysalis longicornis* ticks. Insect Biochem. Mol. Biol. 38, 285-295 (2008).
 - Alim, M.A., Tsuji, N., Miyoshi, T., Islam, M.K., Huang, X., Hatta, T., Fujisaki, K.: HlLgm2, a member of asparaginyl endopeptidases/legumains in the midgut of the ixodid tick *Haemaphysalis longicornis*, is involved in blood-meal digestion. J. Insect Physiol. 54, 573-585 (2008).
 - Tsuji, N., Miyoshi, T., Battsetseg, B., Matsuo, T., Xuan, X., Fujisaki, K.: A cysteine protease is critical for *Babesia* spp. transmission in *Haemaphysalis* ticks. PLoS Pathogen 4, e1000062 (2008).
 - Boldbaatar, D., Musyoka, K.R., Battur, B., Umemiya, R., Tanaka, T., Xuan, X., Fujisaki, K.: Identification of two forms of Cyclophilin from the hard tick *Haemaphysalis longicornis*. Process Bicohem. 43, 615-625 (2008).
 - Boldbaatar, B., Battsetseg, B., Matsuo, T., Hatta, T., Umemiya-Shirafuji, R., Xuan, X., Fujisaki, K.: Tick vitellogenin receptor reveals critical role in oocyte development and transovarial transmission of *Babesia* parasite. Biochem. Cell Biol. 86, 331-344 (2008).
 - Miyoshi, T., Tsuji, N., Islam, M.K., Alim, M.A., Hatta, T., Huang, X., Fujisaki, K.: A set of serine proteinase paralogs are required for blood-digestion in the ixodid tick *Haemaphysalis longicornis*. Parasitol. Int. 57, 499-505 (2008).
 - Alim, M.A., Tsuji, N., Miyoshi, T., Islam, M.K., Hatta, T., Fujisaki, K.: Legumains from the hard tick *Haemaphysalis longicornis* play modulatory roles in blood feeding and gut cellular remodeling and impact on embryogenesis. Int. J. Parasitol. 39, 97-107 (2009).
 - Liao, M., Zhou, J., Gong, H., Boldbaatar, D., Shirafuji-Umemiya, R., Battur, B., Dong, J., Nishikawa, Y., Fujisaki, K.: Hemalin, a thrombin inhibitor isolated from a midgut cDNA library from the hard tick *Haemaphysalis longicornis*. J. Insect Physiol. 55, 165-174 (2009).
 - Zhou, J., Liao, M., Ueda, M., Gong, H., Xuan, X., Fujisaki, K.: Characterization of an intracellular cystatin homolog from the tick *Haemaphysalis longicornis*. Vet. Parasitol. 160, 180-183 (2009).
 - Gong, H., Umemiya, R., Zhou, J., Liao, M., Zhang, H., Jia, H., Nishikawa, Y., Xuan, X., Fujisaki, K.: Blocking the secretion of saliva by silencing the *HlYkt* gene in the tick *Haemaphysalis longicornis*. Insect Biochem. Mol. Biol. 39, 372-381 (2009).
 - Islam, M.K., Tsuji, N., Miyoshi, T., Alim, M.A., Huang, X., Hatta, T., Fujisaki, K.: The hard tick Kunitz inhibitor, Haemangin, is a novel modulator of angiogenesis and wound healing and is crucial for slower blood-feeding arthropods. PLoS Pathogen 5, e1000497 (2009).
 - Battsetseg, B., Boldbaatar, D., Battur, B., Xuan, X., Fujisaki, K.: Cloning and molecular characterization of tick kynureine aminotransferase (HlKAT) from *Haemaphysalis longicornis* (Acar: Ixodidae). Parasitol. Res. 105, 669-679 (2009).
 - Battur, B., Boldbaatar, D., Umemiya-Shirafuji, R., Liao, M., Battsetseg, B., Baymbaa, B., Fujisaki, K.: LKR/SDH plays important roles throughout the tick life cycle including a long starvation period. PLoS One 4, e7136 (2009).
 - Yamaji, K., Tsuji, N., Miyoshi, T., Islam, M.K., Hatta, T., Alim, M.A., Anisuzzaman, Takenaka, A., Fujisaki, K.: Hemoglobinase activity of a cysteine protease from the ixodid tick *Haemaphysalis longicornis*. Parasitol. Int. 58, 232-237 (2009).
 - Boldbaatar, D., Battur, B., Umemiya-Shirafuji, R., Liao, M., Tanaka, T., Xuan, X., Fujisaki, K.: GATA

-
- transcription, translation and regulation in *Haemaphysalis longicornis* tick: analysis of the cDNA and essential role for vitellogenesis. Insect Biochem. Mol. Biol. 40, 49-57 (2010).
- Miyoshi, T., Tsuji, N., Islam, M.K., Alim, M.A., Hatta, T., Yamaji, K., Anisuzzaman, Fujisaki, K.: A Kunitz-type proteinase inhibitor from the midgut of the ixodid tick, *Haemaphysalis longicornis*, and its endogenous target serine proteinase. Mol. Biochem. Parasitol. 170, 112-115 (2010).
 - Umemiya-Shirafuji, R., Matsuo, T., Liao, M., Boldbaatar, D., Battur, B., Suzuki, H., Fujisaki, K.: Increased expression of ATG genes during nonfeeding periods in the tick *Haemaphysalis longicornis*. Autophagy 6, 473-481 (2010).
 - Hatta, T., Tsuji, N., Miyoshi, T., Islam, M.K., Alim, M.A., Yamaji, K., Anisuzzaman, Fujisaki, K.: Leucine aminopeptidase, HILAP, from the ixodid tick *Haemaphysalis longicornis*, plays vital roles in the development of oocytes. Parasitol. Int. 59, 286-289 (2010).
 - Zhou, J., Liao, M., Gong, H., Xuan, X., Fujisaki, K.: Characterization of Hlcyst-3 as a member of cystatins from the tick *Haemaphysalis longicornis*. Exp. Appl. Acarol. 51, 327-333 (2010).
 - Boldbaatar, D., Battur, B., Umemiya-Shirafuji, R., Liao, M., Tanaka, T., Xuan, X., Fujisaki, K.: Multiple vitellogenins from the *Haemaphysalis longicornis* tick are crucial for ovarian development. J. Insect Physiol. 56, 1587-1598 (2010).
 - Aung, K.M., Boldbaatar, D., Liao, M., Nakao, S., Umemiya-Shirafuji, R., Tanaka, T., Fujisaki, K.: Identification and characterization of class B scavenger receptor CD36 from the hard tick, *Haemaphysalis longicornis*. Parasitol. Res. 108, 273-285 (2011).
 - Anisuzzaman, Islam, M.K., Abdul, M.A., Miyoshi, T., Hatta, T., Yamaji, K., Matsumoto, Y., Fujisaki, K., Tsuji, N.: Longistatin, a plasminogen activator, is key to the availability of blood meals for ixodid ticks. PLoS Pathogen 7, e1001312 (2011).