

A NOTE ON THE RELATION BETWEEN THE GRAPH OF
THE DEGREES OF THE GROUP CHARACTERS AND
NON-SIMPLICITY

著者	ATSUMI Tsuyoshi
journal or publication title	鹿児島大学理学部紀要. 数学・物理学・化学
volume	9
page range	39-41
別言語のタイトル	群の既約指標の次数と正規性について
URL	http://hdl.handle.net/10232/00003965

A NOTE ON THE RELATION BETWEEN THE GRAPH OF THE DEGREES OF THE GROUP CHARACTERS AND NON-SIMPLICITY

By

Tsuyoshi ATSUMI*

(Received September 27, 1976)

1. Introduction and Summary

Let G be a finite group, $D(G)$ the set of degrees of the irreducible complex non-principal characters of G . We introduce an ordering in $D(G)$ as follows: let a and b be two elements of $D(G)$. Then $a > b$ if and only if a divides b . Let k be the number of maximal elements in $D(G)$. Then G is k -headed. We form a graph $D(G)$ of G as follows: the points of $D(G)$ are the elements of $D(G)$. The (oriented) edge ab of $D(G)$ exists, where a and b are points of $D(G)$, if and only if $a > b$. Now we shall have the following conjecture.

Conjecture. *If $D(G)$ is a 2-headed graph then G is non-simple.*

In special cases the above problem and the related problems were solved by I.M. Isaacs and D.S. Passman in [2], [3], [4] and [5]. In this note we shall prove the following theorem.

Theorem. *Let G be a finite group with the following properties, the set of degrees of the irreducible complex characters of G is $\{1, m, n, k_1, k_2, \dots, k_i\}$ and $mn \mid k_i$ for all i . Then G is not a simple group.*

2. Proof of the theorem

Suppose the statement is false and let G be a counter example to the theorem. We can assume that $m < n$ and by a result of Thompson [6] $(m, n) = 1$. Let χ be an irreducible non-linear character of G with $\chi(1) = m$. Since G has the irreducible characters of degree n it follows from a theorem of Burnside and Brauer (see Satz 10.8 on p. 519 of [1]) that some power χ^r has an irreducible constituent of degree n . Choose r minimal with this property. Similarly χ^{s_i} has an irreducible constituent of degree k_i and s_i is minimal with this property.

Let $\phi_i \in \text{Irr}(G)$, $\phi_i(1) = k_i$ with ϕ_i a constituent of χ^{s_i} .

If there exists $i \in \mathbb{Z}$ such that $s_i < r$, then let the minimal number of $\{s_i \mid s_i < r\}$ be s_i . For some irreducible constituent ψ of χ^{s_i-1} we must have

* Department of Mathematics, Faculty of Science, Kagoshima University, Kagoshima, Japan.

$$0 \neq [\psi\chi, \phi_i]$$

and by the minimality of s_i we have $\psi(1)=m$ or $\psi(1)=1$.

Then $\psi(1)\chi(1) \leq m^2 < \phi_i(1) = k_i$. This is a contradiction.

So from now on we assume that for all i $s_i \geq r$. As above let $\phi \in \text{Irr}(G)$, $\phi(1)=n$ with ϕ a constituent of χ^r .

For some irreducible constituent ψ of χ^{r-1} we must have

$$0 \neq [\psi\chi, \phi] = \frac{1}{|G|} \sum_{x \in G} \psi(x)\chi(x)\overline{\phi(x)} = [\bar{\psi}, \chi\bar{\phi}]$$

and by the minimality of r we have $\psi(1)=m$. (The case that $\psi=1$ is impossible since then χ is irreducible of degree m .)

Thus $\chi\bar{\phi}$ has an irreducible constituent of degree m and has no linear constituent (in this case this is 1) since otherwise

$$0 \neq [\chi\bar{\phi}, 1] = [\bar{\phi}, \bar{\chi}],$$

contradicting $\bar{\phi}(1)=n > m = \bar{\chi}(1)$. Thus all irreducible constituents of $\chi\bar{\phi}$ have degree m , n or k_i and at least one has degree m . Let a be the number of constituents of degree m , b the number of those of degree n and c_i the number of those of degree k_i . We obtain $mn = am + bn + \sum_{i=1}^l c_i k_i$. Now $n|am$ and since $(m, n)=1$, we have $n|a$. However $a > 0$ and thus $a \geq n$. It follows that $a=n$, $b=0$, $c_i=0$ for all i . So every irreducible constituent of $\chi\bar{\phi}$ has degree m . We may write

$$\chi\bar{\phi} = \sum_{i=1}^n \theta$$

where the $\theta_i \in \text{Irr}(G)$ all have degree m and not necessarily all distinct. Suppose some θ_i is not χ . Then we have

$$0 = [\chi, \theta_i] = [1, \bar{\chi}\theta_i] \text{ and } \bar{\chi}\theta_i \text{ has not } 1.$$

However $0 \neq [\chi\bar{\phi}, \theta_i] = [\bar{\phi}, \bar{\chi}\theta_i]$ so $\bar{\chi}\theta_i$ has a constituent $\bar{\phi}$ of degree n . Let c be the number of the irreducible constituents of $\bar{\chi}\theta_i$ of degree m , d the number of degree n and e_i the number of degree k_i . Then as above we have

$$m = cm + dn + \sum_{i=1}^l e_i k_i.$$

Thus $m|d$ and $d > 0$ so $d \geq m$ and we have $m^2 \geq dn \geq mn$ which contradicts $n > m$. It follows that each θ_i is χ .

This yields

$$\chi\bar{\phi} = n\chi.$$

Since $\bar{\phi}$ is faithful. $\chi(x) = 0$ for $x \in G$, $x \neq 1$.

This yields $[\chi, 1] \neq 0$. This is a contradiction. So we complete the proof of the theorem.

Acknowledgment

The author is grateful to Prof. H. Nagao for suggesting the present form of the theorem in a letter to the author.

References

- [1] B. Huppert, *Endliche Gruppen 1*, Springer-Verlag, Berlin, 1967.
- [2] I.M. Isaacs, Groups having at most three irreducible character degrees, *Proc. Amer. Math. Soc.*, **21** (1969), 185–188.
- [3] I.M. Isaacs and D.S. Passman, Groups with representations of bounded degree, *Canad. J. Math.*, **16** (1964), 299–309.
- [4] ———, A characterization of groups in terms of the degrees of their characters, *Pacific J. Math.*, **15** (1965), 877–903.
- [5] ———, A characterization of groups in terms of the degrees of their characters, II, *Pacific J. Math.*, **24** (1968), 467–510.
- [6] J.G. Thompson, Normal p -complements and irreducible characters, *J. of Algebra*, **14** (1970), 129–134.