

Approximation of Fourier Transforms by Piecewise Linear Functions

著者	SUDA Katsuhiko, KAWAI Toru
journal or publication title	鹿児島大学理学部紀要. 数学・物理学・化学
volume	27
page range	13-16
別言語のタイトル	区分的線形関数によるフーリエ変換の近似
URL	http://hdl.handle.net/10232/00000501

Approximation of Fourier Transforms by Piecewise Linear Functions

Katsuhiko SUDA¹⁾ and Toru KAWAI¹⁾

(Received September 12, 1994)

Abstract

An approximation theorem of Fourier transform by piecewise linear functions is proved.

Key words: Fourier transform, piecewise linear function.

1. Introduction

We denote Banach spaces of all integrable functions and all regular complex finite measures on the real line by $L^1(\mathbf{R})$ and $M(\mathbf{R})$, respectively. For a function $f \in L^1(\mathbf{R})$, the Fourier transform is

$$\hat{f}(\xi) = \int_{-\infty}^{\infty} e^{-i\xi x} f(x) dx \quad (\xi \in \mathbf{R}),$$

and for a measure $\mu \in M(\mathbf{R})$, the Fourier-Stieltjes transform is

$$\hat{\mu}(\xi) = \int_{-\infty}^{\infty} e^{-i\xi x} d\mu(x) \quad (\xi \in \mathbf{R}).$$

We define

$$A(\mathbf{R}) = \{\hat{f}: f \in L^1(\mathbf{R})\}, \quad \|\hat{f}\|_{A(\mathbf{R})} = \|f\|_{L^1(\mathbf{R})}$$

and

$$B(\mathbf{R}) = \{\hat{\mu}: \mu \in M(\mathbf{R})\}, \quad \|\hat{\mu}\|_{B(\mathbf{R})} = \|\mu\|_{M(\mathbf{R})}.$$

¹⁾ Department of Mathematics, Faculty of Science, Kagoshima University, 1-21-35 Korimoto, Kagoshima 890, Japan

We denote Banach spaces of all integrable functions on one dimensional torus \mathbf{T} by $L^1(\mathbf{T})$. For a function $f \in L^1(\mathbf{T})$, the Fourier coefficients are

$$\hat{f}(n) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-inx} f(x) dx \text{ for integers } n.$$

We define

$$A(\mathbf{T}) = \left\{ \varphi: \sum_{n=-\infty}^{\infty} |\hat{\varphi}(n)| < \infty \right\}, \quad \|\varphi\|_{A(\mathbf{T})} = \sum_{n=-\infty}^{\infty} |\hat{\varphi}(n)|.$$

Herz [2] has proved that Cantor's ternary set is a spectral synthesis set. Kahane [3] has proved this result using an approximation of functions in $A(\mathbf{T})$ by piecewise linear functions: if $\varphi \in A(\mathbf{T})$ and each φ_n is a continuous function on \mathbf{T} to equal φ at the multiples of $2\pi/n$ and linear on the remainder, then $\varphi_n \in A(\mathbf{T})$ and $\|\varphi_n - \varphi\|_{A(\mathbf{T})} \rightarrow 0$ ($n \rightarrow \infty$) (See also Kahane and Salem [4] and Benedetto [1]). In this paper, we prove that a similar result holds for $A(\mathbf{R})$.

2. Approximation of Fourier transforms

For a continuous function φ on \mathbf{R} and a positive integer n , we define φ_n to be the function to equal φ at the points $2\pi k/n$, $k=0, \pm 1, \pm 2, \dots$, and linear on the remainder of \mathbf{R} . For a continuous function φ on \mathbf{T} , we define φ_n similarly.

Lemma. *If $\varphi \in B(\mathbf{R})$, then $\varphi_n \in B(\mathbf{R})$ and $\|\varphi_n\|_{B(\mathbf{R})} \leq \|\varphi\|_{B(\mathbf{R})}$ for all n .*

Proof. For all $x \in \mathbf{R}$, define the function e_x by $e_x(\xi) = e^{ix\xi}$ ($\xi \in \mathbf{R}$). We now show that if x is a rational number, then $(e_{-x})_n \in B(\mathbf{R})$ and $\|(e_{-x})_n\|_{B(\mathbf{R})} = 1$. Suppose that x is a rational number. Take a positive integer r and an integer s such that $x = s/r$. We see that $(e_{-s})_{rn}(\xi) = (e_{-x})_n(r\xi)$ for all $\xi \in \mathbf{R}$. Since e_{-s} is 2π -periodic, regarding e_{-s} as a function on \mathbf{T} , we have $e_{-s} \in A(\mathbf{T})$. It follows that $(e_{-s})_{rn} \in A(\mathbf{T})$ and $\|(e_{-s})_{rn}\|_{A(\mathbf{T})} = 1$ (See Benedetto [1, p.168]). Hence $(e_{-x})_n \in B(\mathbf{R})$ and $\|(e_{-x})_n\|_{B(\mathbf{R})} = 1$. Let $\xi_1, \dots, \xi_m \in \mathbf{R}$ and c_1, \dots, c_m be complex numbers such that $\|\sum_{k=1}^m c_k e_{-\xi_k}\|_{\infty} \leq 1$. To prove $\varphi_n \in B(\mathbf{R})$ and $\|\varphi_n\|_{B(\mathbf{R})} \leq \|\varphi\|_{B(\mathbf{R})}$, it suffices to show that

$$\left| \sum_{k=1}^m c_k \varphi_n(\xi_k) \right| \leq \|\varphi\|_{B(\mathbf{R})}.$$

Let $\varphi = \hat{\mu}$, $\mu \in M(\mathbf{R})$, and $\varepsilon > 0$. Take $\delta > 0$ so that $2(1 + \|\varphi\|_{B(\mathbf{R})})\delta \sum_{k=1}^m |c_k| < \varepsilon$. Since

$(e_{-x})_n(\xi_k)$ is uniformly continuous as a function of x for each k , there is a positive number η such that $|(e_{-x})_n(\xi_k) - (e_{-y})_n(\xi_k)| < \delta$ for all x, y and k such that $|x - y| < \eta$. Since $\mu \in M(\mathbf{R})$, there are $t_0, \dots, t_p \in \mathbf{R}$ so that $t_0 < \dots < t_p$, $t_i - t_{i-1} = \eta$ for all i , and $|\mu|((t_0, t_p]^c) < \delta$. Choose rational numbers x_1, \dots, x_p such that $x_i \in (t_{i-1}, t_i]$ ($i=1, \dots, p$). It follows that

$$\begin{aligned} \varphi_n(\xi) &= \sum_{j=-\infty}^{\infty} \varphi\left(\frac{2\pi j}{n}\right) \Delta\left(\xi - \frac{2\pi j}{n}\right) = \sum_{j=-\infty}^{\infty} \int_{-\infty}^{\infty} (e_{-x})\left(\frac{2\pi j}{n}\right) d\mu(x) \Delta\left(\xi - \frac{2\pi j}{n}\right) \\ &= \int_{-\infty}^{\infty} (e_{-x})_n(\xi) d\mu(x), \end{aligned}$$

where $\Delta(\xi) = \max\left(1 - \frac{n}{2\pi}|\xi|, 0\right)$ for $\xi \in \mathbf{R}$. This implies that

$$\begin{aligned} \left| \sum_{k=1}^m c_k \varphi_n(\xi_k) \right| &= \left| \sum_{k=1}^m c_k \int_{-\infty}^{\infty} (e_{-x})_n(\xi_k) d\mu(x) \right| \\ &\leq \left| \sum_{k=1}^m c_k \int_{(t_0, t_p]} (e_{-x})_n(\xi_k) d\mu(x) \right| + \left| \sum_{k=1}^m c_k \int_{(t_0, t_p]^c} (e_{-x})_n(\xi_k) d\mu(x) \right| \\ &\leq \left| \sum_{k=1}^m c_k \sum_{i=1}^p \int_{(t_{i-1}, t_i]} (e_{-x_i})_n(\xi_k) d\mu(x) \right| + \sum_{k=1}^m |c_k| \delta \|\varphi\|_{B(\mathbf{R})} \\ &\quad + \sum_{k=1}^m |c_k| |\mu|((t_0, t_p]^c) \\ &\leq \sum_{i=1}^p \int_{(t_{i-1}, t_i]} d|\mu|(x) \left| \sum_{k=1}^m c_k (e_{-x_i})_n(\xi_k) \right| + \varepsilon. \end{aligned}$$

Since $(e_{-x})_n \in B(\mathbf{R})$ and $\|(e_{-x})_n\|_{B(\mathbf{R})} = 1$ for every rational numbers x , it follows that

$$\left| \sum_{k=1}^m c_k (e_{-x_i})_n(\xi_k) \right| \leq 1 \quad (i=1, \dots, p).$$

Consequently, we have

$$\left| \sum_{k=1}^m c_k \varphi_n(\xi_k) \right| \leq \|\varphi\|_{B(\mathbf{R})} + \varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, this completes the proof of the lemma.

Theorem. If $\varphi \in A(\mathbf{R})$, then $\varphi_n \in A(\mathbf{R})$ for all n and $\|\varphi_n - \varphi\|_{A(\mathbf{R})} \rightarrow 0$ ($n \rightarrow \infty$).

Proof. Let V_m be de la Vallée Poussin's kernel; that is,

$$\tilde{V}_m(\xi) = \begin{cases} 1 & (|\xi| \leq m) \\ 2 - |\xi|/m & (m \leq |\xi| \leq 2m) \\ 0 & (2m \leq |\xi|) \end{cases} \quad (\xi \in \mathbf{R}).$$

By Lemma, we have $(\varphi \widehat{V}_m)_n \in B(\mathbf{R})$. Since each $(\varphi \widehat{V}_m)_n$ has compact support, this means that $(\varphi \widehat{V}_m)_n \in A(\mathbf{R})$. Since $\|\varphi - \varphi \widehat{V}_m\|_{A(\mathbf{R})} \rightarrow 0$ ($m \rightarrow \infty$), Lemma shows that $\{(\varphi \widehat{V}_m)_n\}_m$ is a Cauchy sequence in $A(\mathbf{R})$. Let g denote the limit of $(\varphi \widehat{V}_m)_n$ in $A(\mathbf{R})$. The value $\varphi_n(x)$ is equal to $g(x)$ since it is the limit of $(\varphi \widehat{V}_m)_n(x)$ for each $x \in \mathbf{R}$. Thus we have $\varphi_n \in A(\mathbf{R})$. To prove that $\|\varphi_n - \varphi\|_{A(\mathbf{R})} \rightarrow 0$ ($n \rightarrow \infty$), assume first that φ has compact support. Let E be the subset of $A(\mathbf{R})$ consisting of all functions h with compact support contained in $(-3\pi/4, 3\pi/4)$. Then there exists a linear mapping S of E into $A(\mathbf{T})$ and constant C such that $Sh = h$ on $(-\pi, \pi)$ and $\|h\|_{A(\mathbf{R})} \leq C \|Sh\|_{A(\mathbf{T})}$ for all $h \in E$ (See Rudin [5, p.56]). It follows that $\|h_n - h\|_{A(\mathbf{R})} \leq C \|Sh_n - Sh\|_{A(\mathbf{T})}$ for all $n \geq 8$ and all $h \in A(\mathbf{R})$ with compact support contained in $(-\pi/2, \pi/2)$. To show that $\|\varphi_n - \varphi\|_{A(\mathbf{R})} \rightarrow 0$ ($n \rightarrow \infty$), let $\varepsilon > 0$ and p be a positive integer such that the support of φ is contained in $(-\pi p/2, \pi p/2)$. Let T be a mapping defined by $Tf(\xi) = f(p\xi)$ for $f \in A(\mathbf{R})$ and $\xi \in \mathbf{R}$. Since $T\varphi \in A(\mathbf{R})$ and the support of $T\varphi$ is contained in $(-\pi/2, \pi/2)$, it follows that $ST\varphi \in A(\mathbf{T})$. Therefore there is a positive integer N such that $\|(ST\varphi)_{pn} - ST\varphi\|_{A(\mathbf{T})} < \varepsilon/C$ for every $n \geq N$. (See Benedetto [1, p.168]). For $n > \max(8/p, N)$, we have

$$\begin{aligned} \|\varphi_n - \varphi\|_{A(\mathbf{R})} &= \|T\varphi_n - T\varphi\|_{A(\mathbf{R})} = \|(T\varphi)_{pn} - T\varphi\|_{A(\mathbf{R})} \leq C \|S(T\varphi)_{pn} - ST\varphi\|_{A(\mathbf{T})} \\ &= C \|(ST\varphi)_{pn} - ST\varphi\|_{A(\mathbf{T})} < \varepsilon. \end{aligned}$$

In the general case, we may use the fact that the functions with compact supports are dense in $A(\mathbf{R})$. Let $\varepsilon > 0$, and let ψ be a function with compact support such that $\|\varphi - \psi\|_{A(\mathbf{R})} < \varepsilon/3$. By Lemma, we have $\|\varphi_n - \psi_n\|_{A(\mathbf{R})} \leq \|\varphi - \psi\|_{A(\mathbf{R})}$. Since ψ has compact support, there is a positive integer N such that $\|\psi_n - \psi\|_{A(\mathbf{R})} < \varepsilon/3$ for all $n > N$. It follows that

$$\|\varphi_n - \varphi\|_{A(\mathbf{R})} \leq \|\varphi_n - \psi_n\|_{A(\mathbf{R})} + \|\psi_n - \psi\|_{A(\mathbf{R})} + \|\psi - \varphi\|_{A(\mathbf{R})} < \varepsilon.$$

This completes the proof of the theorem.

References

- [1] J.J. Benedetto, Spectral Synthesis, Academic Press, 1975.
- [2] C.S. Herz, Spectral synthesis for the Cantor set, Proc. Nat. Acad. Sci. USA, 42 (1956), 42–43.
- [3] J.-P. Kahane, Séries de Fourier Absolument Convergentes, Springer-Verlag, 1970.
- [4] J.-P. Kahane and R. Salem, Ensembles Parfaits et Séries Trigonométriques, Hermann, 1963.
- [5] W. Rudin, Fourier Analysis on Groups, Interscience, 1962.