

## 平成26年退職予定者の研究業績

岩井純夫氏（生物生産学科 園芸生産学講座 蔬菜園芸学研究室）

### 学術論文等

- Kawashima, N., Tanabe, Y. and Iwai, S.: Similarities and differences in the primary structure of fraction I proteins in the genus *Nicotiana*. *Biochim Biophys Acta*. 371, 417-31 (1974)
- Kawashima, N., Tanabe, Y. and Iwai, S.: Origin of *Nicotiana tabacum* detected by primary structure of Fraction I protein. *Biochim. Biophys. Acta*, 365, 138-147 (1974)
- Imai H., Iwai S. and Yamada Y.: Comparative studies on photosynthesis of higher plants. IV. Further studies on the photosynthetic sugar formation pathway in C4-plants. *Soil. Sci. Plant Nutri.* 21, 13-19 (1975)
- Iwai, S. Tanabe, Y. and Kawashima, N.: Origin of sequence heterogeneity of the small subunit of fraction 1 protein from *Nicotiana tabacum* L. *Biochem Biophys Res Commun*, 73, 993-996 (1976)
- Iwai S. Kawashima N. and Matsuyama, S.: Effect of water stress on proline catabolism in tobacco leaves. *Phytochemistry* 18, 1155-1157 (1979)
- Iwai, S. Nagao, T. Nakata, T., Kawashima, N. and Matsuyama, S.: Expression of nuclear and chloroplastic genes coding for fraction-1 protein in somatic hybrids of *Nicotiana tabacum + rustica*. *Planta*, 147, 414-417 (1980)
- Iwai, S. Nagao, T. Nakata, T., Kawashima, N. and Matsuyama, S.: Detection of the *Nicotiana rustica* chloroplast genome coding for the large subunit of Fraction I protein in a somatic hybrid in which only the *N. tabacum* chloroplast genome appeared to have been expressed. *Planta*, 152, 478-480 (1981)
- Sano S., Kawashima N. and Iwai S.: Characterization of amino acid distribution in the floral organs of *Nicotiana tabacum*. *Agric. Biol. Chem.*, 48, 323-332 (1983)
- Matsuzaki T., Koiwai, A., Iwai, S. and Yamada, Y.: In vitro proliferation of stigma-like, style-like structures of *Nicotiana tabacum* and its fatty acid composition. *Plant Cell Physiol*. 25, 197-203 (1984)
- Iwai, S., Kishi, C., Nakata K. and Kubo, S.: Production of a hybrid of *Nicotiana repanda x N. tabacum* by ovule culture. *Plant Science*, 41, 175-178 (1985)
- Iwai, S., Kishi, C., Nakata, K., and Kawashima, N.: Production of *Nicotiana tabacum x N. acuminata* hybrid by ovule culture. *Plant Cell Reports*, 5, 403-404 (1986)
- Matsuzaki, T., Iwai, S. and Koiwai, A.: Effects of temperature on seed fatty acid composition in ovary culture of tobacco. *Agric. Biol. Chem.*, 52, 1283-1285 (1988)
- Hamaoka, Y., Fujita, Y. and Iwai, S.: Number of chloroplasts in haploids and diploids produced via anther culture in *Brassica campestris*. *Plant Tissue Culture Letters*, 8, 67-72 (1991)
- Hamaoka, Y., Fujita, Y. and Iwai, S.: Effects of temperature on the mode of pollen development in anther culture of *Brassica campestris* L. *Physiol. Plantarum.*, 82, 67-72 (1991)
- Katoh, N. and Iwai, S.: In vitro induction of stigma-like and style-like structure from receptacle tissue in *Nicotiana tabacum* L. *Plant Cell, Tissue and Organ Culture*, 26, 7-10 (1991)
- Katoh, N., Hagimori, M. and S. Iwai: Production of haploid plants of melon by pseudofertilized ovule culture. *Plant Tissue Culture Letters*, 10: 60-66 (1993)
- Katoh, N., and Iwai, S.: Induction of haploid plants from unpollinated ovules in *Nicotiana rustica*. *Plant Tissue Culture Letters*, 10, 123-129 (1993)
- Katoh, N., Hagimori, M. and Iwai, S.: Effects of various culture condition on developmental patterns of tobacco ovules. *Plant Tissue Culture Letters*, 10: 130-137 (1993)
- Katoh, N. and Iwai, S.: Effect of plant growth substances on the formation of carpelloid structures from tobacco ovules. *J. Plant Res.*, 106: 129-135 (1993)
- Sato, S., Hagimori, M., and Iwai, S.: Recovering vitrified carnation (*Dianthus caryophyllus* L.) shoots using Bacto-Peptone and its subfraction. *Plant Cell Reports*, 12: 370-374 (1993)
- Sato, S., Katoh, N., Iwai, S. and M. Hagimori: Establishment of reliable methods of in vitro pollen germination and pollen preservation of *Brassica rapa* (syn. *B. campestris*). *Euphytica* 103: 29-33 (1998)
- Katoh N. and Iwai, S.: Formation of style- and stigma-like structures by ovule culture in tobacco. *J. Jap. Soc. Horticultural Science* 67, 1157-1160 (1998)
- Sato, S., Katoh, N., Yoshida, H., Iwai, S. and Hagimori, M.: Production of doubled haploid plants of Carnation (*Dianthus caryophyllus* L.) by pseudofertilized ovule culture. *Scientia Horticulturae* 83, 301-310 (2000)

- Etoh T., Watanabe H. and Iwai, S.: RAPD variation of garlic clones in the center of origin and the westernmost area of distribution. Mem. Fac. Agri. Kagoshima Univ. 37, 21-27 (2000)
- Hong, C., Watanabe, H., Etoh, T. and Iwai, S.: Morphological and karyological comparison of garlic clones between the center of origin and the westernmost area of distribution. Mem. Fac. Agri. Kagoshima Univ., 36, 1-10 (2000)
- Hong, C., Watanabe, H., Etoh, T. and Iwai, S.: A search of pollen fertile clones in the Iberian garlic of RAPD markers. Mem. Fac. Agri. Kagoshima Univ., 36, 11-16 (2000)
- Hong, C., Etoh, T. and Iwai, S.: An attempt of crossbreeding in garlic. Mem. Fac. Agri. Kagoshima Univ., 36, 17-24 (2000)
- Sato, S., Katoh, N., Iwai, S. and Hagimori, M.: Effect of low temperature pretreatment of buds or inflorescence on isolated microspore culture in *Brassica rapa* (syn. *B. campestris*). Breeding Science, 52, 23-26 (2002)
- Iwai, S., N. Shimomura, A. Nakashima and T. Etoh: New fava bean guard cell signaling mutant impaired in ABA-induced stomatal closure. Plant Cell Physiol., 44, 212-219 (2003)
- Kariya, H., Mutsumoto, T., Etoh, T. and Iwai, S.: Analysis of recombination frequencies in male and female gametes in *Cucumis melo* L. Japan. Soc. HortSci., 74, 451-457 (2005)
- Sato, S., Katoh, N., Iwai, S. and Hagimori, M.: Frequency of spontaneous polyploidization of embryos regenerated from cultured anthers or microspores of *Brassica rapa* var *pekinensis* L. and *B. oleracea* var. *capitata*. Breeding Science, 55 : 99-102 (2005)
- Yan, J., Tsuichihara, N., Etoh, T. and Iwai, S.: Reactive oxygen species and nitric oxide are involved in ABA-inhibition of stomatal opening. Plant & Cell Environment, 30, 1320-1325 (2007)
- Yang J., Matsumoto Y., Etoh, T. and Iwai, S.: Nitric oxide (NO)-dependent and NO-independent signaling pathways act in ABA- inhibition of stomatal opening. Plant Signaling & Behavior 3, 131-132 (2008)
- Nakashima, M., Shigekuni, Y., Obi, T., Shiraishi, M., Miyamoto, A., Yamasaki, H., Etoh, T. and Iwai, S.: Nitric oxide-dependent hypotensive effects of wax gourd juice. J. Ethnopharmacol., 138, 404-407 (2011)
- Watanabe, Y., Iwai S., Ono Y., Hiradate S., Fujii Y. and Komai E.: Development of in vitro system for evaluation of allelopathic activities of *Asparagus* calluses. J. Japan. Soc. Hort. Sci., 82-88 (2011)
- Watanabe, Y., Iwai S., Ono Y., Hiradate S., Fujii Y. and Komai E. Assesment of allelopathic activities in female and male individuals of asparagus seedlings and regenerants. J. Japan. Soc. Hort. Sci., 80, 169-174 (2011)
- Sugiyama, Y., Uraji, M., Watanabe-Sugimoto, M., Okuma, E., S. Munemasa, S., Y. Nakamura, Y., Mori, I., Iwai, S. and Murata, Y.: FIA functions as an early signal component of abscisic acid signal cascade in *Vicia faba* guard cells. J. Exp. Bot., 63, 1357-1365 (2012)
- Joudoi, T., Y. Shichiri, N. Kamizono, T. Akaike, T. Sawa, J. Yoshitake, N. Yamada, S. Iwai: Nitrated cGMP modulates guard cell signaling in *Arabidopsis*. Plant Cell 25: 58-571 (2013)
- 池澤和広・福元伸一・遠城道雄・吉田理一郎・岩井純夫. ポット栽培における湛水処理がサトイモ‘大吉’ (*Clocasia esculenta* Schott cv. ‘Daikichi’) の生育と収量に及ぼす影響. 園学研. (Hort. Res. (Japan)), 13, 35-40 (2014)

徳永正雄氏 (生物資源化学科 生命機能化学講座 応用分子微生物学研究室)

学術論文等

- Tokunaga, M., Nakano, Y. and Kitaoka, S.: The GABA shunt in the callus derived from soybean cotyledon. *Agric. Biol. Chem.*, 40, 115-120 (1976)
- Tokunaga, M., Nakano, Y. and Kitaoka, S.: Separation and properties of the NAD-linked and NADP-linked isozymes of succinic semialdehyde dehydrogenase in *Euglena gracilis* z. *Biochim. Biophys. Acta*, 429, 55-62 (1976)
- Tokunaga, M., Nakano, Y. and Kitaoka, S.: Preparation of the physiologically intact mitochondria from *Euglena gracilis* z. *Agric. Biol. Chem.*, 40, 1435-1436 (1976)
- Tokunaga, M., Nakano, Y. and Kitaoka, S.: Some properties of physiologically intact mitochondria from *Euglena gracilis* z., *Bull. Univ. Osaka Pref., Ser. B* 29 17-23 (1977)
- Tokunaga, H., Tokunaga, M., Nakano, Y. and Kitaoka, S.: Some further properties of the partially purified gamma-aminobutyrate transaminase from *Bacillus cereus* strain K-22. *Bull. Univ. Osaka Pref., SerB* 29, 24-31 (1977)
- Tokunaga, M., Nakano, Y. and Kitaoka, S.: Subcellular localization of the GABA-shunt enzymes in *Euglena gracilis* z. *J. Protozool.*, 24, 471-473 (1979)
- Nakae, T., Ishii, J. and Tokunaga, M.: Subunit structure of functional porin oligomers that form permeability channels in the outer membrane of *Escherichia coli*. *J. Biol. Chem.*, 254, 1457-1461 (1979)
- Tokunaga, H., Tokunaga, M. and Nakae, T.: Characterization of porins from the outer membrane of *Salmonella typhimurium*. Chemical analysis. *Eur. J. Biochem.*, 95, 433-439 (1979)
- Tokunaga, M., Tokunaga, H., Okajima, Y. and Nakae, T.: Characterization of porins from the outer membrane of *Salmonella typhimurium*. 2. Physical properties of the functional oligomeric aggregates. *Eur. J. Biochem.*, 95, 441-448 (1979)
- Tokunaga, M., Tokunaga, H. and Nakae, T.: The outer membrane permeability of Gram-negative bacteria: Determination of the permeability rate in reconstituted vesicle membranes. *FEBS Lett.*, 106, 85-88 (1979)
- Tokunaga, H., Tokunaga, M. and Nakae, T.: Permeability properties of chemically modified porin trimers from *Escherichia coli*. *J. Biol. Chem.*, 256, 8024-8029 (1981)
- Tokunaga, M., Tokunaga, H. and Wu, H.C.: Post-translational modification and processing of *Escherichia coli* prolipoprotein in vitro. *Proc. Natl. Acad. Sci. USA*, 79, 2255-2259 (1982)
- Tokunaga, M., Loranger, J.M., Wolfe, P.B. and Wu, H.C.: Prolipoprotein signal peptidase in *Escherichia coli* is distinct from the M13 procoat protein signal peptidase. *J. Biol. Chem.*, 257, 9922-9925 (1982)
- Tokunaga, M., Loranger, J.M. and Wu, H.C.: Isolation and characterization of an *Escherichia coli* clone overproducing prolipoprotein signal peptidase. *J. Biol. Chem.*, 258, 12102-12105 (1983)
- Wu, H.C., Tokunaga, M., Tokunaga, H., Hayashi, S. and Giam, C.H.: Posttranslational modification and processing of membrane lipoproteins in bacteria. *J. Cell. Biochem.*, 22, 161-171 (1983)
- Tokunaga, M., Loranger, J.M. and Wu, H.C.: Prolipoprotein modification and processing enzymes in *Escherichia coli*. *J. Biol. Chem.*, 3825-3830 (1984)
- Tokunaga, M., Loranger, J.M. and Wu, H.C.: A distinct signal peptidase for prolipoprotein in *Escherichia coli*. *J. Cell. Biochem.*, 24, 113-120 (1984)
- Regue, M., Remenick, J., Tokunaga, M., Mackie, G. A. and Wu, H.C.: Mapping of the lipoprotein signal peptidase gene (*lsp*). *J. Bacteriol.*, 158, 632-635 (1984)
- Innis, M.A., Tokunaga, M., Williams, M., Loranger, J.M., Chang, S.Y., Chang, C., and Wu, H.C.: Nucleotide sequence of the *Escherichia coli* prolipoprotein signal peptidase (*lsp*) gene. *Proc. Natl. Acad. Sci. USA*, 81, 3708-3712 (1984)
- Tokunaga, M., Loranger, J.M., Chang, S.Y., Regue, M., Chang, S. and Wu, H.C.: Identification of prolipoprotein signal peptidase and genomic organization of the *lsp* gene in *Escherichia coli*. *J. Biol. Chem.*, 260, 5610-5615 (1985)
- Wu, H.C. and Tokunaga, M.: Biogenesis of lipoproteins in bacteria. *Current topics in microbiology and immunology*, 125, 127-157 (1986)
- 林滋・徳永正雄・神尾好是：細菌リポ蛋白質の生合成および分泌機構。蛋白質 核酸 酵素, 32, 41-53 (1987)
- Tokunaga, M., Wada, N. and Hishinuma, F.: Expression and identification of immunity determinants on linear DNA plasmids pGKL1 and pGKL2 in *Kluyveromyces lactis*. *Nucleic Acids Res.*, 15, 1031-1046 (1987)
- Tokunaga, M., Wada, N. and Hishinuma, F.: A novel yeast secretion vector utilizing secretion signal of killer toxin encoded on the yeast linear DNA plasmid pGKL1. *Biochem. Biophys. Res. Commun.*, 144, 613-619 (1987)
- Tokunaga, M., Wada, N. and Hishinuma, F.: A novel yeast secretion signal isolated from 28K killer precursor protein

- encoded on the linear DNA plasmid pGKL1. *Nucleic Acids Res.*, 16, 7499-7511 (1988)
- Tokunaga, M., Kawamura, A. and Hishinuma, F.: Expression of pGKL killer 28k subunit in *Saccharomyces cerevisiae*. *Nucleic Acids Res.*, 17, 3435-3446 (1989).
  - Kanaya, E., Higashizaki, T., Ozawa, F., Hirai, K., Nishizawa, M., Tokunaga, M., Tsukui, H., Hatanaka, H. and Hishinuma, F.: Synthesis and secretion of human nerve growth factor by *Saccharomyces cerevisiae*. *Gene*, 83, 65-74 (1989).
  - 徳永 正雄：酵母線状プラスミドの構造とその機能. *日本農芸化学会誌*, 64, 46-49 (1990).
  - Tokunaga, M., Kawamura, A., Kitada, K. and Hishinuma, F.: Secretion of killer toxin encoded on the linear DNA plasmid pGKL1 from *Saccharomyces cerevisiae*. *J. Biol. Chem.*, 265, 17274-17280 (1990).
  - 徳永 正雄, 河野 憲二：BiP蛋白の構造と機能. *Cell Science*, 8, 61-68 (1991).
  - Tokunaga, M., Kawamura, A., Omori, A. and Hishinuma, F.: Purification and determination of the NH<sub>2</sub>-terminal amino acid sequence of mouse-alpha-amylase secreted from *Saccharomuces cerevisiae*: correct processing of secretion signal from pGKL killer 28 kDa precursor protein. *Biochim. Biophys. Acta*, 1080, 135-137 (1991).
  - 徳永 正雄, 菱沼文男. シグナルペプチドの改変. *蛋白質核酸酵素*. 37, 223-234 (1992).
  - Tokunaga, M., Kawamura, A., Omori, A. and Hishinuma, F.: Characterization of mouse-alpha-amylase secreted from *Saccharomyces cerevisiae* by the pGKL 128kDa killer secretion signal. *Biosci. Biotech.and Biochem.*, 56, 155-156 (1992).
  - Tokunaga, M., Kawamura, A., Omori, A. and Hishinuma, F.: Prepro-structure of yeast pGKL 128 kDa killer secretion signal:processing of 128 kDa killer secretion signal-alpha-amylase fusion protein. *Eur. J. Biochem.*, 203, 415-423 (1992).
  - Tokunaga, M., Kawamura, A. and Kohono, K.: Purification and characterization of BiP/Kar2 protein from *Saccharomyces cerevisiae*. *J. Biol. Chem.*, 267, 17553-17559 (1992).
  - Tokunaga, M., Kawamura, A., Yoneyuu, S., Kishida, M. and Hishinuma, F.: Secretion of mouse a-amylase from fission yeast *Schizosaccharomyces pombe*: presence of chymostatin-sensitive protease activity in the culture medium. *Yeast*, 9, 379-387 (1993).
  - 徳永 正雄. 酵母における蛋白質輸送・分泌生産. *化学と生物*, 33, 427-436 (1995).
  - Kishida M., Tokunaga M., Katayose Y., Yajima H., Kawamura-Watabe A., Hishinuma F.: Isolation and genetic characterization of pGKL killer-insensitive mutants (*iki*) from *Saccharomyces cerevisiae*. *Biosci Biotechnol Biochem.*, 60(5), 798-801 (1996)
  - Yajima H, Tokunaga M, Nakayama-Murayama A, Hishinuma F.: Characterization of IKI1 and IKI3 genes conferring pGKL killer sensitivity on *Saccharomyces cerevisiae*. *Biosci Biotechnol Biochem.*, 61(4), 704-9 (1997)
  - Tokunaga M., Ishibashi M., Tatsuda D., Tokunaga H.: Secretion of mouse alpha-amylase from *Kluyveromyces lactis*. *Yeast.*, 13(8), 699-706 (1997)
  - Tokunaga M., Miyawaki H., Shiraishi Y., Tokunaga H.: Purification and characterization of a GroEL homologue from the moderately eubacterial halophile *Pseudomonas* sp. #43., *FEMS Microbiol Lett.*, 152(2), 321-326 (1997)
  - Tokunaga M., Matsuoka K., Tokunaga H.: Identification and NH<sub>2</sub>-terminal amino acid sequences of DnaK and groEL homologues in moderate eubacterial halophiles. *Biosci Biotechnol Biochem.*, 61(8), 1388-1390 (1997)
  - Tokunaga H., Yamakawa M., Mizukami M., Takagi H., Tokunaga M.: Molecular cloning of the dnaK locus, and purification and characterization of a DnaK protein from *Bacillus brevis* HPD31. *Biochim Biophys Acta.* 1387(1-2), 65-79 (1998)
  - Tokunaga M., Kato S., Kawamura-Watabe A., Tanaka R., Tokunaga H.: Characterization of deletion mutations in the carboxy-terminal peptide-binding domain of the Kar2 protein in *Saccharomyces cerevisiae*. *Yeast*, 14(14), 1285-1295 (1998)
  - Tanaka R., Tokunaga H., Hara S., Arakawa T., Tokunaga M.: Expression and purification of cytokine receptor homology domain of human granulocyte-colony stimulating factor receptor in *Escherichia coli*. *Biosci Biotechnol Biochem.*, 62(9), 1809-1811 (1998)
  - Tokunaga H., Hara S., Arakawa T., Ishibashi M., Gupta RS, Tokunaga M.: Identification and partial purification of DnaK homologue from extremely halophilic archaeobacteria, *Halobacterium cutirubrum*. *J Protein Chem.*, 18(8), 837-844 (1999)
  - Tanaka R., Ishibashi M., Tokunaga H., Tokunaga M.: Secretion of hen egg white lysozyme from *Kluyveromyces lactis*. *Biosci Biotechnol Biochem.*, 64(12), 2716-2718 (2000)
  - Tatsuda D., Arimura H., Tokunaga H., Ishibashi M., Arakawa T., Tokunaga M.: Expression and purification of cytokine receptor homology domain of human granulocyte-colony-stimulating factor receptor fusion protein in *Escherichia coli*. *Protein Expr Purif.*, 21(1), 87-91 (2001)

- Ishibashi M., Tokunaga H., Arakawa T., Tokunaga M.: Expression, purification, and characterization of the active immunoglobulin-like domain of human granulocyte-colony-stimulating factor receptor in *Escherichia coli*. *Protein Expr Purif.*, 21(2), 317-322 (2001)
- Ishibashi M., Tokunaga H., Hiratsuka K., Yonezawa Y., Tsurumaru H., Arakawa T., Tokunaga M.: NaCl-activated nucleoside diphosphate kinase from extremely halophilic archaeon, *Halobacterium salinarum*, maintains native conformation without salt. *FEBS Lett.* 493(2-3), 134-138 (2001)
- Kato S., Ishibashi M., Tatsuda D., Tokunaga H. and Tokunaga M.: Efficient expression, purification and characterization of mouse salivary alpha-amylase secreted from methylotrophic yeast, *Pichia pastoris*. *Yeast*, 18(7), 643-655 (2001)
- Tokunaga M., Shiraishi Y., Odachi M., Mizukami M., Tokunaga H., Philo JS, Arakawa T., Ishibashi M., Tanaka R., Takagi H.: Molecular cloning of groESL locus, and purification and characterization of chaperonins, GroEL and GroES, from *Bacillus brevis*. *Biosci Biotechnol Biochem.*, 65(6), 1379-1387 (2001)
- Yonezawa Y, Tokunaga H, Ishibashi M, Tokunaga M.: Characterization of nucleoside diphosphate kinase from moderately halophilic eubacteria. *Biosci Biotechnol Biochem.*, 65(10), 2343-2346 (2001)
- Ishibashi M., Arakawa T., Philo JS, Sakashita K., Yonezawa Y., Tokunaga H., Tokunaga M.: Secondary and quaternary structural transition of the halophilic archaeon nucleoside diphosphate kinase under high- and low-salt conditions. *FEMS Microbiol Lett.* 216(2), 235-241 (2002)
- Ishibashi M., Arakawa T. and Tokunaga M.: Salting-in effects offset MgCl<sub>2</sub>-induced refolding of nucleoside diphosphate kinase. *Protein Pept Lett.*, 10(6), 575-580 (2003)
- Yonezawa Y., Tokunaga H., Ishibashi M., Taura S., Tokunaga M.: Cloning, expression, and efficient purification in *Escherichia coli* of a halophilic nucleoside diphosphate kinase from the moderate halophile *Halomonas* sp. #593. *Protein Expr Purif.*, 27(1), 128-133 (2003)
- Tanaka R., Mizukami M., Ishibashi M., Tokunaga H. and Tokunaga M.: Cloning and expression of the ccdA-associated thiol-disulfide oxidoreductase (catA) gene from *Brevibacillus choshinensis*: stimulation of human epidermal growth factor production. *J Biotechnol.*, 103(1), 1-10 (2003)
- Ishibashi M., Sakashita K., Tokunaga H., Arakawa T. and Tokunaga M.: Activation of halophilic nucleoside diphosphate kinase by a non-ionic osmolyte, trimethylamine N-oxide. *J Protein Chem.*, 22(4), 345-351 (2003)
- Arakawa T. and Tokunaga M.: Electrostatic and hydrophobic interactions play a major role in the stability and refolding of halophilic proteins. *Protein Pept Lett.*, 11(2), 125-132 (2004)
- Mine S., Koshiba T., Honjo E., Okamoto T., Tamada T., Maeda Y., Matsukura Y., Horie A., Ishibashi M., Sato M., Azuma M., Tokunaga M., Nitta K and Kuroki R.: Thermodynamic analysis of the activation mechanism of the GCSF receptor induced by ligand binding. *Biochemistry.* 43(9), 2458-2464 (2004)
- Ishibashi M., Arakawa T., Tokunaga M.: Facilitated folding and subunit assembly in *Escherichia coli* and *in vitro* of nucleoside diphosphate kinase from extremely halophilic archaeon conferred by amino-terminal extension containing hexa-His-tag. *FEBS Lett.*, 570(1-3), 87-92 (2004)
- Tokunaga H., Ishibashi M., Arakawa T., Tokunaga M.: Highly efficient renaturation of beta-lactamase isolated from moderately halophilic bacteria. *FEBS Lett.*, 558(1-3), 7-12 (2004)
- Tokunaga H., Mitsuo K., Kamekura M. and Tokunaga M.: Major outer membrane proteins in moderately halophilic eubacteria of genera *Chromohalobacter* and *Halomonas*. *J Basic Microbiol.*, 44(3), 232-240 (2004)
- Tokunaga H., Mitsuo K., Ichinose S., Omori A., Ventosa A., Nakae T., Tokunaga M.: Salt-inducible multidrug efflux pump protein in the moderately halophilic bacterium *Chromohalobacter* sp. *Appl Environ Microbiol.*, 70(8), 4424-4431 (2004)
- Tanaka R., Araki Y, Mizukami M., Miyauchi A., Ishibashi M., Tokunaga H., Tokunaga M.: Expression and purification of the *Bacillus subtilis* thioredoxin superfamily protein YkvV. *Biosci Biotechnol Biochem.*, 68(8), 1801-1804 (2004)
- Tanaka R., Kosugi K., Mizukami M., Ishibashi M., Tokunaga H., Tokunaga M.: Expression and purification of thioredoxin (TrxA) and thioredoxin reductase (TrxB) from *Brevibacillus choshinensis*. *Protein Expr Purif.*, 37(2), 385-391 (2004)
- Iwata T., Tanaka R., Suetsugu M., Ishibashi M., Tokunaga H., Kikuchi M., Tokunaga M.: Efficient secretion of human lysozyme from the yeast, *Kluyveromyces lactis*. *Biotechnol Lett.*, 26(23), 1803-1808 (2004)
- Ishibashi M., Tsumoto K., Ejima D., Arakawa T. and Tokunaga M.: Characterization of arginine as a solvent additive: a halophilic enzyme as a model protein. *Protein Pept Lett.*, 12(7), 649-653 (2005)
- Ishibashi M., Tsumoto K., Tokunaga M., Ejima D., Kita Y., Arakawa T.: Is arginine a protein-denaturant? *Protein Expr Purif.*, 42(1), 1-6 (2005)

- Ishibashi M., Yamashita S. and Tokunaga M.: Characterization of halophilic alkaline phosphatase from *Halomonas* sp. 593, a moderately halophilic bacterium. *Biosci Biotechnol Biochem.*, 69(6), 1213-1216 (2005)
- Tokunaga M., Arakawa T., Ishibashi M. and Tokunaga H.: Familiar extremophiles, halophiles, and halophilic enzymes. *Seikagaku*, 77(4), 320-331 (2005)
- Nagayoshi C., Ishibashi M., Kita Y., Matsuoka M., Nishimoto I. and Tokunaga M.: Expression, refolding and characterization of human brain serine racemase in *Escherichia coli* with N-terminal His-tag. *Protein Pept Lett.*, 12(5), 487-490 (2005)
- Honjo E., Tamada T., Maeda Y., Koshiha T., Matsukura Y., Okamoto T., Ishibashi M., Tokunaga M. and Kuroki R.: Crystallization of a 2:2 complex of granulocyte-colony stimulating factor (GCSF) with the ligand-binding region of the GCSF receptor. *Acta Crystallogr Sect F Struct Biol Cryst Commun.*, 61(Pt 8), 7887-7890 (2005)
- Ishibashi M., Tsumoto K., Ejima D., Arakawa T. and Tokunaga M.: Characterization of arginine as a solvent additive: a halophilic enzyme as a model protein. *Protein Pept Lett.*, 12(7), 649-653 (2005)
- Besir H., Zeth K., Bracher A., Heider U., Ishibashi M., Tokunaga M. and Oesterhelt D.: Structure of a halophilic nucleoside diphosphate kinase from *Halobacterium salinarum*. *FEBS Lett.*, 579(29), 6595-6600 (2005)
- Tokunaga H., Arakawa T., Fukada H., Tokunaga M.: Opposing effects of NaCl on reversibility and thermal stability of halophilic beta-lactamase from a moderate halophile, *Chromohalobacter* sp. 560. *Biophys Chem.*, 119(3), 316-320 (2006)
- Tokunaga H., Oda Y., Yonezawa Y., Arakawa T., Tokunaga M.: Contribution of halophilic nucleoside diphosphate kinase sequence to the heat stability of chimeric molecule. *Protein Pept Lett.*, 13(5), 525-530 (2006)
- Tamada T., Honjo E., Maeda Y., Okamoto T., Ishibashi M., Tokunaga M. and Kuroki R.: Homodimeric cross-over structure of the human granulocyte colony-stimulating factor (GCSF) receptor signaling complex. *Proc Natl Acad Sci U S A.*, 103(9), 3135-3140 (2006)
- Nagayoshi C., Tokunaga H., Hayashi A., Harazono H., Hamasaki K., Ando A. and Tokunaga M.: Efficient expression of haloarchaeal nucleoside diphosphate kinase via strong porin promoter in moderately halophilic bacteria. *Protein Pept Lett.*, 13(6), 611-615 (2006)
- Arakawa T., Tsumoto K., Ejima D., Kita Y., Yonezawa Y., Tokunaga M.: Induced binding of proteins by ammonium sulfate in affinity and ion-exchange column chromatography. *J Biochem Biophys Methods.*, 70(3), 493-498 (2007)
- Yonezawa Y., Izutsu K., Tokunaga H., Maeda H., Arakawa T. and Tokunaga M.: Dimeric structure of nucleoside diphosphate kinase from moderately halophilic bacterium: contrast to the tetrameric *Pseudomonas* counterpart. *FEMS Microbiol Lett.*, 268(1), 52-58 (2007)
- Sánchez-Porro C., Tokunaga H., Tokunaga M. and Ventosa A.: *Chromohalobacter japonicus* sp. nov., a moderately halophilic bacterium isolated from a Japanese salty food. *Int J Syst Evol Microbiol.*, 57(Pt 10), 2262-2266 (2007)
- Arakawa T., Ejima D., Tsumoto K., Ishibashi M. and Tokunaga M.: Improved performance of column chromatography by arginine: dye-affinity chromatography. *Protein Expr Purif.*, 52(2), 410-414 (2007)
- Ishibashi M., Tatsuda S., Izutsu K., Kumeda K., Arakawa T., Tokunaga M.: A single Gly114Arg mutation stabilizes the hexameric subunit assembly and changes the substrate specificity of halo-archaeal nucleoside diphosphate kinase. *FEBS Lett.*, 581(21), 4073-4079 (2007)
- Tokunaga H., Ishibashi M., Arisaka F., Arai S., Kuroki R., Arakawa T. and Tokunaga M.: Residue 134 determines the dimer-tetramer assembly of nucleoside diphosphate kinase from moderately halophilic bacteria. *FEBS Lett.*, 582(7), 1049-1054 (2008)
- Tokunaga H., Arakawa T., Tokunaga M.: Engineering of halophilic enzymes: two acidic amino acid residues at the carboxy-terminal region confer halophilic characteristics to *Halomonas* and *Pseudomonas* nucleoside diphosphate kinases. *Protein Sci.*, 17(9), 1603-1610 (2008)
- Tokunaga M., Tokunaga H., Ishibashi M. and Arakawa T.: Halophilic enzymes: negative charges determine halophilicity. *Seikagaku.*, 81(5), 401-406 (2009)
- Nagayoshi C., Ishibashi M. and Tokunaga M.: Purification and characterization of human brain serine racemase expressed in moderately halophilic bacteria. *Protein Pept Lett.* 16(2), 201-206 (2009)
- Ishibashi M., Iwasa T., Kumeda K., Arakawa T., Tokunaga M.: Effects of mutations at Gly114 on the stability and refolding of Haloarchaeal nucleoside diphosphate kinase in low salt solution. *Int J Biol Macromol.*, 44(4), 361-364 (2009)
- Ishibashi M. and Tokunaga M.: Molecular mechanism of halophilic enzyme-- nucleoside diphosphate kinase from extreme halophile. *Seikagaku.*, 81(12), 1080-1086 (2009)
- Tokunaga H., Saito S., Sakai K., Yamaguchi R., Katsuyama I., Arakawa T., Onozaki K., Arakawa T. and Tokunaga M.: Halophilic beta-lactamase as a new solubility- and folding-enhancing tag protein: production of native human interleukin

- alpha and human neutrophil alpha-defensin. *Appl Microbiol Biotechnol.*, 86(2), 649-658 (2010)
- Tokunaga M., Arakawa T. and Tokunaga H.: Recombinant expression in moderate halophiles. *Curr Pharm Biotechnol.*, 11(3), 259-266 (2010)
  - Hirano A., Tokunaga H., Tokunaga M., Arakawa T. and Shiraki K.: The solubility of nucleobases in aqueous arginine solutions. *Arch Biochem Biophys.*, 497(1-2), 90-96 (2010)
  - Tokunaga H., Arakawa T. and Tokunaga M.: Novel soluble expression technologies derived from unique properties of halophilic proteins. *Appl Microbiol Biotechnol.*, 88(6), 1223-1231 (2010)
  - Arakawa T., Tokunaga H., Yamaguchi R. and Tokunaga M.: High solubility supports efficient refolding of thermally unfolded  $\beta$ -lactamase. *Int J Biol Macromol.*, 47(5), 706-709 (2010)
  - Yamaguchi R., Tokunaga H., Ishibashi M., Arakawa T. and Tokunaga M.: Salt-dependent thermo-reversible  $\alpha$ -amylase: cloning and characterization of halophilic  $\alpha$ -amylase from moderately halophilic bacterium, *Kocuria varians*. *Appl Microbiol Biotechnol.* 89(3), 673-684 (2011)
  - Ishibashi M., Oda K., Arakawa T. and Tokunaga M.: Cloning, expression, purification and activation by Na ion of halophilic alkaline phosphatase from moderate halophile *Halomonas* sp. 593. *Protein Expr Purif.*, 76(1), 97-102 (2011)
  - Ishibashi M., Ida K., Tatsuda S., Arakawa T. and Tokunaga M.: Interaction of hexa-His tag with acidic amino acids results in facilitated refolding of halophilic nucleoside diphosphate kinase. *Int J Biol Macromol.*, 49(4), 778-783 (2011)
  - Rajan RS., Tsumoto K., Tokunaga M., Tokunaga H., Kita Y. and Arakawa T.: Chemical and pharmacological chaperones: application for recombinant protein production and protein folding diseases. *Curr Med Chem.*, 18(1), 1-15 (2011)
  - Yamaguchi R., Arakawa T., Tokunaga H., Ishibashi M. and Tokunaga M.: Distinct characteristics of single starch-binding domain SBD1 derived from tandem domains SBD1-SBD2 of halophilic *Kocuria varians* alpha-amylase. *Protein J.*, 31(3), 250-258 (2012)
  - Yamaguchi R., Inoue Y., Tokunaga H., Ishibashi M., Arakawa T., Sumitani J., Kawaguchi T. and Tokunaga M.: Halophilic characterization of starch-binding domain from *Kocuria varians*  $\alpha$ -amylase. *Int J Biol Macromol.*, 50(1), 95-102 (2012)
  - Yamaguchi R., Arakawa T., Tokunaga H., Ishibashi M. and Tokunaga M.: Halophilic properties of metal binding protein characterized by high histidine content from *Chromohalobacter salexigens* DSM3043. *Protein J.*, 31(2), 175-183 (2012)
  - Arai S., Yonezawa Y., Okazaki N., Matsumoto F., Tamada T., Tokunaga H., Ishibashi M., Blaber M., Tokunaga M. and Kuroki R.: A structural mechanism for dimeric to tetrameric oligomer conversion in *Halomonas* sp. nucleoside diphosphate kinase. *Protein Sci.*, 21(4), 498-510 (2012)
  - Yamaguchi R., Arakawa T., Tokunaga H., Ishibashi M. and Tokunaga M.: Effects of salt and ligand concentrations on the thermal unfolding and refolding of halophilic starch-binding domain from *Kocuria varians*  $\alpha$ -amylase. *Protein Pept Lett.*, 19(3), 326-332 (2012)
  - Ishibashi M., Uchino M., Arai S., Kuroki R., Arakawa T. and Tokunaga M.: Reduction of salt-requirement of halophilic nucleoside diphosphate kinase by engineering S-S bond. *Arch Biochem Biophys.*, 525(1), 47-52 (2012)
  - Tokunaga H., Arakawa T. and Tokunaga M.: Surface acidic amino acid of *Pseudomonas/Halomonas* chimeric nucleoside diphosphate kinase leads effective recovery from heat-denaturation. *Protein Pept Lett.*, 20(7), 836-841 (2013)
  - Yamaguchi R., Ishibashi M., Tokunaga H., Arakawa T. and Tokunaga M.: Structure of starch binding domains of halophilic alpha-amylase at low pH. *Protein Pept Lett.*, 20(7), 755-760 (2013).
  - Onishi H., Mizukami M., Hanagata H., Tokunaga M., Arakawa T. and Miyauchi A.: Efficient production of anti-fluorescein and anti-lysozyme as single-chain anti-body fragments (scFv) by *Brevibacillus* expression system. *Protein Expr Purif.*, 91(2), 184-191 (2013)
  - Tokunaga H., Furukawa M., Arakawa T. and Tokunaga M.: Channel forming outer membrane porin protein in halophile: expressed as a soluble form in *Escherichia coli*. *Int J Biol Macromol.*, 54, 44-50 (2013)
  - Tokunaga M., Mizukami M., Yamasaki K., Tokunaga H., Onishi H., Hanagata H., Ishibashi M., Miyauchi A., Tsumoto K. and Arakawa T.: Secretory production of single-chain antibody (scFv) in *Brevibacillus choshinensis* using novel fusion partner. *Appl Microbiol Biotechnol.*, 97(19), 8569-8580 (2013)
  - Ishibashi M., Hayashi T., Yoshida C. and Tokunaga M.: Increase of salt dependence of halophilic nucleoside diphosphate kinase caused by a single amino acid substitution. *Extremophiles.*, 17(4), 585-591 (2013).
  - Tokunaga Y., Matsumoto M., Tokunaga M., Arakawa T. and Sugimoto Y.: Amyloid fibril formation in vitro from halophilic metal binding protein: Its high solubility and reversibility minimized formation of amorphous protein aggregations. *Protein Sci.*, 22(11), 1582-1591 (2013)

## 著 書

- 酒井 明, 徳永 正雄, 菱沼 文男: 酵母における蛋白質の分泌生産. 日本生化学会編 新生化学実験講座 第1巻, p.213-233 東京化学同人, 東京 (1992).
- 徳永 正雄: 酵母ホスト-ベクター系. 日本生化学会編 新生化学実験講座. 第17巻, p.404-408, 東京化学同人, 東京 (1992).
- 徳永 正雄: *In vitro*蛋白質合成系-酵母. 日本生化学会編 新生化学実験講座. 第17巻, p.314-316, 東京化学同人, 東京 (1992).
- 徳永 正雄: 異種蛋白質の分泌系-ERにおけるQuality control mechanism. 21世紀に向かう酵母研究, p.25-35, 学会出版センター, 東京 (1994).
- 徳永 正雄: 酵母による蛋白質の分泌生産. 生物化学実験法37, p.143-165, 学会出版センター, 東京 (1995).



## 高木 東氏 (生物環境学科 生産環境工学講座 農地工学研究室)

## 学術論文等

- 矢橋辰吾・高木 東・中野政詩：浪岡東部丘陵地における土壌侵食－斜面の侵食に関する研究（Ⅰ）－．農業土木学会論文集，76，31-37（1978）
- 高木 東・矢橋辰吾・中野政詩：ガリ網の規則性と発達－斜面の侵食に関する研究（Ⅱ）－．農業土木学会論文集，77，47-52（1978）
- 高木 東・矢橋辰吾・中野政詩：ガリの縦横断形特性とガリ網の発達－斜面の侵食に関する研究（Ⅲ）－．農業土木学会論文集，90，17-25（1980）
- 高木 東・中野政詩：農地斜面のガリ網の形成・発達に関する確率的シミュレーション．農業土木学会論文集，98，22-30（1982）
- Nakano, M., Takagi, A. and Haraguchi, N.: Stochastic simulation of gully networks on eroded land. Soil Conservation Society of America, 178-187（1985）
- Nakano, M., Hagiwara, H., Takagi, A. and Shirai, K.: Computer Analysis on Effects of Land Parcel Shape upon Rill Network Formation. Irrigation Engineering and Rural Planning, 10, 34-39（1986）
- 高木 東・中野政詩：雨裂からの土砂流出に関する解析．農業土木学会論文集，126，51-58（1986）
- 小林裕志・堤 聰・佐藤幸一・高木 東・鈴木創三・杉浦俊弘：土・水保全に有効な採草播種床造成法と流去水処理法－改良山成畑工法について（Ⅳ）－．農業土木学会誌，55(2)，23-30（1987）
- 小林裕志・堤 聰・佐藤幸一・高木 東・鈴木創三・杉浦俊弘：採草畑造成における計画・施工例－改良山成畑工法について（Ⅲ）－．農業土木学会誌，54(10)，11-18（1986）
- 高木 東・堤 聰・中野政詩：リルにおける侵食量と流出土砂量の予測．農業土木学会論文集，141，89-98（1989）
- 堤 聰・大久保博・嶋 栄吉・高木 東：山地畜産における渓流取水の実態．農業土木学会誌，58(7)，29-34（1990）
- 大久保博・堤 聰・嶋 栄吉・江崎 要：畜産用水の需要変動と調整施設容量．農業土木学会誌，58(10)，29-34（1990）
- 友正達美・高木 東・中尾誠司：土壌流亡量式USLEにおける降雨係数の近似計算法．農業土木学会誌，58(12)，25-28（1990）
- 中尾誠司・高木 東・友正達美・細川雅敏：開発農用地における承・集水路周辺部の侵食防止工法．農業土木学会誌，62(4)，33-38（1994）
- 高木 東・中尾誠司・友正達美：流砂モデルによるリル侵食量および流砂量の解析．農業土木学会論文集，181，11-21（1996）
- 中尾誠司・高木 東・友正達美・細川雅敏：傾斜ウネ立て圃場からの土砂流出量予測モデルの提案とその検証，184，1-12（1996）
- 塩野隆弘・高木 東・小倉 力・上村健一郎：畝部の土壌侵食に対するキャベツの植生の影響度．農業土木学会論文集，204，13-20（1999）
- 高木 東・堤 聰・増本隆夫・塩野隆弘：短期間における農地のリル形成およびリル網の発達．農業土木学会論文集，207，1-9（2000）
- 吉村亜希子・石田憲治・高木 東：行方・鹿島台地における低水流出の逓減特性について．農業工学研究所技報，198，51-60（2000）
- 中尾誠司・高木 東・細川雅敏：傾斜畑における土壌侵食防止のための保全管理とその対策．農業土木学会誌，70(7)，35-38（2002）
- Takagi, A., Nakao, S. and Tomosho, T.: Analysis of Soil Erosion and Transport along a Rill Channel using a Sediment Transport Model. JARQ, 35(4), 245-254（2001）
- 中尾誠司・高木 東・松田誠祐：長大ウネ立て圃場に対する土砂流出量予測モデルの適用性．農業土木学会論文集，226，89-95（2003）
- 中尾誠司・高木 東・松田誠祐：土壌侵食モデルにおける受食性パラメータの実験的決定法．農業土木学会論文集，226，97-104（2003）
- 高木 東・増本隆夫・堤 聰：農地におけるリル発生の限界斜面長の予測．農業土木学会論文集，228，25-32（2003）
- 塩野隆弘・奥島修二・高木 東・福本昌人：黒ボク土畝立て圃場の土壌侵食に対するキャベツ栽培の影響．農業土木学会論文集，230，1-9（2004）
- 増本隆夫・久保田富次郎・松田 周・高木 東：水田農業地帯の水資源が持つ生態環境維持機能の評価法．農業工学研究所技報，202，81-90（2004）
- 藤井秀人・石田憲治・中 達雄・高木 東：水田灌漑農業の国際比較と水利用の持続性評価手法．農業土木学会誌，

72(2), 39-44 (2004)

- 伊藤祐二・舛井和朗・中川 啓・高木 東：池田湖水位の推定モデル－水資源としての湖の水収支に関する研究－．農業土木学会論文集, 243, 65-72 (2006)
- Naka, T., Ishida, K., Fujii, H. and Takagi, A. : Evaluation Indicators on Water-use Sustainability in Irrigated Paddy Farming. JARQ, 41(3), 233-245 (2007)
- Takagi, A., Shiono, T. and Nakano, M.: Rill Link Characteristics and Statistical Laws of Geomorphology of Rill Networks in Fields. Trans. of JSIDRE, 253, 1-9 (2008)
- Takagi, A., Shiono, T. and Nakano, M.: A Simplified Method to Predict Sediment Yield from a Developed Rill Network Using Rill Link Characteristics. Trans. of JSIDRE, 253, 11-20 (2008)
- 高木 東・小川茂男・平 瑞樹：農地のリル網の平均リルリンク長と1次元的地形則．農業農村工学会論文集, 267, 59-67 (2010)
- 高木 東・中川 啓・肥山浩樹：農地におけるリル網の集水域に関する地形特性量．地形統計則およびリル密度の予測式：農業農村工学会論文集, 272, 31-42 (2011)
- 肥山浩樹・舛井和朗・中川 啓・高木 東：畑地帯からの降雨流出特性に関する研究－鹿児島県徳之島および大隅圃場の事例－．鹿大農学学術報告, 61, 29-34 (2011)
- 高木 東・舛井和朗・肥山浩樹・伊藤祐二：南九州畑地集水域における降雨流出特性－鹿児島県大隅および徳之島地域の事例－．農業農村工学会論文集, 288, 113-120 (2013)

#### 著 書

- 高木 東：山路永司・塩沢 昌編集 農地環境工学 第6章 農地の保全と防災 1. 水食とその対策 2. 風食とその対策．文永堂, 東京 (2008)
- 高木 東：日本草地学会編 草地科学実験・調査法. 17.4 土壌浸食. 全国農村教育協会, 東京 (2004)