

## ON HAEFLIGER'S OBSTRUCTIONS TO EMBEDDINGS AND TRANSFER MAPS

Dedicated to the memory of Professor Katsuo Kawakubo

YOSHIYUKI KURAMOTO and TSUTOMU YASUI

(Received May 17, 2001)

### 1. Introduction and statement of results

Throughout this article,  $n$ -manifolds mean compact differentiable (or topological) manifolds of dimension  $n$ . The (co-)homology is understood to have  $\mathbf{Z}_2$  for coefficients.

For a manifold  $V$ , we denote by  $w(V)$  and  $\bar{w}(V)(= w(V)^{-1})$ , the total Stiefel-Whitney class and the total normal Stiefel-Whitney class of  $V$ , respectively. Furthermore, we denote by  $U_V \in H^{\dim V}(V \times V)$  the  $\mathbf{Z}_2$ -Thom class (or  $\mathbf{Z}_2$ -diagonal cohomology class) of  $V$  [10, p. 125]. For a (continuous) map  $f: M^n \rightarrow N^{n+k}$  between closed manifolds  $M$  and  $N$ , we define the total Stiefel-Whitney class  $w(f) = \sum_{i \geq 0} w_i(f)$  by the equation

$$w(f) = \bar{w}(M) f^* w(N).$$

For a map  $f: M^n \rightarrow N^{n+k}$ , the transfer map (or Umkehr homomorphism)  $f_!: H^i(M) \rightarrow H^{i+k}(N)$  is defined by the commutative diagram below:

$$\begin{array}{ccc} H^i(M) & \xrightarrow{f^!} & H^{i+k}(N) \\ \cong \downarrow \cap [M] & & \cong \downarrow \cap [N] \\ H_{n-i}(M) & \xrightarrow{f_*} & H_{n-i}(N). \end{array}$$

Here  $[V] \in H_{\dim V}(V)$  denotes the fundamental homology class of a manifold  $V$ .

Our main theorem is the following

**Theorem 1.1.** *For a continuous map  $f: M^n \rightarrow N^{n+k}$  between closed topological manifolds,  $U_M(1 \times w_k(f)) + (f \times f)^* U_N = 0$  if and only if  $f^* f_!(a) = a w_k(f)$  for all  $a \in H^*(M)$ .*

The cohomology elements, appearing in this theorem, are related to the embeddability of  $f$ . A. Haefliger [7, Théorème 5.2] proved the following

**Theorem** (Haefliger). *If a map  $f: M^n \rightarrow N^{n+k}$  between topological manifolds is homotopic to a topological embedding, then  $w_i(f) = 0$  for  $i > k$  and*

$$U_M(1 \times w_k(f)) + (f \times f)^* U_N = 0 \in H^{n+k}(M \times M).$$

Thus we have immediately the following

**Corollary 1.2.** *If a map  $f: M^n \rightarrow N^{n+k}$  between closed topological manifolds is homotopic to a topological embedding, then  $f^* f_!(a) = aw_k(f)$  for all  $a \in H^*(M)$ .*

REMARK 1. It is well-known, e.g., [4, p. 246], that if  $f$  is homotopic to a differentiable embedding then  $f^* f_!(a) = aw_k(f)$  for all  $a \in H^*(M)$ .

REMARK 2. As we will see in §3, the assumption ‘homotopic’ in Haefliger’s theorem or Corollary 1.2 can be weakened to ‘ $R$ -bordant’.

R.L.W. Brown [4] established the conditions that a map  $f: M^n \rightarrow N^{n+k}$  is cobordant to a differentiable embedding in the sense of Stong [12]. Here a map  $f_1: M_1^n \rightarrow N_1^{n+k}$  between differentiable closed manifolds is said to be cobordant to  $f_2: M_2^n \rightarrow N_2^{n+k}$  if there exist two cobordisms  $(W, M_1^n, M_2^n)$ ,  $(V, N_1^{n+k}, N_2^{n+k})$  and a map  $F: W \rightarrow V$  such that  $F|_{M_i} = f_i$  ( $i = 1, 2$ ).

From Theorem 1.1 and Brown’s theorem [4], we infer immediately a result which means the converse of Haefliger’s theorem up to cobordism of maps in the sense of Stong [12].

**Corollary 1.3.** *Let  $k > 0$ . Then a map  $f: M^n \rightarrow N^{n+k}$  between differentiable manifolds is cobordant to a differentiable embedding if  $w_i(f) = 0$  ( $i > k$ ) and  $U_M(1 \times w_k(f)) + (f \times f)^* U_N = 0$ .*

For an  $n$ -manifold  $M$ , we use the same symbol  $M$  as the generator of  $H^n(M) \cong \mathbf{Z}_2$ , i.e.,  $H^n(M) = \mathbf{Z}_2\langle M \rangle$ , and denote the  $H^p(M) \times H^q(M)$ -component of  $u \in H^{p+q}(M \times M)$  by  $[u]_{p,q}$ . To prove Theorem 1.1 we use the following

**Proposition 1.4.** *For a map  $f: M^n \rightarrow N^{n+k}$  and two elements  $x, y \in H^*(M)$  with  $\dim x + \dim y = r \leq n - k$ ,*

$$[(U_M(1 \times w_k(f)) + (f \times f)^* U_N)(x \times y)]_{n,k+r} = M \times (xw_k(f) + f^* f_!(x))y.$$

Using this proposition, we can reformulate Brown’s theorem [4] in case  $k > n/2$ .

**Theorem 1.5.** *Let  $k > n/2$ . Then a differentiable map  $f: M^n \rightarrow N^{n+k}$  is cobordant to a differentiable embedding if and only if the following two conditions hold:*

- (1)  $\langle w_I(M)w_J(f)w_i(f), [M] \rangle = 0$  for any integer  $i (i > k)$  and sequences  $I, J$  of non-negative integers such that  $|I| + |J| + i = n$ .
- (2)  $(U_M(1 \times w_k(f)) + (f \times f)^*U_N)(w_I(M) \times f^*(w_J(N))w_K(M)) = 0$  for any sequences  $I, J, K$  of non-negative integers such that  $|I| + |J| + |K| = n - k$ .

Here,  $w_I(M) = w_{i_1}(M) \cdots w_{i_r}(M)$  and  $|I| = \sum_{1 \leq j \leq r} i_j$  for a finite sequence  $I = (i_1, \dots, i_r)$  of non-negative integers.

The rest of this article is organized as follows: In §2, we will prove Theorem 1.1, Proposition 1.4 and Theorem 1.5. §3 will be devoted to the study of the relation between  $R$ -bordism and Haefliger's obstruction. In §4, we will give some examples of maps  $f: M^n \rightarrow N^{n+k}$ , e.g., a map which is cobordant to a differentiable embedding but not  $R$ -bordant to a topological embedding.

## 2. Proofs

To prove Theorem 1.1 and Proposition 1.4, we use the following two lemmas, the first of which is a slight generalization of [8, Lemma 2].

**Lemma 2.1.** *For a map  $f: M^n \rightarrow N^{n+k}$  and an element  $x \in H^r(M)$ , we have*

$$[(f \times f)^*U_N(x \times 1)]_{n,k+r} = M \times f^*f_1(x).$$

*Proof.* We can choose bases  $\{u_i \mid i \in I\}$  and  $\{v_i \mid i \in I\}$  for  $H^*(N)$  such that  $\langle u_i v_j, [N] \rangle = \delta_{ij}$ . Then the Thom class  $U_N$  of  $N$  can be described as  $U_N = \sum_{i \in I} u_i \times v_i$  by, e.g., [10, Theorem 11.11]. The element  $f_1(x)$  can be described as  $f_1(x) = \sum_{i \in I} \alpha_i v_i (\alpha_i \in \mathbf{Z}_2)$ . Let  $I_0 = \{i \in I \mid f^*(u_i)x = M\}$ . Then

$$\begin{aligned} \alpha_i &= \langle \alpha_i u_i v_i, [N] \rangle = \left\langle u_i \sum_{i \in I} \alpha_i v_i, [N] \right\rangle = \langle u_i f_1(x), [N] \rangle \\ &= \langle u_i, f_1(x) \cap [N] \rangle = \langle u_i, f_*(x \cap [M]) \rangle = \langle f^*(u_i)x, [M] \rangle \\ &= \begin{cases} 1 & i \in I_0 \\ 0 & i \notin I_0. \end{cases} \end{aligned}$$

Thus,  $f_1(x) = \sum_{i \in I_0} v_i$  and so  $f^*f_1(x) = \sum_{i \in I_0} f^*(v_i)$ . Hence, we have

$$\begin{aligned} [(f \times f)^*U_N \cdot (x \times 1)]_{n,k+r} &= \left[ \left( \sum_{i \in I} f^*(u_i) \times f^*(v_i) \right) (x \times 1) \right]_{n,k+r} \\ &= \left[ \sum_{i \in I} f^*(u_i)x \times f^*(v_i) \right]_{n,k+r} \end{aligned}$$

$$\begin{aligned}
&= \sum_{i \in I_0} M \times f^*(v_i) = M \times \sum_{i \in I_0} f^*(v_i) \\
&= M \times f^* f_1(x).
\end{aligned}$$

This completes the proof.  $\square$

**Lemma 2.2.** *For an  $n$ -manifold  $M^n$ , and an element  $x \in H^r(M)$ , we have*

$$[U_M(x \times 1)]_{n,r} = M \times x.$$

Proof. The Thom class  $U_M$  can be described as  $U_M = M \times 1 + \sum_j a_j \times b_j$ , ( $\dim a_j < n$ ) and there is a relation  $U_M(x \times 1) = U_M(1 \times x)$  (e.g., [10, Lemma 11.8]). Thus the lemma follows immediately.  $\square$

Proof of Proposition 1.4. Let  $x, y \in H^*(M)$  with  $\dim x + \dim y = r$ . Then, we have

$$\begin{aligned}
&[(U_M(1 \times w_k(f)) + (f \times f)^* U_N)(x \times y)]_{n,k+r} \\
&= [U_M(1 \times w_k(f))(x \times 1) + (f \times f)^* U_N(x \times 1)]_{n,k+\dim x}(1 \times y) \\
&= M \times x w_k(f) y + M \times f^* f_1(x) y \quad \text{by Lemmas 2.1–2.2} \\
&= M \times (x w_k(f) + f^* f_1(x)) y.
\end{aligned}$$

Thus, the proposition follows.  $\square$

Proof of Theorem 1.1. First we assume that  $U_M(1 \times w_k(f)) + (f \times f)^* U_N = 0$ . Take any  $a \in H^r(M)$ . Then

$$\begin{aligned}
0 &= [(U_M(1 \times w_k(f)) + (f \times f)^* U_N)(a \times 1)]_{n,k+r} \\
&= M \times (a w_k(f) + f^* f_1(a)) \quad \text{by Proposition 1.4}
\end{aligned}$$

Thus we get  $f^* f_1(a) = a w_k(f)$  for all  $a \in H^*(M)$ .

Conversely, suppose that  $f^* f_1(a) = a w_k(f)$  for all  $a \in H^*(M)$ . Since  $U_M(1 \times w_k(f)) + (f \times f)^* U_N \in H^{n+k}(M \times M)$ , it is sufficient for our purpose to show that  $(U_M(1 \times w_k(f)) + (f \times f)^* U_N)u = 0$  for all  $u \in H^{n-k}(M \times M)$ . By the Künneth formula, we may assume that  $u = a \times b$  with  $\dim a + \dim b = n - k$ . Then by Proposition 1.4, we have

$$\begin{aligned}
&(U_M(1 \times w_k(f)) + (f \times f)^* U_N)(a \times b) \\
&= [(U_M(1 \times w_k(f)) + (f \times f)^* U_N)(a \times b)]_{n,n} \\
&= M \times (a w_k(f) + f^* f_1(a)) b = 0.
\end{aligned}$$

Hence we get  $U_M(1 \times w_k(f)) + (f \times f)^* U_N = 0$ .  $\square$

Proof of Theorem 1.5. The condition (1) of Theorem 1.5 is just a restatement of the condition (i) of Brown's theorem. On the other hand, by the assumption that  $k > n/2$ , we have only to consider the case  $r = 2$  in the condition (ii) of Brown's theorem, which is reduced to

$$\langle f^*(w_J(N))f^*f_i(w_I(M))w_K(M), [M] \rangle = \langle f^*(w_J(N))w_I(M)w_K(M)w_k(f), [M] \rangle.$$

Applying Proposition 1.4 for  $x = w_I(M)$  and  $y = f^*(w_J(N))w_K(M)$ , we see that this equality is equivalent to the condition (2) of Theorem 1.5.  $\square$

### 3. Relations between $R$ -bordisms and Haefliger's obstructions

The concept of  $R$ -bordism of maps is introduced in [3, §3]. Let  $f_i: M_i^n \rightarrow N^{n+k}$  ( $i = 1, 2$ ) be maps between topological manifolds, where  $M_i$ 's are closed (while  $N$  is not necessarily closed). The two maps are said to be  $R$ -bordant if there exist a topological cobordism  $(W, M_1, M_2)$  and a continuous map  $F: W \rightarrow N$  such that (1)  $F|_{M_i} = f_i$  ( $i = 1, 2$ ) and (2) there exist retractions  $r_i: W \rightarrow M_i$  ( $i = 1, 2$ ).

Let  $j_i: M_i \rightarrow W$  be the natural inclusion ( $i = 1, 2$ ). Then by [6, Theorem 1.2],

$$(r_2j_1)_*: H_*(M_1) \rightarrow H_*(M_2)$$

is an isomorphism, and by [3, §3]

$$f_{1*} = f_{2*}(r_2j_1)_*: H_*(M_1) \rightarrow H_*(N).$$

In this section, we will prove

**Theorem 3.1.** *Let  $f: M^n \rightarrow N^{n+k}$  be a map between closed topological manifolds. If  $f$  is  $R$ -bordant to a topological embedding, then  $w_i(f) = 0$  ( $i > k$ ) and*

$$U_M(1 \times w_k(f)) + (f \times f)^*U_N = 0.$$

This theorem, together with Corollary 1.3, leads to the following

**Corollary 3.2.** *Let  $f: M^n \rightarrow N^{n+k}$  be a map between closed differentiable manifolds. If  $f$  is  $R$ -bordant to a topological embedding, then  $f$  is cobordant to a differentiable embedding.*

REMARK 3. If we consider *cobordism* and *embeddings* in topological category, the conclusion of this corollary is rather trivial.

Theorem 3.1 follows from Proposition 3.3 (or Corollary 3.4) below and Haefliger's theorem.

**Proposition 3.3.** *Let  $f_i: M_i^n \rightarrow N^{n+k}$  ( $i = 1, 2$ ) and  $g: M_1^n \rightarrow M_2^n$  be maps such that  $g_*: H_*(M_1) \rightarrow H_*(M_2)$  is an isomorphism and  $f_{1*} = f_{2*}g_*: H_*(M_1) \rightarrow H_*(N)$ . Then  $w(f_1) = g^*w(f_2)$  and*

$$\begin{aligned} U_{M_1}(1 \times w_k(f_1)) + (f_1 \times f_1)^*U_N \\ = (g \times g)^*(U_{M_2}(1 \times w_k(f_2)) + (f_2 \times f_2)^*U_N). \end{aligned}$$

*Proof.* Let  $\{u_i \mid i \in I\}$  and  $\{v_i \mid i \in I\}$  be two bases for  $H^*(M_2)$  such that  $\langle u_i v_j, [M_2] \rangle = \delta_{ij}$ . Then the Thom class  $U_{M_2}$  of  $M_2$  can be described as  $U_{M_2} = \sum_{i \in I} u_i \times v_i$  (see [10, Theorem 11.11]). Since  $g_*[M_1] = [M_2]$  and  $g^*$  is an isomorphism, because so is  $g_*$ , we have the two bases  $\{g^*u_i \mid i \in I\}$  and  $\{g^*v_i \mid i \in I\}$  for  $H^*(M_1)$  with  $\langle (g^*u_i)(g^*v_j), [M_1] \rangle = \delta_{ij}$ . Hence,

$$U_{M_1} = \sum_{i \in I} g^*u_i \times g^*v_i = (g \times g)^* \sum_{i \in I} u_i \times v_i = (g \times g)^*U_{M_2}.$$

Since  $f_{1*} = f_{2*}g_*$ , we have  $f_1^* = g^*f_2^*$  and  $w(f_1) = g^*w(f_2)$  by [3, Theorem 4.2]. Hence we have

$$\begin{aligned} U_{M_1}(1 \times w_k(f_1)) + (f_1 \times f_1)^*U_N \\ = (g \times g)^*U_{M_2}(1 \times g^*w_k(f_2)) + (g \times g)^*(f_2 \times f_2)^*U_N \\ = (g \times g)^*(U_{M_2}(1 \times w_k(f_2)) + (f_2 \times f_2)^*U_N). \end{aligned}$$

This completes the proof.  $\square$

**Corollary 3.4.** *Let  $f_i: M_i^n \rightarrow N^{n+k}$  ( $i = 1, 2$ ) be maps between closed topological manifolds. If  $f_1$  is  $R$ -bordant to  $f_2$ , then,  $w_i(f_1)$  ( $i \geq 0$ ) and  $U_{M_1}(1 \times w_k(f_1)) + (f_1 \times f_1)^*U_N$  correspond to  $w_i(f_2)$  ( $i \geq 0$ ) and  $U_{M_2}(1 \times w_k(f_2)) + (f_2 \times f_2)^*U_N$ , respectively, by the canonical isomorphisms.*

**REMARK 4.** By virtue of Proposition 1.4 and the fact that for  $f: M^n \rightarrow N^{n+k}$ ,  $w_k(f) + f^*f_!(1)$  is the Poincaré dual to the element  $\theta(f) \in H_{n-k}(M)$  in [3], the results in Theorem 3.1, Proposition 3.3 and Corollary 3.4 are, respectively, somewhat stronger than those in [3, Corollary 4.4, Theorem 4.2 and Corollary 4.3] in case  $N$  is a closed manifold.

#### 4. Relations among obstructions to embeddings

For a map  $f: M^n \rightarrow N^{n+k}$ , we describe conditions (0)–(3) below:

- (0)  $w_i(f) = 0$  for  $i > k$ .
  - (1)  $f^*f_!(a) + aw_k(f) = 0$  for all  $a \in H^*(M)$ .
- (or equivalently,  $U_M(1 \times w_k(f)) + (f \times f)^*U_N = 0$  by Theorem 1.1.)

(2)  $f^* f_!(w_I(M)) + w_I(M)w_k(f) = 0$  for all sequences  $I$  of non-negative integers, where  $w_I(M) = w_{i_1}(M) \cdots w_{i_r}(M)$  if  $I = (i_1, \dots, i_r)$ .

(3)  $f^* f_!(1) + w_k(f) = 0$ .

So far, for a map  $f: M^n \rightarrow N^{n+k}$  between closed differentiable manifolds, we know

$$\begin{array}{l}
 f \text{ is homotopic to a topological embedding} \\
 \Downarrow \\
 f \text{ is } R\text{-bordant to a topological embedding} \Rightarrow (0) + (1) \\
 \Downarrow \\
 f \text{ is cobordant to a differentiable embedding} \Leftarrow (0) + (2) \\
 \Downarrow \\
 f \text{ is cobordant to a differentiable embedding} \Leftrightarrow (0) + (3)
 \end{array}$$

REMARK 5. If  $k \geq n - 4$ ,  $2k > n$  and if  $f$  satisfies the conditions (0) and (3), then  $f$  is cobordant to a differentiable embedding ([1, Theorems (3.6) and (3.9)] and [9, Corollary 1.3]).

REMARK 6. Even if  $f$  is cobordant to an embedding, the conditions (0) and (3) do not necessarily hold ([8, Remark 2]).

In this section, we will show that

- (a) even if  $f$  is  $R$ -bordant to an embedding,  $f$  is not necessarily homotopic to an embedding (see Example 1 below),
- (b) the conditions (0) and (2) do not imply the conditions (1) (see Example 2),
- (c) the condition (3) does not lead to the condition (2) (see Example 3), and
- (d) the conditions (0) and (3) induce the relation (see Proposition 4.1)

$$f^* f_!(v_i(M)) = v_i(M)w_k(f),$$

where  $v_i(M)$  stands for the  $i$ -th Wu class of  $M$  defined by  $Sq(\sum_{0 \leq i} v_i(M)) = w(M)$ .

EXAMPLE 1. Let  $S^1 = \{z \in \mathbf{C}^1 \mid |z| = 1\}$  be the circle, and let  $f: S^1 \rightarrow S^1 \times S^1$  be a map defined by  $f(z) = (f_1(z), f_2(z)) = (z^2, 1)$ . Then  $f$  is not homotopic to an embedding. But  $f$  is  $R$ -bordant to an embedding.

REMARK 7. This example is a modification of an example appearing in earlier versions of [3], but omitted in the final one.

Proof. Suppose that  $f$  is homotopic to a topological embedding  $g = (g_1, g_2): S^1 \rightarrow S^1 \times S^1$ . Then  $g_2$  is homotopic to the constant map  $f_2$ . Hence,  $g_2$  has a lifting  $g'_2: S^1 \rightarrow \mathbf{R}^1$ . If we put  $g' = (g_1, g'_2): S^1 \rightarrow S^1 \times \mathbf{R}^1$ , then  $g'$  is also an embedding.

Identifying  $S^1 \times \mathbf{R}^1$  with  $\mathbf{C}^1 - \{0\}$ , we have a topological embedding  $g': S^1 \rightarrow \mathbf{C}^1 - \{0\}$ . From now on, the authors owe C. Biasi, J. Daccach and O. Saeki for the proof. Note that  $g'_*: H_1(S^1, \mathbf{Z})(\cong \mathbf{Z}) \rightarrow H_1(\mathbf{C}^1 - \{0\}, \mathbf{Z})(\cong \mathbf{Z})$  maps  $a \in \mathbf{Z}$  to  $2a$ . By the Schoenflies theorem,  $g'(S^1)$  bounds a region  $U$  in  $\mathbf{C}^1$  homeomorphic to the closed 2-dimensional disk. If  $0 \notin U$ , then  $g'$  is null-homotopic in  $\mathbf{C}^1 - \{0\}$ , which is a contradiction. If  $0 \in U$ , then  $g'$  represents a generator of  $H_1(\mathbf{C}^1 - \{0\})$ , which is also a contradiction. Thus  $f$  is not homotopic to an embedding. On the other hand,  $f$  is  $R$ -bordant to an embedding by [3, Example 4.8].  $\square$

EXAMPLE 2. We denote by  $P^m$  the real projective  $m$ -space. Furthermore,  $\pi: P^3 \rightarrow P^3/P^2 = S^3$  and  $j: P^l \subset P^{l+k}$  stand for the natural projection and inclusion, respectively. Let  $M^n = P^3 \times P^l$ ,  $N = S^3 \times P^{l+k}$  and let  $f = \pi \times j: M^n \rightarrow N^{n+k}$ . Then  $f$  satisfies (0) and (2), but  $f$  does not satisfy (1).

Proof. Put

$$H^1(P^3) = \mathbf{Z}_2\langle x_1 \rangle, \quad H^1(P^l) = \mathbf{Z}_2\langle x_2 \rangle, \quad H^3(S^3) = \mathbf{Z}_2\langle s \rangle, \quad H^1(P^{l+k}) = \mathbf{Z}_2\langle y \rangle.$$

Then

$$f^*(s) = x_1^3, \quad f^*(y) = x_2, \quad w(f) = (1 + x_2)^{-l-1}(1 + x_2)^{l+k+1} = (1 + x_2)^k.$$

Therefore

$$w_i(f) = 0 \quad \text{for } i > k, \quad w_k(f) = x_2^k.$$

The Thom classes of  $M$  and  $N$  are given by

$$\begin{aligned} U_M &= \sum_{0 \leq i \leq l} x_1^3 x_2^i \times x_2^{l-i} + \sum_{0 \leq i \leq l} x_1^2 x_2^i \times x_1 x_2^{l-i} \\ &\quad + \sum_{0 \leq i \leq l} x_1 x_2^i \times x_1^2 x_2^{l-i} + \sum_{0 \leq i \leq l} x_2^i \times x_1^3 x_2^{l-i}, \\ U_N &= \sum_{0 \leq i \leq l+k} s y^i \times y^{l+k-i} + \sum_{0 \leq i \leq l+k} y^i \times s y^{l+k-i}. \end{aligned}$$

Hence, and because  $f^*(y^{l+1}) = x_2^{l+1} = 0$ , we have

$$[U_M(1 \times w_k(f)) + (f \times f)^* U_N]_{n,k} = M \times (w_k(f) + f^* f_l(1)) = 0,$$

$$[U_M(1 \times w_k(f)) + (f \times f)^* U_N]_{n-1,k+1} = x_1^2 x_2^l \times x_1 x_2^k,$$

$$M \times f^* f_l(x_2^i) = [((f \times f)^* U_N)(x_2^i \times 1)]_{n,k+i} = M \times x_2^{k+i}.$$



Thus  $f$  does not satisfy the condition (1). But  $f$  satisfies (2), because  $w_i(M) = \binom{l+1}{i} x_2^i$  and  $f^* f_i(x_2^r) = x_2^{r+k} = x_2^r w_k(f)$ .  $\square$

REMARK 8. The above example shows that a map  $f$  satisfying the conditions (0) and (2) is not necessarily  $R$ -bordant to an embedding, in particular that a map which is cobordant to a differentiable embedding is not necessarily  $R$ -bordant to a topological embedding.

EXAMPLE 3. Let  $\pi: P^2 \rightarrow P^2/P^1 = S^2$  and  $j: P^l \subset P^{l+k}$  be the natural projection and inclusion, respectively and let  $f = \pi \times j: M = P^2 \times P^l \rightarrow S^2 \times P^{l+k}$ . Then, if  $k$  is even, the relation  $f^* f_i(1) = w_k(f)$  holds, however (2) does not hold.

Proof. As in Example 2, put

$$H^1(P^2) = \mathbf{Z}_2\langle x_1 \rangle, \quad H^1(P^l) = \mathbf{Z}_2\langle x_2 \rangle, \quad H^2(S^2) = \mathbf{Z}_2\langle s \rangle, \quad H^1(P^{l+k}) = \mathbf{Z}_2\langle y \rangle.$$

Then

$$w_1(M) = x_1 + (l+1)x_2, \quad f^*(s) = x_1^2, \quad f^*(y) = x_2, \quad w_k(f) = x_2^k.$$

Just as in Example 2, we have

$$M \times (w_k(f) + f^* f_i(1)) = [U_M(1 \times w_k(f)) + (f \times f)^* U_N]_{n,k} = 0,$$

$$\begin{aligned} M \times (w_1(M)w_k(f) + f^* f_i(w_1(M))) \\ &= [(U_M(1 \times w_k(f)) + (f \times f)^* U_N)(w_1(M) \times 1)]_{n,k+1} \\ &= M \times x_1 x_2^k. \end{aligned}$$

Thus the relation  $f^* f_i(1) = w_k(f)$  holds, however  $f^* f_i(w_1(M)) \neq w_1(M)w_k(f)$ .  $\square$

**Proposition 4.1.** *Assume that  $f: M^n \rightarrow N^{n+k}$  satisfies the conditions that  $w_i(f) = 0$  ( $k < i$ ) and  $f^* f_i(1) = w_k(f)$ , then*

$$f^* f_i(v_i(M)) = v_i(M)w_k(f) \quad (0 < i).$$

Proof. For each  $x \in H^{n-k-i}(M)$ , we have

$$\begin{aligned} x f^* f_i(v_i(M)) &= v_i(M) f^* f_i(x) \quad \text{by, e.g., [9, Lemma 2.1, (4)]} \\ &= Sq^i f^* f_i(x) \quad \text{because } \dim f^* f_i(x) = n - i \\ &= [Sq f^* f_i(x)]_n \end{aligned}$$

$$\begin{aligned}
&= [f^* f_i(Sq(x)w(f))]_n \quad \text{by, e.g., [9, Lemma 2.1, (2)]} \\
&= f^* f_i \left( \sum_{0 \leq j} Sq^j(x)w_{i-j}(f) \right) \\
&= \sum_{0 \leq j} Sq^j(x)w_{i-j}(f)f^* f_i(1) \quad \text{by, e.g., [9, Lemma 2.1, (4)]} \\
&= \sum_{0 \leq j} Sq^j(x)w_{i-j}(f)w_k(f) \quad \text{because } f^* f_i(1) = w_k(f) \\
&= \sum_{0 \leq j} Sq^j(x)Sq^{i-j}w_k(f) \quad \text{because } w_i(f) = 0 \ (k < i) \\
&= Sq^i(xw_k(f)) = v_i(M)xw_k(f).
\end{aligned}$$

Here,  $[y]_j$  for  $y \in \sum_{0 \leq i} H^i(M)$  means the  $j$ -dimensional component of  $y$ . Thus  $xf^* f_i(v_i(M)) = xv_i(M)w_k(f)$  for all  $x \in H^{n-k-i}(M)$ . Hence  $f^* f_i(v_i(M)) = v_i(M)w_k(f)$  by the Poincaré duality.  $\square$

For  $k = 1$ , the conditions (0) and (3) imply the condition (2), i.e. we have

**Proposition 4.2.** *Assume that  $f: M^n \rightarrow N^{n+1}$  satisfies the conditions that  $w_i(f) = 0$  ( $1 < i$ ) and  $f^* f_i(1) = w_1(f)$ , then for all sequences  $I$  of non-negative integers, we have*

$$f^* f_i(w_I(M)) = w_I(M)w_1(f).$$

*Proof.* By the assumption we have  $\bar{w}(M)f^*w(N) = w(f) = 1 + w_1(f) = 1 + f^* f_i(1)$ . Hence  $w(M) = f^*w(N)(1 + f^* f_i(1))^{-1} = f^*(w(N)(1 + f_i(1))^{-1}) \in f^*H^*(N)$ . Thus  $w_I(M) \in f^*H^*(N)$  for all  $I$ , and therefore we obtain the result since  $f_i(f^*y) = yf_i(1)$  for all  $y \in H^*(N)$ .  $\square$

---

#### References

- [1] M.A. Aguilar and G. Pastor: *On maps cobordant to embeddings*, Bol. Soc. Mat. Mexicana **31** (1986), 61–67.
- [2] M.F. Atiyah and F. Hirzebruch: *Cohomologie-operationen und charakteristische klassen*, Math. Z. **77** (1961), 149–187.
- [3] C. Biasi, J. Daccach and O. Saeki: *On R-Bordism of maps and obstruction to topological embeddings*, Osaka J. Math. **37** (2000), 527–535.
- [4] R.L. W. Brown: *Stiefel-Whitney numbers and maps cobordant to embeddings*, Proc. Amer. Math. Soc. **48** (1975), 245–250.
- [5] E. Dyer: *Cohomology Theories*, Benjamin, New York, 1969.
- [6] D.R. Gibbs: *Cobordism with retractions*, Math. Ann. **203** (1973), 77–88.

- [7] A. Haefliger: *Points multiples d'une application et produit cyclique reduit*, Amer. J. Math. **83** (1961), 57–70.
- [8] Y. Kuramoto, T. Murata and T. Yasui: *Non-embeddability of manifolds to projective spaces*, Math. J. Toyama Univ. **23** (2000), 79–83.
- [9] Y. Kuramoto and T. Yasui: *On conditions that a map is cobordant to an embedding*, Far East J. Math. Sci. Special vol. (2000), Part II, 189–206.
- [10] J.W. Milnor and J. D. Stasheff: *Characteristic Classes*, Ann. of Math. Stud. **76**, Princeton Univ. Press, 1974.
- [11] E.H. Spanier: *Algebraic Topology*, McGraw-Hill, New York, 1966.
- [12] R.E. Stong: *Cobordism of maps*, Topology **5** (1966), 245–258.

Yoshiyuki Kuramoto  
Faculty of Education  
Kagoshima University  
Kagoshima 890-0065, Japan  
e-mail: kuramoto@edu.kagoshima-u.ac.jp

Tsutomu Yasui  
Faculty of Education  
Kagoshima University  
Kagoshima 890-0065, Japan  
e-mail: yasui@edu.kagoshima-u.ac.jp